

Home Assignment 4

Jaan Tollander de Balsch

December 5, 2018

Question 1

A kite is a graph on an even number of vertices, say $2n$, in which n of the vertices form a clique and the remaining n vertices are connected in a *tail* that consists of a path joined to one of the vertices of the clique. Given a graph and a goal g , the kite problem asks for a subgraph which is a kite and which contains $2g$ vertices. Prove that kite is NP-complete.

Question 2

Consider two problems:

- 1) Given a graph $G = (V, E)$ and integer k , find an independent set of size at least k .
- 2) Given a graph $G = (V, E)$, compute the size of maximum independent set.
- 3) Given a graph $G = (V, E)$ and integer k , decide whether graph G has an independent set of size k .

Prove that these three problems are equally hard. That is, show that if one of them admits an efficient algorithm, then all of them do.

Question 3

In the EXACT 4SAT problem, the input is a set of clauses, each of which is a disjunction of exactly four literals, and such that each variable occurs at most once in each clause. The goal is to find a satisfying assignment, if one exists. Prove that EXACT 4SAT is NP-complete.

Question 4

Consider the following two problems:

- 1) **Problem X:** Given an undirected graph $G = (V, E)$ and integer k , find a cut $(C, V - S)$ such that there are at least k edges across the cut, i.e. $|\{(u, v) \in E : u \in S, v \in V - S\}| \geq k$.
- 2) **Problem Y:** Given a collection of m equations over n variables x_1, \dots, x_n under modulo 2:

$$\begin{aligned} x_1 + x_2 &= 1 \pmod{2} \\ x_2 + x_3 + x_4 &= 0 \pmod{2} \\ &\dots \\ x_i + x_j + x_l &= 1 \pmod{2} \end{aligned}$$

The goal is to decide whether there is an assignment $\sigma : \{x_1, x_2, \dots, x_n\} \rightarrow \{0, 1\}$ that satisfies at least k equations.

Part 1: Describe a reduction from Problem X to Problem Y. Since Problem X is NP-hard, this reduction implies that Problem Y is NP-hard.

Part 2: Consider a special case of problem Y where each equation involves at most 2 variables. Present an $O(n + m)$ time algorithm that decides whether all m equations can be simultaneously satisfied.

Does the existence of this algorithm contradict Part 1 where we just proved that Problem Y is NP-hard?

Question 5

This question is taken directly from Ericson's [chapter](#). Please use the definitions of universal, uniform and pairwise independent hash functions from his book.

Each of the following question is worth one point:

- a) Describe a set of hash functions that is uniform but not (near-)universal.
- b) Describe a set of hash functions that is universal but not (near)uniform.
- c) Describe a set of hash functions that is universal but (near-)3-universal.
- d) Describe a set of hash functions that is uniform but not pairwise independent.
- e) Describe a set of hash functions that is pairwise independent but not (near-)uniform.
- f) Describe a set of hash functions that is universal but not pairwise independent.
- g) Describe a set of hash functions that is pairwise independent but not (near-)uniform.
- h) Describe a set of hash functions that is universal and pairwise independent but not uniform, or prove no such set exists.

Note: For this particular question, an answer without proof will not receive any point.

Question 6

In a student chess competition, there are n students (we call these students by numbers $1, 2, 3, \dots, n$) who played one-on-one decisive matches against each other (i.e. each game only ends in win/loss but not a draw). There were m matches in total.

The competition is over. To encourage a more fierceful competition among students in the next year, an evil coach wants to group these students into two groups: He calls the groups, the elite and the sucker groups respectively. The coach wants to group the students in such a way that the grouping is as *consistent* with the competition results as possible. In particular, each match record of the form *i lost to j* is said to be consistent with the grouping if student *i* is placed in the sucker group and student *j* is placed in the elite group. The coach aims at maximizing the number of consistent match results.

Part 1: A randomized algorithm picks a partition into two groups at random. Prove that the expected number of consistent matches is at least $1/4$ of the total number of matches.

Part 2: Describe an efficient deterministic algorithm that partitions students into two groups such that the number of consistent matches is at least $1/4$ of the total number of matches.

Part 3: Now, instead of partitioning into only two groups, the coach wants to rank the students (no ties are allowed). A ranking is said to be consistent with the match *i lost to j* if the rank of student *i* is worse than the rank of student *j*. The objective of the coach is to produce a ranking that is as consistent as possible.

A randomized algorithm picks a random ranking, i.e. pick a random permutation σ and uses this as a ranking. Prove that the expected number of consistent match results is at least $1/2$ of the total number of matches.

Part 4: Describe a deterministic algorithm that, on an input, produces a ranking that is always consistent with at least $1/2$ of the total number of matches.