

MATRIX ALGEBRA

J. Alexander Branham

Fall 2016

WHAT IS A MATRIX?

- Let's start with something familiar, with a new word

- Let's start with something familiar, with a new word
- One number (12, for example) is referred to as a *scalar*

- Let's start with something familiar, with a new word
- One number (12, for example) is referred to as a *scalar*
- This can be thought of as a 1x1 matrix

- Let's start with something familiar, with a new word
- One number (12, for example) is referred to as a *scalar*
- This can be thought of as a 1x1 matrix
 - More on that in a bit...

$$\begin{bmatrix} 12 \end{bmatrix} = c$$

- We can put several scalars together to make a vector

- We can put several scalars together to make a vector
- For example:

$$\begin{bmatrix} 12 \\ 14 \\ 15 \end{bmatrix} = b$$

- We can put several scalars together to make a vector
- For example:

$$\begin{bmatrix} 12 \\ 14 \\ 15 \end{bmatrix} = b$$

- Since this is a column of numbers, we cleverly refer to it as a *column vector*

If we take b and arrange it so that it is a row of numbers instead of a column, we refer to it as a *row vector*:

$$\begin{bmatrix} 12 & 14 & 15 \end{bmatrix} = d$$

OPERATORS

OPERATIONS ON VECTORS:

The **summation operator** \sum lets us perform an operation (sum) on a sequence of numbers (often but not always a vector):

$$\sum_{i=1}^5 i$$

SUMMATION OPERATOR

Let:

$$x_i = \begin{bmatrix} 12 & 7 & -2 & 0 & 1 \end{bmatrix}$$

Find:

$$\sum_{i=1}^3 x_i$$

SUMMATION OPERATOR

```
x <- c(12, 7, -2, 0, 1)
```

```
sum(x[1:3])
```

```
## [1] 17
```

YOU TRY!

Let

$$y = \begin{bmatrix} 1 & 0 & -1 & 4 \end{bmatrix}$$

Find:

$$\sum_{i=1}^{\infty} y^2$$

YOU TRY (ANSWER)

```
y <- c(1, 0, -1, 4)
```

```
sum(y ^ 2)
```

```
## [1] 18
```


We might want to multiply instead of add, in which case we can use the product operator \prod

Let:

$$z = \begin{bmatrix} 6 & -2 & 0 & 1 \end{bmatrix}$$

Find:

$$\prod_{i=1}^2 z_i$$

```
z <- c(6, -2, 0, 1)
```

```
prod(z[1:2])
```

```
## [1] -12
```

YOU TRY!

Let:

$$a = [1 \ 2 \ 3 \ 4 \ 5 \ 4 \ 3 \ 2 \ 1]$$

Find:

$$\prod_{i=1}^3 a^2 - 2a + 3$$

YOU TRY (ANSWER)

```
a <- c(1, 2, 3, 4, 5, 4, 3, 2, 1)
prod(a[1:3] ^ 2 - 2 * a[1:3] + 3)
```

```
## [1] 36
```

MATRICES

We can put multiple vectors together to get a *matrix*:

$$\begin{bmatrix} 12 & 14 & 15 \\ 115 & 22 & 127 \\ 193 & 29 & 219 \end{bmatrix} = A$$

- We refer to the *dimensions* of matrices by *row x column*

- We refer to the *dimensions* of matrices by *row x column*
- So A is a 3×3 matrix.

- We refer to the *dimensions* of matrices by *row x column*
- So A is a 3×3 matrix.
- Note that matrices are usually designated by capital letters

- We refer to the *dimensions* of matrices by *row x column*
- So A is a 3×3 matrix.
- Note that matrices are usually designated by capital letters
 - And sometimes bolded as well

ROW x COLUMN

- How do we refer to specific elements of the matrix???

- How do we refer to specific elements of the matrix???
- Solution: come up with a clever indexing scheme

- How do we refer to specific elements of the matrix???
- Solution: come up with a clever indexing scheme
- Matrix A is an $n \times m$ matrix where $n = m = 3$.

- How do we refer to specific elements of the matrix???
- Solution: come up with a clever indexing scheme
- Matrix A is an $n \times m$ matrix where $n = m = 3$.
- More generally, matrix B is an $m \times n$ matrix where the elements look like this:

$$B = \begin{bmatrix} b_{11} & b_{12} & b_{13} & \dots & b_{1n} \\ b_{21} & b_{22} & b_{23} & \dots & b_{2n} \\ \vdots & \vdots & \vdots & \dots & \vdots \\ b_{m1} & b_{m2} & b_{m3} & \dots & b_{mn} \end{bmatrix}$$

MATRIX OPERATIONS

ADDITION AND SUBTRACTION ARE EASY!

- Requirement: Must have *exactly* the same dimensions

ADDITION AND SUBTRACTION ARE EASY!

- Requirement: Must have *exactly* the same dimensions
- To do the operation, just add or subtract each element with the corresponding element from the other matrix:

$$A \pm B$$

ADDITION AND SUBTRACTION

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \pm \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{bmatrix}$$
$$= \begin{bmatrix} a_{11} \pm b_{11} & a_{12} \pm b_{12} & a_{13} \pm b_{13} \\ a_{21} \pm b_{21} & a_{22} \pm b_{22} & a_{23} \pm b_{23} \\ a_{31} \pm b_{31} & a_{32} \pm b_{32} & a_{33} \pm b_{33} \end{bmatrix}$$

YOU TRY!

Let

$$A = \begin{bmatrix} 1 & 4 & 2 \\ -2 & -1 & 0 \\ 0 & -1 & 3 \end{bmatrix} \quad B = \begin{bmatrix} 5 & 1 & 0 \\ 2 & -1 & 0 \\ 7 & 1 & 2 \end{bmatrix}$$

Find $A + B$

YOU TRY (ANSWERS)

```
A <- matrix(c(1, 4, 2, -2, -1, 0, 0, -1, 3),  
            byrow = TRUE, nrow = 3)  
B <- matrix(c(5, 1, 0, 2, -1, 0, 7, 1, 2),  
            byrow = TRUE, nrow = 3)  
  
A + B
```

```
##      [,1] [,2] [,3]  
## [1,]    6    5    2  
## [2,]    0   -2    0  
## [3,]    7    0    5
```

Easy - just multiply each element of the matrix by the scalar

$$cA = \begin{bmatrix} ca_{11} & ca_{12} & ca_{13} \\ ca_{21} & ca_{22} & ca_{23} \\ ca_{31} & ca_{32} & ca_{33} \end{bmatrix}$$

SCALAR MULTIPLICATION

A

```
##      [,1] [,2] [,3]
## [1,]    1    4    2
## [2,]   -2   -1    0
## [3,]    0   -1    3
```

3 * A

```
##      [,1] [,2] [,3]
## [1,]    3   12    6
## [2,]   -6   -3    0
## [3,]    0   -3    9
```


- Requirement: the two matrices must be *conformable*

- Requirement: the two matrices must be *conformable*
- This means that the number of columns in the first matrix equals the number of rows in the second

- Requirement: the two matrices must be *conformable*
- This means that the number of columns in the first matrix equals the number of rows in the second
- The resulting matrix will have the number of rows in the first, and the number of columns in the second!

- Which can we multiply? What will the resulting dimensions be?

$$b = \begin{bmatrix} 2 \\ 3 \\ 4 \\ 1 \end{bmatrix} \quad M = \begin{bmatrix} 1 & 0 & 2 \\ 1 & 2 & 4 \\ 2 & 3 & 2 \end{bmatrix} \quad L = \begin{bmatrix} 6 & 5 & -1 \\ 1 & 4 & 3 \end{bmatrix}$$

POP QUIZ

- Which can we multiply? What will the resulting dimensions be?

$$b = \begin{bmatrix} 2 \\ 3 \\ 4 \\ 1 \end{bmatrix} \quad M = \begin{bmatrix} 1 & 0 & 2 \\ 1 & 2 & 4 \\ 2 & 3 & 2 \end{bmatrix} \quad L = \begin{bmatrix} 6 & 5 & -1 \\ 1 & 4 & 3 \end{bmatrix}$$

- **ONLY** LM and **NOT** ML

POP QUIZ

- Which can we multiply? What will the resulting dimensions be?

$$b = \begin{bmatrix} 2 \\ 3 \\ 4 \\ 1 \end{bmatrix} \quad M = \begin{bmatrix} 1 & 0 & 2 \\ 1 & 2 & 4 \\ 2 & 3 & 2 \end{bmatrix} \quad L = \begin{bmatrix} 6 & 5 & -1 \\ 1 & 4 & 3 \end{bmatrix}$$

- **ONLY** LM and **NOT** ML
- The dimensions will be 2×3

HOW TO ACTUALLY DO THIS?

- Multiply each row by each column

HOW TO ACTUALLY DO THIS?

- Multiply each row by each column
- (Board examples)

HOW TO ACTUALLY DO THIS?

- Multiply each row by each column
- (Board examples)

HOW TO ACTUALLY DO THIS?

- Multiply each row by each column
- (Board examples)

```
L <- matrix(c(6, 5, -1, 1, 4, 3),  
            nrow = 2, byrow = TRUE)  
M <- matrix(c(1, 0, 2, 1, 2, 4, 2, 3, 2),  
            nrow = 3, byrow = TRUE)  
  
L %*% M
```

```
##      [,1] [,2] [,3]  
## [1,]    9    7   30  
## [2,]   11   17   24
```


HAHAHA... NOPE

- Addition and subtraction

- Addition and subtraction
 - Associative $(A \pm B) \pm C = A \pm (B \pm C)$

- Addition and subtraction
 - Associative $(A \pm B) \pm C = A \pm (B \pm C)$
 - Commutative $A \pm B = B \pm A$

PROPERTIES OF OPERATORS

- Addition and subtraction
 - Associative $(A \pm B) \pm C = A \pm (B \pm C)$
 - Commutative $A \pm B = B \pm A$
- Multiplication

- Addition and subtraction
 - Associative $(A \pm B) \pm C = A \pm (B \pm C)$
 - Commutative $A \pm B = B \pm A$
- Multiplication
 - $AB \neq BA$

- Addition and subtraction
 - Associative $(A \pm B) \pm C = A \pm (B \pm C)$
 - Commutative $A \pm B = B \pm A$
- Multiplication
 - $AB \neq BA$
 - $A(BC) = (AB)C$

- Addition and subtraction
 - Associative $(A \pm B) \pm C = A \pm (B \pm C)$
 - Commutative $A \pm B = B \pm A$
- Multiplication
 - $AB \neq BA$
 - $A(BC) = (AB)C$
 - $A(B + C) = AB + AC$

PROPERTIES OF OPERATORS

- Addition and subtraction
 - Associative $(A \pm B) \pm C = A \pm (B \pm C)$
 - Commutative $A \pm B = B \pm A$
- Multiplication
 - $AB \neq BA$
 - $A(BC) = (AB)C$
 - $A(B + C) = AB + AC$
 - $(A + B)C = AC + BC$

TRANSPOSITION

- Switch the rows and columns

QU'EST-CE QUE C'EST?

- Switch the rows and columns
- So a $n \times m$ matrix becomes $m \times n$

QU'EST-CE QUE C'EST?

- Switch the rows and columns
- So a $n \times m$ matrix becomes $m \times n$
- Typically denoted L' or L^T

TRANSPOSITION

L

```
##      [,1] [,2] [,3]
```

```
## [1,]    6    5   -1
```

```
## [2,]    1    4    3
```

$t(L)$

```
##      [,1] [,2]
```

```
## [1,]    6    1
```

```
## [2,]    5    4
```

```
## [3,]   -1    3
```

- Matrix is *always* conformable for multiplication with its transpose in both directions

- Matrix is *always* conformable for multiplication with its transpose in both directions
- $(A \pm B)' = A' \pm B'$

- Matrix is *always* conformable for multiplication with its transpose in both directions
- $(A \pm B)' = A' \pm B'$
- $A'' = A$

- Matrix is *always* conformable for multiplication with its transpose in both directions
- $(A \pm B)' = A' \pm B'$
- $A'' = A$
- $(AB)' = B'A'$

PROPERTIES OF TRANSPOSING

- Matrix is *always* conformable for multiplication with its transpose in both directions
- $(A \pm B)' = A' \pm B'$
- $A'' = A$
- $(AB)' = B'A'$
- $(cA)' = cA'$ where c is a scalar

PROPERTIES OF TRANSPOSING

- Matrix is *always* conformable for multiplication with its transpose in both directions
- $(A \pm B)' = A' \pm B'$
- $A'' = A$
- $(AB)' = B'A'$
- $(cA)' = cA'$ where c is a scalar
- AA' and $A'A$ will always result in a symmetric matrix

YOU TRY!

Let:

$$A = \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix} \quad B = \begin{bmatrix} 2 & 3 & 4 \\ -1 & 0 & 1 \end{bmatrix}$$

Find:

1. $A'A$

YOU TRY!

Let:

$$A = \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix} \quad B = \begin{bmatrix} 2 & 3 & 4 \\ -1 & 0 & 1 \end{bmatrix}$$

Find:

1. $A'A$
2. AB

YOU TRY!

Let:

$$A = \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix} \quad B = \begin{bmatrix} 2 & 3 & 4 \\ -1 & 0 & 1 \end{bmatrix}$$

Find:

1. $A'A$
2. AB
3. BA

YOU TRY (ANSWERS)

```
A <- matrix(c(2, 4, 3, 1), byrow = TRUE, nrow = 2)
B <- matrix(c(2, 3, 4, -1, 0, 1), byrow = TRUE, nrow = 2)
A %*% t(A)
```

```
##      [,1] [,2]
## [1,]   20  10
## [2,]   10  10
```

YOU TRY (ANSWERS)

```
A %*% B
```

```
##      [,1] [,2] [,3]  
## [1,]    0    6   12  
## [2,]    5    9   13
```

YOU TRY (ANSWERS)

```
B %*% A
```

```
## Error in B %*% A: non-conformable arguments
```

MATRIX INVERSE

- I kinda lied when I said there isn't matrix division

MATRIX INVERSES

- I kinda lied when I said there isn't matrix division
- We use matrix inverses all the time

MATRIX INVERSES

- I kinda lied when I said there isn't matrix division
- We use matrix inverses all the time
- If A is an $n \times n$ square matrix:

$$AB = BA = I_n$$

MATRIX INVERSES

- I kinda lied when I said there isn't matrix division
- We use matrix inverses all the time
- If A is an $n \times n$ square matrix:

$$AB = BA = I_n$$

- Then B is said to be the *inverse* of A

MATRIX INVERSES

- I kinda lied when I said there isn't matrix division
- We use matrix inverses all the time
- If A is an $n \times n$ square matrix:

$$AB = BA = I_n$$

- Then B is said to be the *inverse* of A
 - This is usually denoted A^{-1}

MATRIX INVERSES

- I kinda lied when I said there isn't matrix division
- We use matrix inverses all the time
- If A is an $n \times n$ square matrix:

$$AB = BA = I_n$$

- Then B is said to be the *inverse* of A
 - This is usually denoted A^{-1}
 - So $AA^{-1} = I_n$

MATRIX INVERSES

- I kinda lied when I said there isn't matrix division
- We use matrix inverses all the time
- If A is an $n \times n$ square matrix:

$$AB = BA = I_n$$

- Then B is said to be the *inverse* of A
 - This is usually denoted A^{-1}
 - So $AA^{-1} = I_n$
- If B doesn't exist, then the matrix is *singular*

MATRIX INVERSES

- I kinda lied when I said there isn't matrix division
- We use matrix inverses all the time
- If A is an $n \times n$ square matrix:

$$AB = BA = I_n$$

- Then B is said to be the *inverse* of A
 - This is usually denoted A^{-1}
 - So $AA^{-1} = I_n$
- If B doesn't exist, then the matrix is *singular*
- Finding inverses by hand is super hard (especially as n increases), so we let computers do this for us

SOME PROPERTIES

- Let A be $n \times n$ square matrix:

SOME PROPERTIES

- Let A be $n \times n$ square matrix:
- If A^{-1} exists:

SOME PROPERTIES

- Let A be $n \times n$ square matrix:
- If A^{-1} exists:
- A is full rank: $\text{rank}(A) = n$

SOME PROPERTIES

- Let A be $n \times n$ square matrix:
- If A^{-1} exists:
- A is full rank: $\text{rank}(A) = n$
- A' is also invertible

SOME PROPERTIES

- Let A be $n \times n$ square matrix:
- If A^{-1} exists:
- A is full rank: $\text{rank}(A) = n$
- A' is also invertible
- $(A^{-1})^{-1} = A$

SOME PROPERTIES

- Let A be $n \times n$ square matrix:
- If A^{-1} exists:
- A is full rank: $\text{rank}(A) = n$
- A' is also invertible
- $(A^{-1})^{-1} = A$
- $(cA)^{-1} = c^{-1}A^{-1}$ for nonzero scalar c

SOME PROPERTIES

- Let A be $n \times n$ square matrix:
- If A^{-1} exists:
- A is full rank: $\text{rank}(A) = n$
- A' is also invertible
- $(A^{-1})^{-1} = A$
- $(cA)^{-1} = c^{-1}A^{-1}$ for nonzero scalar c
- $(A')^{-1} = (A^{-1})'$

SPECIAL MATRICES

Some matrices get more love than others

SQUARE MATRIX

Any $n \times n$ matrix (same number rows and columns)

$$\begin{bmatrix} 1 & 0 & 2 \\ 1 & 2 & 4 \\ 2 & 3 & 2 \end{bmatrix}$$

SYMMETRIC MATRIX

A square matrix that is the same as its transpose

$$\begin{bmatrix} 2 & 5 & 7 \\ 5 & 9 & 6 \\ 7 & 6 & 7 \end{bmatrix}$$

A symmetric matrix with zeros everywhere but the main diagonal

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & 9 & 0 \\ 0 & 0 & 7 \end{bmatrix}$$

A diagonal matrix with the same number all along the diagonal

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

IDENTITY MATRIX

A scalar matrix where the diagonal elements are 1.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

IDENTITY MATRIX

A scalar matrix where the diagonal elements are 1.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- This is a super important type of matrix.

IDENTITY MATRIX

A scalar matrix where the diagonal elements are 1.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- This is a super important type of matrix.
- It gets its own notation: I_n where n is the number of rows and columns

IDENTITY MATRIX

A scalar matrix where the diagonal elements are 1.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- This is a super important type of matrix.
- It gets its own notation: I_n where n is the number of rows and columns
- Note that $I_n A = A$ and also $A I_n = A$