FUNCTIONS

J. Alexander Branham

Fall 2016



• Exponents tell you to multiply that thing by its base x times:

• Exponents tell you to multiply that thing by its base x times:

$$\cdot 3^4 = 3 * 3 * 3 * 3 =$$

• Exponents tell you to multiply that thing by its base x times:

$$\cdot 3^4 = 3 * 3 * 3 * 3 =$$

 $\boldsymbol{\cdot}$ Logarithms ask how many times you must raise the base to get x:

- Exponents tell you to multiply that thing by its base x times:
 - $\cdot 3^4 = 3 * 3 * 3 * 3 =$
- Logarithms ask how many times you must raise the base to get x:
 - $log_3(81) = 4$

- Exponents tell you to multiply that thing by its base x times:
 - $\cdot 3^4 = 3 * 3 * 3 * 3 =$
- Logarithms ask how many times you must raise the base to get x:
 - $\cdot \log_3(81) = 4$
 - $\boldsymbol{\cdot}$ Note that logarithms with negative arguments are undefined

• Exponents tell you to multiply that thing by its base x times:

$$\cdot 3^4 = 3 * 3 * 3 * 3 =$$

- Logarithms ask how many times you must raise the base to get x:
 - $\cdot \log_3(81) = 4$
 - · Note that logarithms with negative arguments are undefined
 - Sometimes $\log(n)$ means $\log_{10}(n)$

• Exponents tell you to multiply that thing by its base x times:

$$\cdot 3^4 = 3 * 3 * 3 * 3 =$$

- Logarithms ask how many times you must raise the base to get x:
 - $\cdot \log_3(81) = 4$
 - · Note that logarithms with negative arguments are undefined
 - Sometimes log(n) means $log_{10}(n)$
 - · Othertimes, it means $\log_e(n) = ln(n)$

$$a^m a^n = a^{m+n}$$

$$a^m a^n = a^{m+n}$$

$$\frac{a^m}{a^n}=a^{m-n}$$

$$a^m a^n = a^{m+n}$$

$$\frac{a^m}{a^n}=a^{m-n}$$

$$(a^m)^n = a^{mn}$$

$$a^{m}a^{n} = a^{m+n}$$

$$\frac{a^{m}}{a^{n}} = a^{m-n}$$

$$(a^{m})^{n} = a^{mn}$$

 $a^0 = 1$

PROPERTIES OF EXPONENTS (CONTINUED)

$$a^{1/n} = \sqrt[n]{a}$$

PROPERTIES OF EXPONENTS (CONTINUED)

$$a^{1/n} = \sqrt[n]{a}$$

$$\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n} = a^n b^{-n} \qquad \forall b \neq 0$$

PROPERTIES OF EXPONENTS (CONTINUED)

$$a^{1/n} = \sqrt[n]{a}$$

$$\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n} = a^n b^{-n} \qquad \forall b \neq 0$$

$$\left(\frac{a}{b}\right)^{-n} = \left(\frac{b}{a}\right)^n \qquad \forall a, b \neq 0$$

$$\log_{x}(ab) = \log_{x} a + \log_{x} b$$

$$\log_{x}(ab) = \log_{x} a + \log_{x} b$$

$$\log_{\mathsf{x}}\left(\frac{a}{b}\right) = \log_{\mathsf{x}} a - \log_{\mathsf{x}} b$$

$$\log_{x}(ab) = \log_{x} a + \log_{x} b$$

$$\log_{x} \left(\frac{a}{b}\right) = \log_{x} a - \log_{x} b$$

$$\log_{x} a^{b} = b \log_{x} a$$

$$\log_{x}(ab) = \log_{x} a + \log_{x} b$$

$$\log_{x} \left(\frac{a}{b}\right) = \log_{x} a - \log_{x} b$$

$$\log_{x} a^{b} = b \log_{x} a$$

$$\log_{x} 1 = 0$$

$$\log_{x}(ab) = \log_{x} a + \log_{x} b$$

$$\log_{x} \left(\frac{a}{b}\right) = \log_{x} a - \log_{x} b$$

$$\log_{x} a^{b} = b \log_{x} a$$

$$\log_{x} 1 = 0$$

$$m^{\log_{m}(a)} = a$$

$$\log_{x}(ab) = \log_{x} a + \log_{x} b$$

$$\log_{x} \left(\frac{a}{b}\right) = \log_{x} a - \log_{x} b$$

$$\log_{x} a^{b} = b \log_{x} a$$

$$\log_{x} 1 = 0$$

$$m^{\log_{m}(a)} = a$$

$$\frac{\log_{x} n}{1 + \log_{m} n} = \log_{m} n$$

5

You try!

 2^{4} $\log(100)$ $a^{3} \times a^{7}$ $\log_{10}(10z)$

Functions


 \cdot Anything that takes input and gives one output

- · Anything that takes input and gives one output
- In math, this usually looks something like f(x,z) = y

- · Anything that takes input and gives one output
- In math, this usually looks something like f(x, z) = y
 - \cdot x and z are the arguments that the function takes

- Anything that takes input and gives one output
- In math, this usually looks something like f(x, z) = y
 - x and z are the *arguments* that the function takes
 - *y* is the *output* from the function

FUNCTION MACHINE



--

 $\boldsymbol{\cdot}$ We can make a function that describes a line pretty easily

- $\boldsymbol{\cdot}$ We can make a function that describes a line pretty easily
- $\cdot y = mx + b$

- · We can make a function that describes a line pretty easily
- $\cdot y = mx + b$
 - *m* is the slope (for every one unit increase in *x*, *y* increases *m* units)

- · We can make a function that describes a line pretty easily
- $\cdot y = mx + b$
 - *m* is the slope (for every one unit increase in *x*, *y* increases *m* units)
 - b is the y-intercept: the value of y when x = 0

```
ggplot(data.frame(x = c(-3, 3)), aes(x)) +
  stat function(fun = function(x) -2 * x + 3, geom = "line") +
  stat function(fun = function(x) (1 / 2) * x + 1)
 0
```

QUADRATICS


· These lines have one curve

QUADRATICS

- · These lines have one curve
- $\cdot y = ax^2 + bx + c$


QUADRATICS

```
ggplot(data.frame(x = c(-3, 4)), aes(x)) +
  stat_function(fun = function(x) x ^ 2, color = "green") +
  stat_function(fun = function(x) 2 * x ^ 2, color = "red") +
  stat_function(fun = function(x) -2 * x ^ 2 + 6 * x - 4) +
  ylim(c(-5, 5))
```



EXPONENTIAL

• General form: $y = ab^x$



Logs

• General form: $y = a * \log(bx) + k$

```
ggplot(data.frame(x = c(0, 5)), aes(x)) +
    stat_function(fun = function(x) log(x)) +
    stat_function(fun = function(x) log10(x), color = "red") +
    stat_function(fun = function(x) 2 * log(x), color = "green")
```



14

 $\boldsymbol{\cdot}$ The log and exponent charts are obviously related

- · The log and exponent charts are obviously related
- In fact, these functions are each others "inverse" function

- · The log and exponent charts are obviously related
- In fact, these functions are each others "inverse" function
 - Plug in y to find x

- · The log and exponent charts are obviously related
- In fact, these functions are each others "inverse" function
 - Plug in y to find x
- · Exponents have horizontal asymptote

- · The log and exponent charts are obviously related
- In fact, these functions are each others "inverse" function
 - Plug in y to find x
- · Exponents have horizontal asymptote
- Logs have vertical asymptote

You try!

