

# Set Theory & Combinations

J. Alexander Branham

Fall 2015

# Intro to Set Theory

- What is set theory?

# Intro to Set Theory

- What is set theory?
- A branch of mathematics

# Intro to Set Theory

- What is set theory?
- A branch of mathematics
- Collects objects into *sets* and studies the properties

# What's a set?

- A **set** is a collection of objects

$$S = \{s_1, s_2, s_3, \dots s_n\}$$

# What's a set?

- A **set** is a collection of objects

$$S = \{s_1, s_2, s_3, \dots s_n\}$$

- The objects can be anything

# What's a set?

- A **set** is a collection of objects

$$S = \{s_1, s_2, s_3, \dots s_n\}$$

- The objects can be anything
- We usually use variables or units of observation

# Elements in or not

- We can say whether an object is in a set or not:

$$s_{13} \in S$$



# Elements in or not

- We can say whether an object is in a set or not:

$$s_{13} \in S$$

- Or not:

$$q_1 \notin S$$

# Subsets

- We can divide sets into *subsets*

$$M \subset S$$

# Subsets

- We can divide sets into *subsets*

$$M \subset S$$

- $M$  is a *proper subset* of  $S$  iff all elements of  $M$  are in  $S$  but not all elements of  $S$  are in  $M$ .

# Subsets

- We can divide sets into *subsets*

$$M \subset S$$

- $M$  is a *proper subset* of  $S$  iff all elements of  $M$  are in  $S$  but not all elements of  $S$  are in  $M$ .
- We can also define non-proper subsets:

$$L \subseteq S$$

# Empty sets

- Some sets contain no elements

# Empty sets

- Some sets contain no elements
- For example, cities on Earth with average temperatures of greater than 1000 degrees

# Empty sets

- Some sets contain no elements
- For example, cities on Earth with average temperatures of greater than 1000 degrees
- Though Austin might be close...

$$Z = \{\emptyset\}$$

# Set Universes

- We can talk about how subsets fill a certain **universe** of possibilities



# Set Universes

- We can talk about how subsets fill a certain **universe** of possibilities
- For example:

$$R = 1, 2, 3, 4, 5, 6$$

# Set Universes

- We can talk about how subsets fill a certain **universe** of possibilities
- For example:

$$R = 1, 2, 3, 4, 5, 6$$

- $R$  represents all the possibilities of a (single) roll of a die

# Set Universes

- We can talk about how subsets fill a certain **universe** of possibilities
- For example:

$$R = 1, 2, 3, 4, 5, 6$$

- $R$  represents all the possibilities of a (single) roll of a die
- We can define sets for the even possibilities and the odd possibilities

$$E = \{2, 4, 6\} \quad O = \{1, 3, 5\}$$

# Compliments

- A *compliment* is that together, they contain all the elements of the relevant universe

$$E = O^C \quad ; \quad O = E^C$$

# Universe

- Board examples of how to draw sets

# Intersection, Disjoint, unions

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$

# Intersection, Disjoint, unions

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$ 
  - If  $A \cap B = \emptyset$ , then they are **disjoint**

## Intersection, Disjoint, unions

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$ 
  - If  $A \cap B = \emptyset$ , then they are **disjoint**
- Unions:  $A \cup B$  contains all the elements of both  $A$  and  $B$



## Intersection, Disjoint, unions

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$ 
  - If  $A \cap B = \emptyset$ , then they are **disjoint**
- Unions:  $A \cup B$  contains all the elements of both  $A$  and  $B$ 
  - Two sets unions can't be empty (unless both sets are empty to begin with)

## Intersection, Disjoint, unions

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$ 
  - If  $A \cap B = \emptyset$ , then they are **disjoint**
- Unions:  $A \cup B$  contains all the elements of both  $A$  and  $B$ 
  - Two sets unions can't be empty (unless both sets are empty to begin with)
- Size:  $n(R) = 6$  tells us that there are 6 elements in  $R$

## Intersection, Disjoint, unions

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$ 
  - If  $A \cap B = \emptyset$ , then they are **disjoint**
- Unions:  $A \cup B$  contains all the elements of both  $A$  and  $B$ 
  - Two sets unions can't be empty (unless both sets are empty to begin with)
- Size:  $n(R) = 6$  tells us that there are 6 elements in  $R$
- Sizes of unions

## Intersection, Disjoint, unions

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$ 
  - If  $A \cap B = \emptyset$ , then they are **disjoint**
- Unions:  $A \cup B$  contains all the elements of both  $A$  and  $B$ 
  - Two sets unions can't be empty (unless both sets are empty to begin with)
- Size:  $n(R) = 6$  tells us that there are 6 elements in  $R$
- Sizes of unions
  - $n(X \cup Y) = n(X) + n(Y) - n(X \cap Y)$

# Intersection, Disjoint, unions

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$ 
  - If  $A \cap B = \emptyset$ , then they are **disjoint**
- Unions:  $A \cup B$  contains all the elements of both  $A$  and  $B$ 
  - Two sets unions can't be empty (unless both sets are empty to begin with)
- Size:  $n(R) = 6$  tells us that there are 6 elements in  $R$
- Sizes of unions
  - $n(X \cup Y) = n(X) + n(Y) - n(X \cap Y)$
  - Have to subtract the last term, otherwise we'd double-count elements in the intersection of the two sets

# Properties

- $A \cup B = B \cup A$

# Properties

- $A \cup B = B \cup A$
- $A \cap B = B \cap A$

# Properties

- $A \cup B = B \cup A$
- $A \cap B = B \cap A$
- $(A \cup B) \cup C = A \cup (B \cup C)$



# Properties

- $A \cup B = B \cup A$
- $A \cap B = B \cap A$
- $(A \cup B) \cup C = A \cup (B \cup C)$
- $(A \cap B) \cap C = A \cap (B \cap C)$

# Properties

- $A \cup B = B \cup A$
- $A \cap B = B \cap A$
- $(A \cup B) \cup C = A \cup (B \cup C)$
- $(A \cap B) \cap C = A \cap (B \cap C)$
- $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

# Properties

- $A \cup B = B \cup A$
- $A \cap B = B \cap A$
- $(A \cup B) \cup C = A \cup (B \cup C)$
- $(A \cap B) \cap C = A \cap (B \cap C)$
- $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
- $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$

# Combinatorics - What is it?

- How to count things?

# Combinatorics - What is it?

- How to count things?
- Countable *discrete* structures

# Combinatorics - What is it?

- How to count things?
- Countable *discrete* structures
- Let's say we have the set  $R$  from above

$$R = \{1, 2, 3, 4, 5, 6\}$$

# Combinatorics - What is it?

- How to count things?
- Countable *discrete* structures
- Let's say we have the set  $R$  from above

$$R = \{1, 2, 3, 4, 5, 6\}$$

- How many different combinations of 3 dice rolls are there?

# Order

- First thing to ask is if order matters



# Order

- First thing to ask is if order matters
- Is rolling 1, 4, 2 the same as rolling 4, 2, 1?

# Order

- First thing to ask is if order matters
- Is rolling 1, 4, 2 the same as rolling 4, 2, 1?
- If order matters, we talk about **permutations**

# Order

- First thing to ask is if order matters
- Is rolling 1, 4, 2 the same as rolling 4, 2, 1?
- If order matters, we talk about **permutations**
- If not, it's a **combination**

# Order

- First thing to ask is if order matters
- Is rolling 1, 4, 2 the same as rolling 4, 2, 1?
- If order matters, we talk about **permutations**
- If not, it's a **combination**
- think “permutation” = “position”

# Repetition?

- Can we repeat things?

# Repetition?

- Can we repeat things?
- Is rolling 3, 3, 3, permissible?

# Repetition?

- Can we repeat things?
- Is rolling 3, 3, 3, permissible?
- If yes, we choose one die roll of six three times:

$$6 * 6 * 6 = 6^3$$

# Repetition?

- Can we repeat things?
- Is rolling 3, 3, 3, permissible?
- If yes, we choose one die roll of six three times:

$$6 * 6 * 6 = 6^3$$

- more generally,  $n^r$



# Repetition?

- Can we repeat things?
- Is rolling 3, 3, 3, permissible?
- If yes, we choose one die roll of six three times:

$$6 * 6 * 6 = 6^3$$

- more generally,  $n^r$ 
  - $n$  is number of things to choose from

# Repetition?

- Can we repeat things?
- Is rolling 3, 3, 3, permissible?
- If yes, we choose one die roll of six three times:

$$6 * 6 * 6 = 6^3$$

- more generally,  $n^r$ 
  - $n$  is number of things to choose from
  - choose  $r$

# Permutations without replacement

- We need to reduce the number of things from which to choose each time

# Permutations without replacement

- We need to reduce the number of things from which to choose each time
- How many different orderings of 9 horses in a race??

# Permutations without replacement

- We need to reduce the number of things from which to choose each time
- How many different orderings of 9 horses in a race??
  - $9 * 8 * 7 \dots = 9! = 362,880$

# Permutations without replacement

- We need to reduce the number of things from which to choose each time
- How many different orderings of 9 horses in a race??
  - $9 * 8 * 7 \dots = 9! = 362,880$
- Alternatively, maybe we just care about the first three:

# Permutations without replacement

- We need to reduce the number of things from which to choose each time
- How many different orderings of 9 horses in a race??
  - $9 * 8 * 7 \dots = 9! = 362,880$
- Alternatively, maybe we just care about the first three:
  - $9 * 8 * 7 = 504$

# Permutations without replacement

- We need to reduce the number of things from which to choose each time
- How many different orderings of 9 horses in a race??
  - $9 * 8 * 7 \dots = 9! = 362,880$
- Alternatively, maybe we just care about the first three:
  - $9 * 8 * 7 = 504$
- More generally:

$$\frac{n!}{(n-r)!}$$



# Combinations

- We use combinations when the order does **not** matter

# Combinations

- We use combinations when the order does **not** matter
- So 4, 3, 1 is the same as 3, 1, 4

## Repetition not allowed

- Easiest way: do permutation without repetition, then get rid of all the “same” outcomes

## Repetition not allowed

- Easiest way: do permutation without repetition, then get rid of all the “same” outcomes
- We’ve already figured out the permutation part, so need to figure out the second part

$$\frac{n!}{(n-r)!} * \frac{1}{r!} = \frac{n!}{r!(n-r)!} = \binom{n}{r}$$

## Repetition allowed

- These aren't used much

## Repetition allowed

- These aren't used much
- Also the most difficult to calculate

## Repetition allowed

- These aren't used much
- Also the most difficult to calculate
- Formula:

$$\binom{n+r-1}{r} = \frac{(n+r-1)!}{r!(n-1)!}$$