

FUNCTIONS

J. Alexander Branham

Fall 2017

EXPONENTS AND LOGARITHMS

- Exponents tell you to multiply that thing by its base x times:

- Exponents tell you to multiply that thing by its base x times:

- $3^4 = 3 * 3 * 3 * 3 =$

EXPONENTS AND LOGARITHMS

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 3 * 3 * 3 * 3 =$
- Logarithms ask how many times you must raise the base to get x:

EXPONENTS AND LOGARITHMS

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 3 * 3 * 3 * 3 =$
- Logarithms ask how many times you must raise the base to get x :
 - $\log_3(81) = 4$

EXPONENTS AND LOGARITHMS

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 3 * 3 * 3 * 3 =$
- Logarithms ask how many times you must raise the base to get x :
 - $\log_3(81) = 4$
 - Note that logarithms with negative arguments are undefined

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 3 * 3 * 3 * 3 =$
- Logarithms ask how many times you must raise the base to get x :
 - $\log_3(81) = 4$
 - Note that logarithms with negative arguments are undefined
 - Sometimes $\log(n)$ means $\log_{10}(n)$

EXPONENTS AND LOGARITHMS

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 3 * 3 * 3 * 3 =$
- Logarithms ask how many times you must raise the base to get x :
 - $\log_3(81) = 4$
 - Note that logarithms with negative arguments are undefined
 - Sometimes $\log(n)$ means $\log_{10}(n)$
 - Othertimes, it means $\log_e(n) = \ln(n)$

$$a^m a^n = a^{m+n}$$

PROPERTIES OF EXPONENTS

$$a^m a^n = a^{m+n}$$

$$\frac{a^m}{a^n} = a^{m-n}$$

PROPERTIES OF EXPONENTS

$$a^m a^n = a^{m+n}$$

$$\frac{a^m}{a^n} = a^{m-n}$$

$$(a^m)^n = a^{mn}$$

PROPERTIES OF EXPONENTS

$$a^m a^n = a^{m+n}$$

$$\frac{a^m}{a^n} = a^{m-n}$$

$$(a^m)^n = a^{mn}$$

$$a^0 = 1$$

PROPERTIES OF EXPONENTS (CONTINUED)

$$a^{1/n} = \sqrt[n]{a}$$

PROPERTIES OF EXPONENTS (CONTINUED)

$$a^{1/n} = \sqrt[n]{a}$$

$$\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n} = a^n b^{-n} \quad \forall b \neq 0$$

PROPERTIES OF EXPONENTS (CONTINUED)

$$a^{1/n} = \sqrt[n]{a}$$

$$\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n} = a^n b^{-n} \quad \forall b \neq 0$$

$$\left(\frac{a}{b}\right)^{-n} = \left(\frac{b}{a}\right)^n \quad \forall a, b \neq 0$$

$$\log_x(ab) = \log_x a + \log_x b$$

$$\log_x(ab) = \log_x a + \log_x b$$

$$\log_x \left(\frac{a}{b} \right) = \log_x a - \log_x b$$

$$\log_x(ab) = \log_x a + \log_x b$$

$$\log_x \left(\frac{a}{b} \right) = \log_x a - \log_x b$$

$$\log_x a^b = b \log_x a$$

$$\log_x(ab) = \log_x a + \log_x b$$

$$\log_x \left(\frac{a}{b} \right) = \log_x a - \log_x b$$

$$\log_x a^b = b \log_x a$$

$$\log_x 1 = 0$$

PROPERTIES OF LOGS (CONTINUED)

$$m^{\log_m(a)} = a$$

PROPERTIES OF LOGS (CONTINUED)

$$m^{\log_m(a)} = a$$

$$\frac{\log_x n}{\log_x m} = \log_m n$$

YOU TRY!

$$2^4$$

$$\log(100)$$

$$a^3 \times a^7$$

$$\log_{10}(10z)$$

FUNCTIONS

WHAT'S A FUNCTION?

- Anything that takes input and gives one output

WHAT'S A FUNCTION?

- Anything that takes input and gives one output
- In math, this usually looks something like $f(x, z) = y$


WHAT'S A FUNCTION?

- Anything that takes input and gives one output
- In math, this usually looks something like $f(x, z) = y$
 - x and z are the *arguments* that the function takes

WHAT'S A FUNCTION?

- Anything that takes input and gives one output
- In math, this usually looks something like $f(x, z) = y$
 - x and z are the *arguments* that the function takes
 - y is the *output* from the function

FUNCTION MACHINE



- We can make a function that describes a line pretty easily


- We can make a function that describes a line pretty easily
- $y = mx + b$

- We can make a function that describes a line pretty easily
- $y = mx + b$
 - m is the slope (for every one unit increase in x , y increases m units)

- We can make a function that describes a line pretty easily
- $y = mx + b$
 - m is the slope (for every one unit increase in x , y increases m units)
 - b is the y-intercept: the value of y when $x = 0$

LINEAR FUNCTIONS

```
ggplot(data.frame(x = c(-3, 3)), aes(x)) +  
  stat_function(fun = function(x) -2 * x + 3, geom = "line") +  
  stat_function(fun = function(x) (1 / 2) * x + 1)
```




- These lines have one curve

- These lines have one curve
- $y = ax^2 + bx + c$

QUADRATICS


```
ggplot(data.frame(x = c(-3, 4)), aes(x)) +  
  stat_function(fun = function(x) x ^ 2, color = "green") +  
  stat_function(fun = function(x) 2 * x ^ 2, color = "red") +  
  stat_function(fun = function(x) -2 * x ^ 2 + 6 * x - 4) +  
  ylim(c(-5, 5))
```



EXPONENTIAL

- General form: $y = ab^x$


```
ggplot(data.frame(x = c(-1, 5)), aes(x)) +  
  stat_function(fun = function(x) exp(x)) +  
  stat_function(fun = function(x) 2 * exp(x), color = "red")
```



LOGS

- General form: $y = a * \log(bx) + k$

```
ggplot(data.frame(x = c(0, 5)), aes(x)) +  
  stat_function(fun = function(x) log(x)) +  
  stat_function(fun = function(x) log10(x), color = "red") +  
  stat_function(fun = function(x) 2 * log(x), color = "green")
```



- The log and exponent charts are obviously related

- The log and exponent charts are obviously related
- In fact, these functions are each others “inverse” function

- The log and exponent charts are obviously related
- In fact, these functions are each others “inverse” function
 - Plug in y to find x

- The log and exponent charts are obviously related
- In fact, these functions are each others “inverse” function
 - Plug in y to find x
- Exponents have horizontal asymptote

- The log and exponent charts are obviously related
- In fact, these functions are each others “inverse” function
 - Plug in y to find x
- Exponents have horizontal asymptote
- Logs have vertical asymptote

YOU TRY!

What function describes this line?

