

Marginal, Joint, and Conditional Probabilities

J. Alexander Branham

Fall 2015

Marginal probability

- ▶ The Pr of an event occurring:

Marginal probability

- ▶ The Pr of an event occurring:
 - ▶ $Pr(A)$

Marginal probability

- ▶ The Pr of an event occurring:
 - ▶ $Pr(A)$
- ▶ Unconditional probability

Marginal probability

- ▶ The Pr of an event occurring:
 - ▶ $Pr(A)$
- ▶ Unconditional probability
- ▶ Probability of drawing a red card from a standard deck:

$$Pr(red) = \frac{1}{2}$$

Joint probability

- ▶ The probability of both A **and** B

Joint probability

- ▶ The probability of both A **and** B
 - ▶ $Pr(A, B)$

Joint probability

- ▶ The probability of both A **and** B
 - ▶ $Pr(A, B)$
- ▶ This is the intersection of the two sets

Joint probability

- ▶ The probability of both A **and** B
 - ▶ $Pr(A, B)$
- ▶ This is the intersection of the two sets
- ▶ $Pr(A \cap B)$

Joint probability

- ▶ The probability of both A **and** B
 - ▶ $Pr(A, B)$
- ▶ This is the intersection of the two sets
- ▶ $Pr(A \cap B)$
 - ▶ Probability of drawing a red card and a 4:

Joint probability

- ▶ The probability of both A **and** B
 - ▶ $Pr(A, B)$
- ▶ This is the intersection of the two sets
- ▶ $Pr(A \cap B)$
 - ▶ Probability of drawing a red card and a 4:
 - ▶ $Pr(\text{red}, 4) = \frac{2}{52} = \frac{1}{26}$

Conditional probability

- ▶ The probability of some event happening **given** some other event having occurred

Conditional probability

- ▶ The probability of some event happening **given** some other event having occurred
 - ▶ $Pr(A|B)$

Conditional probability

- ▶ The probability of some event happening **given** some other event having occurred
 - ▶ $Pr(A|B)$
- ▶ You've drawn a red card. What's the probability it's a four?

Conditional probability

- ▶ The probability of some event happening **given** some other event having occurred
 - ▶ $Pr(A|B)$
- ▶ You've drawn a red card. What's the probability it's a four?
 - ▶ $Pr(4|red) = \frac{2}{26} = \frac{1}{13}$

How to convert marginal, joint, and conditional probabilities

$$Pr(A|B) = \frac{Pr(A, B)}{Pr(B)}$$

- * The conditional probability is equal to the joint probability over the probability of the condition

Bayes' Law (Theorem)

$$Pr(A|B) = \frac{Pr(B|A)Pr(A)}{Pr(B)} = \frac{Pr(B|A)Pr(A)}{Pr(B|A)Pr(A) + Pr(B|\neg A)Pr(\neg A)}$$

Bayes Example

- ▶ Police set up roadblock to randomly screen for potentially drunk drivers

Bayes Example

- ▶ Police set up roadblock to randomly screen for potentially drunk drivers
- ▶ Person stopped and fails breathalyzer test

Bayes Example

- ▶ Police set up roadblock to randomly screen for potentially drunk drivers
- ▶ Person stopped and fails breathalyzer test
 - ▶ Test gives positive 98 percent of the time when person is drunk

Bayes Example

- ▶ Police set up roadblock to randomly screen for potentially drunk drivers
- ▶ Person stopped and fails breathalyzer test
 - ▶ Test gives positive 98 percent of the time when person is drunk
 - ▶ Also gives positive 5 percent of the time when the person is actually sober

Bayes Example

- ▶ Police set up roadblock to randomly screen for potentially drunk drivers
- ▶ Person stopped and fails breathalyzer test
 - ▶ Test gives positive 98 percent of the time when person is drunk
 - ▶ Also gives positive 5 percent of the time when the person is actually sober
 - ▶ We think that 1 percent of people going through the checkpoint are drunk

Bayes Example

- ▶ Police set up roadblock to randomly screen for potentially drunk drivers
- ▶ Person stopped and fails breathalyzer test
 - ▶ Test gives positive 98 percent of the time when person is drunk
 - ▶ Also gives positive 5 percent of the time when the person is actually sober
 - ▶ We think that 1 percent of people going through the checkpoint are drunk
- ▶ What is the probability that the man was actually drunk?

DWI Example

- ▶ Since he's randomly stopped, we think there is a 1 percent chance he's drunk

DWI Example

- ▶ Since he's randomly stopped, we think there is a 1 percent chance he's drunk
 - ▶ $Pr(drunk) = 0.01$

DWI Example

- ▶ Since he's randomly stopped, we think there is a 1 percent chance he's drunk
 - ▶ $Pr(drunk) = 0.01$
- ▶ Test accuracy:

DWI Example

- ▶ Since he's randomly stopped, we think there is a 1 percent chance he's drunk
 - ▶ $Pr(drunk) = 0.01$
- ▶ Test accuracy:
 - ▶ $Pr(positive|drunk) = 0.98$

DWI Example

- ▶ Since he's randomly stopped, we think there is a 1 percent chance he's drunk
 - ▶ $Pr(drunk) = 0.01$
- ▶ Test accuracy:
 - ▶ $Pr(positive|drunk) = 0.98$
 - ▶ $Pr(positive|\neg drunk) = 0.05$

DWI Example

- ▶ Since he's randomly stopped, we think there is a 1 percent chance he's drunk
 - ▶ $Pr(drunk) = 0.01$
- ▶ Test accuracy:
 - ▶ $Pr(positive|drunk) = 0.98$
 - ▶ $Pr(positive|\neg drunk) = 0.05$
- ▶ Want to know:

DWI Example

- ▶ Since he's randomly stopped, we think there is a 1 percent chance he's drunk

- ▶ $Pr(drunk) = 0.01$

- ▶ Test accuracy:

- ▶ $Pr(positive|drunk) = 0.98$

- ▶ $Pr(positive|\neg drunk) = 0.05$

- ▶ Want to know:

- ▶ $Pr(drunk|positive) = \frac{Pr(drunk)Pr(positive|drunk)}{Pr(positive)}$

DWI Example

- ▶ Since he's randomly stopped, we think there is a 1 percent chance he's drunk
 - ▶ $Pr(drunk) = 0.01$
- ▶ Test accuracy:
 - ▶ $Pr(positive|drunk) = 0.98$
 - ▶ $Pr(positive|\neg drunk) = 0.05$
- ▶ Want to know:
 - ▶ $Pr(drunk|positive) = \frac{Pr(drunk)Pr(positive|drunk)}{Pr(positive)}$
- ▶ Everything known, except $Pr(positive)$

Pr(positive)

$$\begin{aligned}Pr(\textit{positive}) &= Pr(\textit{positive}|\textit{drunk})Pr(\textit{drunk}) \\&\quad + Pr(\textit{positive}|\neg\textit{drunk})Pr(\neg\textit{drunk}) \\&= .98(.01) + .05(.99) \\&\approx .0593\end{aligned}$$

$Pr(drunk|positive)$

$$\blacktriangleright Pr(drunk|positive) = \frac{Pr(drunk)Pr(positive|drunk)}{Pr(positive)}$$

$Pr(drunk|positive)$

- ▶ $Pr(drunk|positive) = \frac{Pr(drunk)Pr(positive|drunk)}{Pr(positive)}$
- ▶ $Pr(drunk|positive) = \frac{0.01(.98)}{.0593}$

$Pr(drunk|positive)$

- ▶ $Pr(drunk|positive) = \frac{Pr(drunk)Pr(positive|drunk)}{Pr(positive)}$
- ▶ $Pr(drunk|positive) = \frac{0.01(.98)}{.0593}$
- ▶ .165

$Pr(drunk|positive)$

- ▶ $Pr(drunk|positive) = \frac{Pr(drunk)Pr(positive|drunk)}{Pr(positive)}$
- ▶ $Pr(drunk|positive) = \frac{0.01(.98)}{.0593}$
- ▶ .165
- ▶ So there is a 16.5 percent chance that the man is drunk given that he tested positive

Example, take two: Monty Hall

- ▶ *Let's Make a Deal* was a game show where contestants were asked to choose one of three doors (A, B, and C). Behind one is a car and behind the other two are goats

Example, take two: Monty Hall

- ▶ *Let's Make a Deal* was a game show where contestants were asked to choose one of three doors (A, B, and C). Behind one is a car and behind the other two are goats
- ▶ Once contestants chose a door, Monty would open one of the doors they *didn't* choose that had a goat behind it

Example, take two: Monty Hall

- ▶ *Let's Make a Deal* was a game show where contestants were asked to choose one of three doors (A, B, and C). Behind one is a car and behind the other two are goats
- ▶ Once contestants chose a door, Monty would open one of the doors they *didn't* choose that had a goat behind it
- ▶ The contestant could then either stick with their original choice or switch to the unopened door

Example, take two: Monty Hall

- ▶ *Let's Make a Deal* was a game show where contestants were asked to choose one of three doors (A, B, and C). Behind one is a car and behind the other two are goats
- ▶ Once contestants chose a door, Monty would open one of the doors they *didn't* choose that had a goat behind it
- ▶ The contestant could then either stick with their original choice or switch to the unopened door
- ▶ Does switching doors increase your probability of winning?

Example, take two: Monty Hall

- ▶ *Let's Make a Deal* was a game show where contestants were asked to choose one of three doors (A, B, and C). Behind one is a car and behind the other two are goats
- ▶ Once contestants chose a door, Monty would open one of the doors they *didn't* choose that had a goat behind it
- ▶ The contestant could then either stick with their original choice or switch to the unopened door
- ▶ Does switching doors increase your probability of winning?
 - ▶ YES!

Example, take two: Monty Hall

- ▶ *Let's Make a Deal* was a game show where contestants were asked to choose one of three doors (A, B, and C). Behind one is a car and behind the other two are goats
- ▶ Once contestants chose a door, Monty would open one of the doors they *didn't* choose that had a goat behind it
- ▶ The contestant could then either stick with their original choice or switch to the unopened door
- ▶ Does switching doors increase your probability of winning?
 - ▶ YES!
 - ▶ Nearly 10,000 people wrote in refusing to believe this (including 1,000 PhDs!)

Bayes's solution to Monty Hall

- ▶ $Pr(A) = Pr(B) = Pr(C) = \frac{1}{3}$ (prior belief about where the car is)

Bayes's solution to Monty Hall

- ▶ $Pr(A) = Pr(B) = Pr(C) = \frac{1}{3}$ (prior belief about where the car is)
- ▶ WLOG assume contestant picks door A

Bayes's solution to Monty Hall

- ▶ $Pr(A) = Pr(B) = Pr(C) = \frac{1}{3}$ (prior belief about where the car is)
- ▶ WLOG assume contestant picks door A
- ▶ WLOG assume Monty opens door B

Bayes's solution to Monty Hall

- ▶ $Pr(A) = Pr(B) = Pr(C) = \frac{1}{3}$ (prior belief about where the car is)
- ▶ WLOG assume contestant picks door A
- ▶ WLOG assume Monty opens door B
- ▶ Then: $Pr(B_{Monty}|A) = \frac{1}{2}$ and $Pr(B_{Monty}|B = 0)$ and $Pr(C_{Monty}|C) = 1$

Bayes's solution to Monty Hall

- ▶ $Pr(A) = Pr(B) = Pr(C) = \frac{1}{3}$ (prior belief about where the car is)
- ▶ WLOG assume contestant picks door A
- ▶ WLOG assume Monty opens door B
- ▶ Then: $Pr(B_{Monty}|A) = \frac{1}{2}$ and $Pr(B_{Monty}|B = 0)$ and $Pr(C_{Monty}|C) = 1$

Bayes's solution to Monty Hall

- ▶ $Pr(A) = Pr(B) = Pr(C) = \frac{1}{3}$ (prior belief about where the car is)
- ▶ WLOG assume contestant picks door A
- ▶ WLOG assume Monty opens door B
- ▶ Then: $Pr(B_{Monty}|A) = \frac{1}{2}$ and $Pr(B_{Monty}|B = 0)$ and $Pr(C_{Monty}|C) = 1$

$$Pr(B_{Monty}, A) = Pr(B_{Monty}|A)Pr(A) = \frac{1}{2} * \frac{1}{3} = \frac{1}{6}$$

$$Pr(B_{Monty}, B) = Pr(B_{Monty}|B)Pr(B) = 0 * \frac{1}{3} = 0$$

$$Pr(B_{Monty}, C) = Pr(B_{Monty}|C)Pr(C) = 1 * \frac{1}{3} = \frac{1}{3}$$

Bayes's solution to Monty Hall

- ▶ $Pr(A) = Pr(B) = Pr(C) = \frac{1}{3}$ (prior belief about where the car is)
- ▶ WLOG assume contestant picks door A
- ▶ WLOG assume Monty opens door B
- ▶ Then: $Pr(B_{Monty}|A) = \frac{1}{2}$ and $Pr(B_{Monty}|B = 0)$ and $Pr(C_{Monty}|C) = 1$

$$Pr(B_{Monty}, A) = Pr(B_{Monty}|A)Pr(A) = \frac{1}{2} * \frac{1}{3} = \frac{1}{6}$$

$$Pr(B_{Monty}, B) = Pr(B_{Monty}|B)Pr(B) = 0 * \frac{1}{3} = 0$$

$$Pr(B_{Monty}, C) = Pr(B_{Monty}|C)Pr(C) = 1 * \frac{1}{3} = \frac{1}{3}$$

- ▶ Note that $Pr(B_{Monty}) =$

$$Pr(B_{Monty}, A) + Pr(B_{Monty}, B) + Pr(B_{Monty}, C) = \frac{1}{6} + 0 + \frac{1}{3} = \frac{1}{2}$$

$Pr(A|B_{Monty})$ and $Pr(C|B_{Monty})$

$$\blacktriangleright Pr(A|B_{Monty}) = \frac{Pr(A)Pr(B_{Monty}|A)}{Pr(B_{Monty})} = \frac{\frac{1}{3} * \frac{1}{2}}{\frac{1}{2}} = \frac{1}{3}$$

$Pr(A|B_{Monty})$ and $Pr(C|B_{Monty})$

$$\begin{aligned}\blacktriangleright Pr(A|B_{Monty}) &= \frac{Pr(A)Pr(B_{Monty}|A)}{Pr(B_{Monty})} = \frac{\frac{1}{3} * \frac{1}{2}}{\frac{1}{2}} = \frac{1}{3} \\ \blacktriangleright Pr(C|B_{Monty}) &= \frac{Pr(C)Pr(B_{Monty}|C)}{Pr(B_{Monty})} = \frac{\frac{1}{3} * 1}{\frac{1}{2}} = \frac{2}{3}\end{aligned}$$

$Pr(A|B_{Monty})$ and $Pr(C|B_{Monty})$

$$\begin{aligned}\text{▶ } Pr(A|B_{Monty}) &= \frac{Pr(A)Pr(B_{Monty}|A)}{Pr(B_{Monty})} = \frac{\frac{1}{3} * \frac{1}{2}}{\frac{1}{2}} = \frac{1}{3} \\ \text{▶ } Pr(C|B_{Monty}) &= \frac{Pr(C)Pr(B_{Monty}|C)}{Pr(B_{Monty})} = \frac{\frac{1}{3} * 1}{\frac{1}{2}} = \frac{2}{3}\end{aligned}$$

▶ So switch!