

# SET THEORY & COMBINATIONS

---

J. Alexander Branham

Fall 2016

# SET THEORY

---

- What is set theory?

- What is set theory?
- A branch of mathematics

- What is set theory?
- A branch of mathematics
- Collects objects into *sets* and studies the properties

# WHAT'S A SET?

- A **set** is a collection of objects

$$S = \{s_1, s_2, s_3, \dots s_n\}$$

# WHAT'S A SET?

- A **set** is a collection of objects

$$S = \{s_1, s_2, s_3, \dots s_n\}$$

- The objects can be anything

# WHAT'S A SET?

- A **set** is a collection of objects

$$S = \{s_1, s_2, s_3, \dots s_n\}$$

- The objects can be anything
- We usually use variables or units of observation



- We can say whether an object is in a set:

$$s_{13} \in S$$

- We can say whether an object is in a set:

$$s_{13} \in S$$

- Or not:

$$q_1 \notin S$$

- We can divide sets into *subsets*

$$M \subset S$$

# SUBSETS

- We can divide sets into *subsets*

$$M \subset S$$

- $M$  is a *proper subset* of  $S$  iff all elements of  $M$  are in  $S$  but not all elements of  $S$  are in  $M$ .

# SUBSETS

- We can divide sets into *subsets*

$$M \subset S$$

- $M$  is a *proper subset* of  $S$  iff all elements of  $M$  are in  $S$  but not all elements of  $S$  are in  $M$ .
- We can also define non-proper subsets:

$$L \subseteq S$$

- Some sets contain no elements

- Some sets contain no elements
- For example, cities on Earth with average temperatures of greater than 1000 degrees

- Some sets contain no elements
- For example, cities on Earth with average temperatures of greater than 1000 degrees
- Though Austin might be close...

$$Z = \{\emptyset\}$$



- We can talk about how subsets fill a certain **universe** of possibilities

- We can talk about how subsets fill a certain **universe** of possibilities
- For example:

$$R = \{1, 2, 3, 4, 5, 6\}$$

- We can talk about how subsets fill a certain **universe** of possibilities
- For example:

$$R = \{1, 2, 3, 4, 5, 6\}$$

- $R$  represents all the possibilities of a (single) roll of a die

- We can talk about how subsets fill a certain **universe** of possibilities
- For example:

$$R = \{1, 2, 3, 4, 5, 6\}$$

- $R$  represents all the possibilities of a (single) roll of a die
- We can define sets for the even possibilities and the odd possibilities

$$E = \{2, 4, 6\} \quad O = \{1, 3, 5\}$$

- A *compliment* is that together, they contain all the elements of the relevant universe

$$E = O^c \qquad O = E^c$$

- Board examples of how to draw sets

## INTERSECTION, DISJOINT, UNIONS

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$

## INTERSECTION, DISJOINT, UNIONS

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$ 
  - If  $A \cap B = \emptyset$ , then they are **disjoint**



## INTERSECTION, DISJOINT, UNIONS

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$ 
  - If  $A \cap B = \emptyset$ , then they are **disjoint**
- Unions:  $A \cup B$  contains all the elements of both  $A$  and  $B$

## INTERSECTION, DISJOINT, UNIONS

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$ 
  - If  $A \cap B = \emptyset$ , then they are **disjoint**
- Unions:  $A \cup B$  contains all the elements of both  $A$  and  $B$ 
  - Two sets unions can't be empty (unless both sets are empty to begin with)

## INTERSECTION, DISJOINT, UNIONS

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$ 
  - If  $A \cap B = \emptyset$ , then they are **disjoint**
- Unions:  $A \cup B$  contains all the elements of both  $A$  and  $B$ 
  - Two sets unions can't be empty (unless both sets are empty to begin with)
- Size:  $n(R) = 6$  tells us that there are 6 elements in  $R$

## INTERSECTION, DISJOINT, UNIONS

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$ 
  - If  $A \cap B = \emptyset$ , then they are **disjoint**
- Unions:  $A \cup B$  contains all the elements of both  $A$  and  $B$ 
  - Two sets unions can't be empty (unless both sets are empty to begin with)
- Size:  $n(R) = 6$  tells us that there are 6 elements in  $R$
- Sizes of unions

## INTERSECTION, DISJOINT, UNIONS

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$ 
  - If  $A \cap B = \emptyset$ , then they are **disjoint**
- Unions:  $A \cup B$  contains all the elements of both  $A$  and  $B$ 
  - Two sets unions can't be empty (unless both sets are empty to begin with)
- Size:  $n(R) = 6$  tells us that there are 6 elements in  $R$
- Sizes of unions
  - $n(X \cup Y) = n(X) + n(Y) - n(X \cap Y)$

## INTERSECTION, DISJOINT, UNIONS

- Intersections:  $A \cap B$  includes all the common elements of  $A$  and  $B$ 
  - If  $A \cap B = \emptyset$ , then they are **disjoint**
- Unions:  $A \cup B$  contains all the elements of both  $A$  and  $B$ 
  - Two sets unions can't be empty (unless both sets are empty to begin with)
- Size:  $n(R) = 6$  tells us that there are 6 elements in  $R$
- Sizes of unions
  - $n(X \cup Y) = n(X) + n(Y) - n(X \cap Y)$
  - Have to subtract the last term, otherwise we'd double-count elements in the intersection of the two sets

- $A \cup B = B \cup A$

- $A \cup B = B \cup A$
- $A \cap B = B \cap A$



- $A \cup B = B \cup A$
- $A \cap B = B \cap A$
- $(A \cup B) \cup C = A \cup (B \cup C)$

- $A \cup B = B \cup A$
- $A \cap B = B \cap A$
- $(A \cup B) \cup C = A \cup (B \cup C)$
- $(A \cap B) \cap C = A \cap (B \cap C)$

- $A \cup B = B \cup A$
- $A \cap B = B \cap A$
- $(A \cup B) \cup C = A \cup (B \cup C)$
- $(A \cap B) \cap C = A \cap (B \cap C)$
- $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

- $A \cup B = B \cup A$
- $A \cap B = B \cap A$
- $(A \cup B) \cup C = A \cup (B \cup C)$
- $(A \cap B) \cap C = A \cap (B \cap C)$
- $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
- $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$

## YOU TRY!

1. Define the universe of cards in a standard 52-card deck —  $\Omega$

## YOU TRY!

1. Define the universe of cards in a standard 52-card deck —  $\Omega$
2. Define the set containing all the spades —  $S$  — the set containing the black cards —  $B$  — and the set of queens  $Q$

## YOU TRY!

1. Define the universe of cards in a standard 52-card deck —  $\Omega$
2. Define the set containing all the spades —  $S$  — the set containing the black cards —  $B$  — and the set of queens  $Q$
3. What is  $n(B)$

## YOU TRY!

1. Define the universe of cards in a standard 52-card deck —  $\Omega$
2. Define the set containing all the spades —  $S$  — the set containing the black cards —  $B$  — and the set of queens  $Q$
3. What is  $n(B)$
4. What is  $S \cup B$ ?



## YOU TRY!

1. Define the universe of cards in a standard 52-card deck —  $\Omega$
2. Define the set containing all the spades —  $S$  — the set containing the black cards —  $B$  — and the set of queens  $Q$
3. What is  $n(B)$
4. What is  $S \cup B$ ?
5. What is  $B \cap Q$ ?

## YOU TRY!

1. Define the universe of cards in a standard 52-card deck —  $\Omega$
2. Define the set containing all the spades —  $S$  — the set containing the black cards —  $B$  — and the set of queens  $Q$
3. What is  $n(B)$
4. What is  $S \cup B$ ?
5. What is  $B \cap Q$ ?
6. What is  $B \cap (Q \cup S)$ ?

## YOU TRY!

1. Define the universe of cards in a standard 52-card deck —  $\Omega$
2. Define the set containing all the spades —  $S$  — the set containing the black cards —  $B$  — and the set of queens  $Q$
3. What is  $n(B)$
4. What is  $S \cup B$ ?
5. What is  $B \cap Q$ ?
6. What is  $B \cap (Q \cup S)$ ?
7. Is  $Q \subset S$ ?

## YOU TRY (ANSWERS)

1.  $\Omega$  is a set with 52 elements, one for each card

## YOU TRY (ANSWERS)

1.  $\Omega$  is a set with 52 elements, one for each card
2.  $S$  is a set with 13 elements, one for each spade.  $B$  has 26 elements (the black cards) and  $Q$  has just four - the queens

## YOU TRY (ANSWERS)

1.  $\Omega$  is a set with 52 elements, one for each card
2.  $S$  is a set with 13 elements, one for each spade.  $B$  has 26 elements (the black cards) and  $Q$  has just four - the queens
3.  $n(B) = 26$

## YOU TRY (ANSWERS)

1.  $\Omega$  is a set with 52 elements, one for each card
2.  $S$  is a set with 13 elements, one for each spade.  $B$  has 26 elements (the black cards) and  $Q$  has just four - the queens
3.  $n(B) = 26$
4.  $S \cup B = B$

## YOU TRY (ANSWERS)

1.  $\Omega$  is a set with 52 elements, one for each card
2.  $S$  is a set with 13 elements, one for each spade.  $B$  has 26 elements (the black cards) and  $Q$  has just four - the queens
3.  $n(B) = 26$
4.  $S \cup B = B$
5.  $B \cap Q$  is the queen of spades and clubs



## YOU TRY (ANSWERS)

1.  $\Omega$  is a set with 52 elements, one for each card
2.  $S$  is a set with 13 elements, one for each spade.  $B$  has 26 elements (the black cards) and  $Q$  has just four - the queens
3.  $n(B) = 26$
4.  $S \cup B = B$
5.  $B \cap Q$  is the queen of spades and clubs
6.  $B \cap (Q \cup S) = B$

## YOU TRY (ANSWERS)

1.  $\Omega$  is a set with 52 elements, one for each card
2.  $S$  is a set with 13 elements, one for each spade.  $B$  has 26 elements (the black cards) and  $Q$  has just four - the queens
3.  $n(B) = 26$
4.  $S \cup B = B$
5.  $B \cap Q$  is the queen of spades and clubs
6.  $B \cap (Q \cup S) = B$
7. No

# COMBINATORICS

---

- How to count things?

# COMBINATORICS - WHAT IS IT?

- How to count things?
- Countable *discrete* structures

- How to count things?
- Countable *discrete* structures
- Let's say we have the set  $R$  from above

$$R = \{1, 2, 3, 4, 5, 6\}$$

- How to count things?
- Countable *discrete* structures
- Let's say we have the set  $R$  from above

$$R = \{1, 2, 3, 4, 5, 6\}$$

- How many different combinations of 3 dice rolls are there?

- First thing to ask is if order matters



- First thing to ask is if order matters
- Is rolling 1, 4, 2 the same as rolling 4, 2, 1?

- First thing to ask is if order matters
- Is rolling 1, 4, 2 the same as rolling 4, 2, 1?
- If order matters, we talk about **permutations**

- First thing to ask is if order matters
- Is rolling 1, 4, 2 the same as rolling 4, 2, 1?
- If order matters, we talk about **permutations**
- If not, it's a **combination**

- First thing to ask is if order matters
- Is rolling 1, 4, 2 the same as rolling 4, 2, 1?
- If order matters, we talk about **permutations**
- If not, it's a **combination**
- think “permutation” = “position”

# REPITITION?

- Can we repeat things?

## REPITITION?

- Can we repeat things?
- Is rolling 3, 3, 3, permissable?

## REPITITION?

- Can we repeat things?
- Is rolling 3, 3, 3, permissable?
- If yes, we choose one die roll of six three times:

$$6 * 6 * 6 = 6^3$$

# REPITITION?

- Can we repeat things?
- Is rolling 3, 3, 3, permissable?
- If yes, we choose one die roll of six three times:

$$6 * 6 * 6 = 6^3$$

- more generally,  $n^r$



# REPITITION?

- Can we repeat things?
- Is rolling 3, 3, 3, permissable?
- If yes, we choose one die roll of six three times:

$$6 * 6 * 6 = 6^3$$

- more generally,  $n^r$ 
  - $n$  is number of things to choose from

# REPITITION?

- Can we repeat things?
- Is rolling 3, 3, 3, permissable?
- If yes, we choose one die roll of six three times:

$$6 * 6 * 6 = 6^3$$

- more generally,  $n^r$ 
  - $n$  is number of things to choose from
  - choose  $r$

## PERMUTATIONS WITHOUT REPLACEMENT

- We need to reduce the number of things from which to choose each time

## PERMUTATIONS WITHOUT REPLACEMENT

- We need to reduce the number of things from which to choose each time
- How many different orderings of 9 horses in a race??

## PERMUTATIONS WITHOUT REPLACEMENT

- We need to reduce the number of things from which to choose each time
- How many different orderings of 9 horses in a race??
  - $9 * 8 * 7 \dots = 9! = 362,880$

## PERMUTATIONS WITHOUT REPLACEMENT

- We need to reduce the number of things from which to choose each time
- How many different orderings of 9 horses in a race??
  - $9 * 8 * 7 \dots = 9! = 362,880$
- Alternatively, maybe we just care about the first three:

## PERMUTATIONS WITHOUT REPLACEMENT

- We need to reduce the number of things from which to choose each time
- How many different orderings of 9 horses in a race??
  - $9 * 8 * 7 \dots = 9! = 362,880$
- Alternatively, maybe we just care about the first three:
  - $9 * 8 * 7 = 504$

## PERMUTATIONS WITHOUT REPLACEMENT

- We need to reduce the number of things from which to choose each time
- How many different orderings of 9 horses in a race??

- $9 * 8 * 7 \dots = 9! = 362,880$

- Alternatively, maybe we just care about the first three:

- $9 * 8 * 7 = 504$

- More generally:

$$\frac{n!}{(n-r)!}$$



- We use combinations when the order does **not** matter

- We use combinations when the order does **not** matter
- So 4, 3, 1 is the same as 3, 1, 4

- Easiest way: do permutation without repitition, then get rid of all the “same” outcomes

- Easiest way: do permutation without repitition, then get rid of all the “same” outcomes
- We’ve already figured out the permutation part, so need to figure out the second part

$$\frac{n!}{(n-r)!} * \frac{1}{r!} = \frac{n!}{r!(n-r)!} = \binom{n}{r}$$

- These aren't used much

- These aren't used much
- Also the most difficult to calculate

- These aren't used much
- Also the most difficult to calculate
- Formula:

$$\binom{n+r-1}{r} = \frac{(n+r-1)!}{r!(n-1)!}$$

## YOU TRY!

Using a standard 52-card deck:

1. How many different 3-card hands exist?



## YOU TRY!

Using a standard 52-card deck:

1. How many different 3-card hands exist?
2. How many ways are there to get dealt the queen of spades, then the king of diamonds, then the eight of spades?

## YOU TRY (ANSWERS)

```
# 1
```

```
choose(52, 3)
```

```
## [1] 22100
```

```
# 2 - one
```