



Applied Data Science Program

TIME SERIES

Munther A. Dahleh

Learning Time-Series


- Time Series are everywhere
 - Finance, weather, control systems
- Questions:
 - Modeling
 - Forecasting
 - Decisions and policy

Learning Time-Series: Outline


- Part I: Introduction to Time Series
 - Stationarity
 - Trends
- Part II: Models of Time Series
- Part 3: Learning Time Series

Introduction

Johnson & Johnson quarterly earnings per share




Exchange Rates Against the Dollar




Economist.com

House Prices



Consumer Price Index (billion Prices)



...What is new here?

- What you measure today depends on yesterday
 - Could be simple or complicated dependence
 - Memory can be high
 - Memory is typically unknown
- The variation in the data can be due to a time-varying average: trends
 - Linear, quadratic (deterministic)
 - Periodic (seasonal)
- Transformation of the data may help

Stationarity

- Time series need to have some structure
- Stationarity: Some variables are ‘constant’ over time
 - Mean
 - Autocovariance
- What happens if such variables are changing
 - Transform to a stationary series
 - Assume slow variation

Mean and Autocovariance

- Mean of process

$$\mu_t = \mathbf{E}(X_t)$$

- Autocovariance

$$R_X(t_1, t_2) = \mathbf{E}((X_{t_1} - \mu_{t_1})(X_{t_2} - \mu_{t_2}))$$

Stationarity

- Mean of process is constant over time
- Sample Mean (constant over λ)

$$\mathbf{E}(X_t) = \mu$$

$$\hat{\mu} = \frac{1}{N - \lambda} \sum_{i=\lambda}^{N-1} X_i$$

Testing Stationarity: Should not see large variations over λ

Mean and Autocovariance: Stationarity


- Autocovariance is a function of the time difference
- Sample Autocovariance

$$R_X(t, t + \tau) = R_X(\tau)$$



$$\hat{R}_X(\tau) = \frac{1}{N - \lambda} \sum_{i=\lambda}^{N-1} (X_i - \hat{\mu})(X_{i+\tau} - \hat{\mu})$$

Testing Stationarity: Should not see large variations over λ





Testing Stationarity



Demonstrate: 100 samples




Demonstrate: 1000 samples




This file is meant for personal use by jacesca@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Trends in the Consumer Price Index (CPI) data




De-trended



Removing Seasonality

- $X_t = S_t + Y_t$
- Estimate S_t using a periodic regression
- Model the difference $X_t - X_{t-h} = Y_t - Y_{t-h}$



End


Models of time series

Models of Time-Series

- White noise
- Auto-regressive (AR)
- Moving Average (MA)
- Auto-regressive moving average (ARMA)
- Autocovariance of these examples

White noise

- Process $X_t = w$ $\mathbf{E}(w_t) = 0$




- Autocovariance =

$$\mathbf{E}(w_{t_1} w_{t_2}) = \sigma^2 \delta(t_1 - t_2)$$

Random Walk

$$X_t = X_{t-1} + w_t$$





Autoregressive Process AR(p)

- Process





$$X_t = \sum_{i=1}^p a_i X_{t-i} + w_t$$

- w_t is white
- Interpretation

Example AR(1)







AR(2)



Example: AR(2)

$$X_t = 0.7X_{t-1} + 0.1X_{t-2} + w_t$$



Autocovariance of AR(p)

- How do you compute the exact Autocovariance of


$$X_t = \sum_{i=1}^p a_i X_{t-i} + w_t$$

- Autocovariance

$$R_X(\tau) = \sum_{i=1}^p a_i R_X(\tau - i) + \sigma^2 \delta(\tau)$$

- Sum of decaying exponentials!

Geometric Shape of ACF for AR(p)



This file is meant for personal use by jacesca@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.





Moving Average MA(q)

- Process

$$X_t = \sum_{i=0}^q b_i w_{t-i}$$


- w_t is white
- Interpretation

Example MA(2)



Sample vs Actual

$$X_t = w_t + 0.5w_{t-1} - 0.7w_{t-2}$$



This file is meant for personal use by jacesca@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Autocovariance of MA


- How do you compute the exact Autocovariance of

$$X_t = \sum_{i=0}^q b_i w_{t-i}$$

- Finite support

$$R_X(\tau) = \sigma^2 \sum_{j=0}^q b_j b_{j-\tau}$$

Finite lag Shape of ACF for MA







Autoregressive Moving Average Process (ARMA)

- We can combine the two processes

$$X_t = \sum_{i=1}^p a_i X_{t-i} + \sum_{j=0}^q b_j w_{t-j}$$

- Superposition of different processes

Example: ARMA



End

Learning Time Series

Learning a Time Series Model

- Data to Models
 - Autoregressive (AR) models
 - Moving Average (MA)
- AR learning Looks like a standard Least Squares
 - What's the catch?
 - What can we learn?
- How about other models
- Example: Consumer Price Data

Learning AR(p)

- Data

$$X_0, X_1, X_2, \dots X_N$$

- Model (known p)

$$X_t = \sum_{i=1}^p a_i X_{t-i} + w_t$$

- Least squares to estimate

$$(a_1, a_2, \dots a_p)$$

Example

- Generate 100 data points according to

$$x_t = .7 x_{t-1} + .1 x_{t-2} + w_t$$

- Fit an AR(1) model
 - Answer: $a = .8704$
- Fit an AR(2) model
 - Answer $a_1 = 0.7448$ and $a_2 = .1742$
- Observations?

Order Estimation




- Example highlights the importance of order estimation
- Derive multiple estimates and choose based on error on cross validation data
- Add penalty to model complexity (MDL, AIC)
- Use the Autocovariance function (ACF) as a guidance!
 - Can we do better than ACF?

Partial Autocovariance Function (PACF)

- $X_t \mid X_{t+1}, X_{t+2}, \dots, X_{t+k-1} \mid X_{t+k}$
- Project X_{t+1} and X_{t+k} on the variables in between:
 P_t and P_{t+k}
- In the case of ARX, the new PACF is zero outside p
- *Use to estimate order!*




Example: AR(3)

$$X_t = .7 X_{t-1} + .2 X_{t-2} - .1 X_{t-3}$$




Example: AR(3)

$$X_t = -.7 X_{t-1} + .2 X_{t-2} + .1 X_{t-3}$$



Geometric Shape of ACF and PACF for AR(p)



This file is meant for personal use by jacesca@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Learning a MA model

- Data

$$X_0, X_1, X_2, \dots X_N$$




- Model (known p)

$$X_t = \sum_{i=0}^p b_i w_{t-i}$$

- Recall ACF has finite support




Example: MA(2)

$$X_t = .7 W_{t-1} + W_t$$





Example: MA(3)

$$X_t = W_t - .7 W_{t-1} + .2 W_{t-2} + .1 W_{t-3}$$



What about ARMA Models?

$$X_t = .7 X_{t-1} + .2 X_{t-2} + .7 W_t + .2 W_{t-1}$$




ACF vs PACF

	ACF	PACF
AR(p)	decays	zero for $h > p$
MA(q)	zero for $h > q$	decays
ARMA(p,q)	decays	decays

Billion Price Data


- The goal of this problem is to analyze the PriceStats data from the MIT Billion Prices Project, provided by Professor Rigobon.
- Consumer Price Index Data: (consumer price index, the price of a "market basket of consumer goods and services" - a proxy for inflation)

Consumer Price Index




This file is meant for personal use by jacesca@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.


Identify Trends in the CPI data




Empirical Autocovariance



Empirical Partial Autocovariance




AR(1) or AR(2)



This file is meant for personal use by jacesca@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Prediction of AR(2)



Conclusions

- Time Series are everywhere
- Stationarity is important for modeling
- AR, MA, and ARMA are powerful models
 - Estimate the order
 - Compare across models
- Trends

End

Thank You