

Boltzmann Equation for WIMP Dark Matter: Approximate Analytical Solution and Numerical Solution Using Python

Jacinto Paulo Neto

1 Radiation-dominated freeze-out

The evolution of the DM number density n_χ is described by the Boltzmann equation,

$$\frac{dY_\chi(x)}{dx} = -\frac{s(x)\langle\sigma v\rangle}{x H(x)} [Y_\chi^2(x) - Y_\chi^{\text{eq}2}(x)] \quad (1)$$

where $x = m_\chi/T$ is a “time” variable that helps us to simplify the integration and physical interpretations and the equilibrium comoving number density

$$Y_\chi^{\text{eq}}(x) = \frac{45}{4\pi^4} \frac{g_\chi}{g_{\star s}} x^2 K_2(x), \quad (2)$$

g_χ corresponds to the DM degrees of freedom and $K_2(x)$ is the modified Bessel function. In this parametrization, the entropy density and the Hubble parameter are

$$s(x) = \frac{2\pi^2}{45} g_{\star s}(x) m_\chi^3 x^{-3}, \quad (3)$$

$$H(x) = \sqrt{\frac{\pi^2 g_\star(x)}{90}} \frac{m_\chi^2}{M_{\text{P}}} x^{-2}, \quad (4)$$

where $M_{\text{P}} = \sqrt{1/(8\pi G)}$ is the reduced Planck mass. Using these equations we can manipulate the right-handed term,

$$\frac{s(x)}{xH(x)} = \frac{2\pi g_{\star s}(x)}{45} \sqrt{\frac{90}{g_\star(x)}} M_{\text{Pl}} \frac{m_\chi}{x^2}. \quad (5)$$

Hence, by rewriting the Boltzmann equation we obtain

$$\frac{dY_\chi}{dx} = -\frac{\lambda(x)}{x^2} [Y_\chi^2(x) - Y_\chi^{\text{eq}2}(x)], \quad (6)$$

where

$$\lambda(x) = \frac{2\pi g_{\star s}(x)}{45} \sqrt{\frac{90}{g_\star(x)}} M_{\text{Pl}} m_\chi \langle\sigma v\rangle \quad (7)$$

1.1 Approximate DM Yield

Step I Since $n_\chi^{\text{eq}}(x) \propto e^{-x}$, it decreases much faster than $n_\chi(x)$. Therefore, we can safely approximate the Boltzmann equation to

$$\frac{dY_\chi(x)}{dx} \approx -\frac{\lambda(x)}{x^2} Y_\chi^2(x). \quad (8)$$

Step II We can also assume that at the freeze-out time, $m_\chi \gtrsim 10$ GeV, hence $g_\star \approx g_{\star,s} \approx 106.8$. If we assume s-wave thermal cross-section,

$$\langle \sigma v \rangle \simeq a + \frac{3}{2}bx^{-1} + \mathcal{O}(x^{-2}), \quad v^2 = \frac{3}{2}x^{-1}, \quad (9)$$

only the zeroth-order term a contributes to $\lambda(x)$. Therefore, $\lambda(x) \approx \bar{\lambda}$ does not depend on x . Hence, we rewrite again

$$\frac{dY_\chi(x)}{dx} \approx -\frac{\bar{\lambda}}{x^2}Y_\chi^2(x), \quad \bar{\lambda} = \frac{2\pi}{15}\sqrt{10g_\star}M_{\text{Pl}}m_\chi\langle\sigma v\rangle. \quad (10)$$

Step III Here, we change variables $\bar{Y}_\chi \equiv 1/Y_\chi$. This results in

$$\frac{dY_\chi}{dx} = \frac{d}{dx} \left(\frac{1}{\bar{Y}_\chi} \right) = -\frac{1}{\bar{Y}_\chi^2} \frac{d\bar{Y}_\chi}{dx} \quad (11)$$

$$\approx -\frac{\bar{\lambda}}{x^2}Y_\chi^2(x) \quad (12)$$

$$\approx -\frac{\bar{\lambda}}{x^2} \frac{1}{\bar{Y}_\chi^2}. \quad (13)$$

On the right-handed side, we equate the first and the last lines to get

$$\frac{d\bar{Y}_\chi}{dx} = \frac{\bar{\lambda}}{x^2}. \quad (14)$$

Now we have to place boundary conditions on this equation somehow we can match the solution in the asymptotic future to a good approximation at $x \sim 1$. Solving this equation, we have

$$\int_{\bar{Y}_{\chi,f}}^{\bar{Y}_{\chi,\infty}} d\bar{Y}_\chi = \bar{\lambda} \int_{x_f}^{\infty} \frac{1}{x^2}, \quad (15)$$

where the subscript f stands for *freeze-out* time and ∞ is the today asymptotic time. Hence, using $Y_\chi = 1/\bar{Y}_\chi$,

$$\frac{1}{Y_{\chi,\infty}} - \frac{1}{Y_{\chi,f}} = \frac{\bar{\lambda}}{x_f}. \quad (16)$$

Typically, we have $Y_{\chi,f} \gg Y_{\chi,\infty}$. Consequently,

$$Y_{\chi,\infty} \approx \frac{\bar{\lambda}}{x_f}. \quad (17)$$

Using the expression for $\bar{\lambda}$, we got the usual formula for the approximate solution of the Boltzmann equation for freeze-out taking place during a radiation-dominated era,

$$Y_{\chi,\infty} \approx \frac{15}{2\pi\sqrt{10g_\star}} \frac{x_{\text{fo}}}{m_\chi M_{\text{Pl}} \langle \sigma v \rangle}. \quad (18)$$

1.2 Freeze-out temperature

The freeze-out temperature is obtained via

$$\left. \frac{n_{\text{eq}} \langle \sigma v \rangle}{H} \right|_{x=x_f} = 1, \quad (19)$$

where

$$\begin{aligned} n_{\text{eq}} &\approx \frac{g_\chi}{(2\pi)^{3/2}} (m_\chi T)^{3/2} e^{-m_\chi/T} \\ &\approx \frac{g_\chi}{(2\pi)^{3/2}} \frac{m_\chi^3}{x^{3/2}} e^{-x} \end{aligned} \quad (20)$$

Using $H(x)$, s-wave annihilation cross-section, and $x \rightarrow x_f$ we obtain

$$e^{-x_f} \approx \sqrt{\frac{8\pi^5 g_\star}{90}} \frac{1}{g_\chi \langle \sigma v \rangle m_\chi M_{\text{P}} x_f^{1/2}}. \quad (21)$$

Using

$$\frac{8}{90} = \frac{4}{45} = \frac{2^2}{3^2 \times 5}, \quad (22)$$

Finally, we get

$$x_f \approx -\ln \left[\frac{2}{3} \sqrt{\frac{\pi^5 g_\star}{5}} \frac{1}{g_\chi \langle \sigma v \rangle m_\chi M_{\text{P}} x_f^{1/2}} \right] \quad (23)$$

$$\approx \ln \left[\frac{3}{2} \sqrt{\frac{5}{\pi^5 g_\star}} g_\chi \langle \sigma v \rangle m_\chi M_{\text{P}} x_f^{1/2} \right]. \quad (24)$$

2 Numerical solution – Python

```

1 import numpy as np
2 from scipy.integrate import solve_ivp
3 from scipy.special import kn
4 import matplotlib.pyplot as plt
5 from matplotlib import rc
6 rc('font', **{'family': 'serif', 'serif': ['Computer Modern']})
7 rc('text', usetex=True)
8 rc('text.latex', preamble=r'\usepackage{amsmath}')
9
10 # Functions
11 def s(x):
12     return (2 * np.pi**2 / 45) * gstar * mchi**3 * x**-3
13
14 def H(x):
15     return np.sqrt(np.pi**2 * gstar / 90) * mchi**2 / (Mp * x**2)
16
17 def Yeq(x):
18     return (45 / (4 * np.pi**4)) * (gchi / gstar) * x**2 * kn(2, x)
19
20 # Boltzmann equation
21 def boltzmann_eq(x, Y, sigmav):

```

```

22     Yeqx = Yeq(x)
23     return -s(x) * sigmav / (x * H(x)) * (Y**2 - Yeqx**2)
24
25 Markdown:  $\Omega_{\text{DM}} h^2 = 2.74385 \times 10^8 \frac{m_{\text{DM}}}{\text{GeV}} Y_{\text{DM}}$ 
26
27 def YDMPlanck(mchi): # DM in GeV
28     return (0.12/2.74385)*(1e-8/mchi)
29
30 YDMPlanck(100)
31
32 # Constants
33 mchi = 100.0 # GeV
34 Mp = 2.435e18 # GeV
35 gchi = 4; gstar = 106.8
36
37 # Annihilation cross-sections
38 sigmav_values = [1.8e-10, 1.8e-9, 1.8e-8] # GeV-2
39
40 # Initial conditions and integration range
41 xinit = 1e-2; xend = 1e4; Y0 = Yeq(xinit)
42
43 # Solving the differential equation for each cross-section
44 xvalues = np.logspace(np.log10(xinit), np.log10(xend), 1000)
45 solutions = {}
46
47 for sigmav in sigmav_values:
48     sol = solve_ivp(boltzmann_eq, [xinit, xend], [Y0], args=(sigmav,),
49                     dense_output=True, method='BDF', atol=1e-12, rtol=1e-12)
50     Yvalues = sol.sol(xvalues)[0]
51     solutions[sigmav] = Yvalues
52
53 # Plotting the solution
54 fig = plt.figure()
55 fig, ax = plt.subplots(figsize=(7,5), tight_layout=True);
56
57 for sigmav in sigmav_values:
58     if(sigmav == 1.8e-10):
59         plt.plot(xvalues, solutions[sigmav], label=r'$\langle \sigma v \rangle = 10^{-10} \, \text{GeV}^{-2}$')
60     if(sigmav == 1.8e-9):
61         plt.plot(xvalues, solutions[sigmav], label=r'$\langle \sigma v \rangle = 10^{-9} \, \text{GeV}^{-2}$')
62     if(sigmav == 1.8e-8):
63         plt.plot(xvalues, solutions[sigmav], label=r'$\langle \sigma v \rangle = 10^{-8} \, \text{GeV}^{-2}$')
64
65 plt.plot(xvalues, Yeq(xvalues), label=r'$Y_{\chi}(\text{eq})(x)$', linestyle='--',
66         color='black')
67 #plt.axhline(y=YDMPlanck(100), color='r', linestyle='--')
68 fig.text(0.2, 0.32, r"$\Omega_{\text{DM}} h^2 \approx 0.12$", fontsize=12, color='gray')
69 plt.axhspan(YDMPlanck(100)+1.2e-12, YDMPlanck(100)-1.2e-12, facecolor='gray',
70             alpha=0.3)
71 plt.xscale('log')
72 plt.yscale('log')
73 plt.xlabel(r'$x = m_{\chi} / T$', fontsize=14)
74 plt.ylabel(r'$Y_{\chi}(x)$', fontsize=14)
75 plt.xlim(1, 1e4)

```

```

73 plt.ylim(1e-14,1e-1)
74 plt.legend()
75 plt.savefig('WIMPDM-BEQ.pdf')
76 plt.show()

```

Listing 1: Solution of the Boltzmann Equation for WIMP Dark Matter




Figure 1: Numerical solution of the Boltzmann equation for WIMP Dark Matter.