

## GROUP ACTIONS

**DEFINITION:** Let  $G$  be a group and  $X$  be a set. A **group action** of  $G$  on  $X$  is a function  $G \times X \rightarrow X$  typically written as  $(g, x) \mapsto g \cdot x$  such that

- (1)  $g \cdot (h \cdot x) = (gh) \cdot x$  for all  $g, h \in G$  and  $x \in X$ , and
- (2)  $e_G \cdot x = x$  for all  $x \in X$ .

Given a group action of  $G$  on  $X$  and  $x \in X$ , the **orbit** of  $x$  is

$$\text{Orb}_G(x) := \{g \cdot x \mid g \in G\}.$$

**LEMMA:** Given a group action of  $G$  on  $X$ ,

- for  $x, y \in X$ , either  $\text{Orb}_G(x) = \text{Orb}_G(y)$  or  $\text{Orb}_G(x) \cap \text{Orb}_G(y) = \emptyset$ .
- $X = \bigcup_{x \in X} \text{Orb}_G(x)$ .

**DEFINITION:** A group action of  $G$  on  $X$  is

- **transitive** if  $\text{Orb}_G(x) = X$  for some  $x \in X$ .
- **faithful** if  $g \cdot x = x$  for all  $x \in X$  implies that  $g = e$ .

**(1)** Let  $G$  be a group acting on a set  $X$ . For  $x, y \in X$ , write  $x \sim y$  if there exists  $g \in G$  such that  $g \cdot x = y$ .

**(a)** Show that  $\sim$  is an equivalence relation<sup>1</sup>.

**(b)** Relate the previous part to the Lemma.

**(c)** Suppose that  $X$  is a finite set, and  $X_1, \dots, X_\ell$  are the distinct orbits of  $G$  acting on  $X$ . Explain:

$$|X| = \sum_{i=1}^{\ell} |X_i|.$$

**(2)** Dihedral group actions: Let  $D_n$  be the group of symmetries of a regular  $n$ -gon  $P_n$  in  $\mathbb{R}^2$ .

**(a)** Explain why/how  $D_n$  acts naturally on  $P_n$ . Is this action transitive? Is it faithful?

**(b)** Explain why/how  $D_n$  acts naturally on the set of vertices of  $P_n$ . Is this action transitive? Is it faithful?

**(3)** Group actions on  $X \longleftrightarrow$  homomorphisms to  $\text{Perm}(X)$ :

**(a)** Let  $G$  be a group acting on a set  $X$ . For  $g \in G$ , let  $\mu_g : X \rightarrow X$  be the function  $\mu_g(x) = g \cdot x$ , which we made out of the group action. Consider the function

$$\rho : G \rightarrow \text{Perm}(X)$$

$$g \mapsto \mu_g$$

Show that  $\rho$  is a group homomorphism<sup>2</sup>. We call  $\rho$  the **permutation representation** associated to the given group action.

**(b)** Label the vertices of a square counterclockwise by  $\{1, 2, 3, 4\}$ . Write out the induced homomorphism  $D_4 \rightarrow S_4$  coming from the action of  $D_4$  on the vertices as in (2.b) above.

<sup>1</sup>Recall that a relation on a set is an **equivalence relation** if it is *reflexive*, *symmetric*, and *transitive*.

<sup>2</sup>Warning: you should also show that  $\mu_g$  is actually an element of  $\text{Perm}(X)$ . One good way to do this is to show that  $\mu_{g^{-1}}$  is the inverse function of  $\mu_g$ .

- (c) Let  $G$  be a group,  $X$  a set, and  $\rho : G \rightarrow \text{Perm}(X)$  a group homomorphism. Give a natural recipe for a group action of  $G$  on  $X$ , and verify that this is indeed a group action.
- (4) Let  $G$  be a group acting on a set  $X$ . Complete the following sentence, and prove your answer:  
The action of  $G$  on  $X$  is faithful if and only if the associated permutation representation  $\rho : G \rightarrow \text{Perm}(X)$  is \_\_\_\_\_.
- (5) Linear representations on  $K^n \longleftrightarrow$  homomorphisms to  $\text{GL}_n(K)$ :
- (a) Let  $G$  be a group and  $K$  be a field (you can assume  $K = \mathbb{R}$  if you want.) A **linear action** of  $G$  on  $K^n$  is a group action of  $G$  on  $K^n$  such that for each  $g \in G$ , the function  $\mu_g : K^n \rightarrow K^n$  as in (3) is a linear transformation over  $K$ . Given a linear action of  $G$  on  $K^n$ , show that there is natural group homomorphism  $\rho : G \rightarrow \text{GL}_n(K)$ .
- (b) Conversely, given a group homomorphism  $\rho : G \rightarrow \text{GL}_n(K)$ , give a natural recipe for a linear action of  $G$  on  $K^n$ .