

§4.14: NOETHER NORMALIZATION AND ZARISKI'S LEMMA

NOETHER NORMALIZATION: Let K be a field, and R be a finitely-generated K -algebra. Then there exists a finite¹ set of elements $f_1, \dots, f_m \in R$ that are algebraically independent over K such that $K[f_1, \dots, f_m] \subseteq R$ is module-finite; equivalently, there is a module-finite injective K -algebra map from a polynomial ring $K[X_1, \dots, X_m] \hookrightarrow R$. Such a ring S is called a **Noether normalization** for R .

LEMMA: Let A be a ring, and $F \in R := A[X_1, \dots, X_n]$ be a nonzero polynomial. Then there exists an A -algebra automorphism ϕ of R such that $\phi(F)$, viewed as a polynomial in X_n with coefficients in $A[X_1, \dots, X_{n-1}]$, has top degree term aX_n^t for some $a \in A \setminus 0$ and $t \geq 0$.

- If $A = K$ is an infinite field, one can take $\phi(X_n) = X_n$ and $\phi(X_i) = X_i + \lambda_i X_n$ for some $\lambda_1, \dots, \lambda_{n-1} \in K$.
- In general, if the top degree of F (with respect to the standard grading) is D , one can take $\phi(X_n) = X_n$ and $\phi(X_i) = X_i + X_n^{D^{n-i}}$ for $i < n$.

ZARISKI'S LEMMA: An algebra-finite extension of fields is module-finite.

USEFUL VARIATIONS ON NOETHER NORMALIZATION:

- **NN FOR DOMAINS:** Let $A \subseteq R$ be a module-finite inclusion of domains². Then there exists $a \in A \setminus 0$ and $f_1, \dots, f_m \in R[1/a]$ that are algebraically independent over $A[1/a]$ such that $A[1/a][f_1, \dots, f_m] \subseteq R[1/a]$ is module-finite.
- **GRADED NN:** Let K be an infinite field, and R be a standard graded K -algebra. Then there exist algebraically independent elements $L_1, \dots, L_m \in R_1$ such that $K[L_1, \dots, L_m] \subseteq R$ is module-finite.
- **NN FOR POWER SERIES:** Let K be an infinite field, and $R = K[[X_1, \dots, X_n]]/I$. Then there exists a module-finite injection $K[[Y_1, \dots, Y_m]] \hookrightarrow R$ for some power series ring in m variables.

(1) Examples of Noether normalizations: Let K be a field.

- (a)** Show that $K[x, y]$ is a Noether normalization of $R = \frac{K[X, Y, Z]}{(X^3 + Y^3 + Z^3)}$, where x, y are the classes of X and Y in R , respectively.
- (b)** Show that $K[x]$ is *not* a Noether normalization of $R = \frac{K[X, Y]}{(XY)}$. Then show that $K[x + y] \subseteq R$ is a Noether normalization.
- (c)** Show that $K[X^4, Y^4]$ is a Noether normalization for $R = K[X^4, X^3Y, XY^3, Y^4]$.

(2) Use Noether Normalization³ to prove Zariski's Lemma.

¹Possibly empty!

²The assumption that R is a domain is actually not necessary, but can't quite state the general statement yet. We assume that R is a domain so that there is fraction field of R in which to take $R[1/a]$.

³and a suitable fact about integral extensions...

- (3)** Proof of Noether Normalization (using the Lemma): Proceed by induction on the number of generators of R as a K -algebra; write $R = K[r_1, \dots, r_n]$.
- (a)** Deal with the base case $n = 0$.
 - (b)** For the inductive step, first do the case that r_1, \dots, r_n are algebraically independent over K .
 - (c)** Let $\alpha : K[X_1, \dots, X_n] \rightarrow R$ be the K -algebra homomorphism such that $\alpha(X_i) = r_i$, and let ϕ be a K -algebra automorphism of $K[X_1, \dots, X_n]$. Let $r'_i = \alpha(\phi(X_i))$ for each i . Explain⁴ why $R = K[r'_1, \dots, r'_n]$, and for any K -algebra relation F on r_1, \dots, r_n , the polynomial $\phi^{-1}(F)$ is a K -algebra relation on r'_1, \dots, r'_n .
 - (d)** Use the Lemma to find a K -subalgebra R' of R with $n - 1$ generators such that the inclusion $R' \subseteq R$ is module-finite.
 - (e)** Conclude the proof.
- (4)** Proof of Lemma:
- (a)** in the General case: Consider the automorphism ϕ from the general case of the Lemma. Show that for a monomial, we have $\phi(aX_1^{d_1} \cdots X_n^{d_n})$ is a polynomial with unique highest degree term $aX_1^{d_1 D^{n-1} + d_2 D^{n-2} + \cdots + d_n}$. Can two monomials μ, ν in F , have $\phi(\mu)$ and $\phi(\nu)$ with the same highest degree term? Complete the proof.
 - (b)** Prove the Lemma in the infinite field case.
- (5)** Variations on NN.
- (a)** Adapt the proof of NN to show Graded NN.
 - (b)** Adapt the proof of NN to show NN for domains.
 - (c)** Adapt the proof of NN to show NN for power series.

⁴Say α' is the K -algebra map given by $\alpha'(X_i) = r'_i$. Observe that $\alpha' = \alpha \circ \phi$. Why is this surjective?