

Use Centering and Scaling to Improve Numerical Properties

Create a table of population data for the years 1750 - 2000 and plot the data points.

```
year = (1750:25:2000)';  
pop = 1e6*[791 856 978 1050 1262 1544 1650 2532 6122 8170 11560]';  
T = table(year, pop)  
plot(year,pop, 'o')
```


T =

<i>year</i>	<i>pop</i>
<hr/>	<hr/>
1750	7.91e+08
1775	8.56e+08
1800	9.78e+08
1825	1.05e+09
1850	1.262e+09
1875	1.544e+09
1900	1.65e+09
1925	2.532e+09
1950	6.122e+09
1975	8.17e+09
2000	1.156e+10



Use `polyfit` with three outputs to fit a 5th-degree polynomial using centering and scaling, which improves the numerical properties of the problem. `polyfit` centers the data in year at 0 and scales it to have a standard deviation of 1, which avoids an ill-conditioned Vandermonde matrix in the fit calculation.

```
[p,~,mu] = polyfit(T.year, T.pop, 5);  
% Use |polyval| with four inputs to evaluate |p| with the scaled  
% years,  
% |(year-mu(1))/mu(2)|. Plot the results against the original years.  
% mu = (1.87500.0829) 1.0e+03  
f = polyval(p,year,[],mu);  
hold on  
plot(year,f)  
hold off
```



*Copyright 2015 The MathWorks, Inc.
Published with MATLAB® R2016a*