


Modelling RNA Circuits

J. Binysh¹

¹Centre for Complexity Science
University of Warwick


September 25, 2015

The RNA System



- ▶ mRNA produced 'self repressed' - tail folded over the RBS
- ▶ sRNA binds to it - new complex has RBS uncovered.

ODE Model for system (1) - Complex Formation



- Hybridization of sRNA (s) and mRNA (m), into unstable complex ($s : m$) then stable one (c).


$$\frac{ds}{dt} = \frac{N_{\alpha_T}}{f_T} y(t) - (\mu + \delta_s)s - k_{on}sm + k_{off}s : m$$

$$\frac{dm}{dt} = \frac{N_{\alpha_L}}{f_L} x(t) - (\mu + \delta_m)m - k_{on}sm + k_{off}s : m$$

$$\frac{ds : m}{dt} = k_{on}sm - (k_{off} + k_{hyb})s : m - (\mu + \delta_{sm})s : m$$

$$\frac{dc}{dt} = k_{hyb}s : m - (\mu + \delta_c)c$$

ODE Model for system (2) - Translation



- ▶ Translation of stable complex (c) into immature GFP (p).
- ▶ maturation of GFP (g), machine calibration giving measured fluorescence (z).

$$\frac{dp}{dt} = \beta m + f_s \beta c - (\gamma + \mu + \delta_g)p - \frac{v_z p}{K_z + p + g}$$

$$\frac{dg}{dt} = \gamma p - (\mu + \delta_g)g - \frac{v_z g}{K_z + p + g}$$

$$z = z_0 + \frac{g}{\Theta}$$

Full model, with Parameters to be estimated

$$\frac{ds}{dt} = \frac{N_{\alpha_T}}{f_T} y(t) - (\mu + \delta_s)s - k_{\text{on}}sm + k_{\text{off}}s : m \quad (1)$$

$$\frac{dm}{dt} = \frac{N_{\alpha_L}}{f_L} x(t) - (\mu + \delta_m)m - k_{\text{on}}sm + k_{\text{off}}s : m \quad (2)$$

$$\frac{ds : m}{dt} = k_{\text{on}}sm - (k_{\text{off}} + k_{\text{hyb}})s : m - (\mu + \delta_m)s : m \quad (3)$$



$$\frac{dc}{dt} = k_{\text{hyb}}s : m - (\mu + \delta_m)c \quad (4)$$

$$\frac{dp}{dt} = \beta m + f_s \beta c - (\gamma + \mu + \delta_g)p - \frac{v_z p}{K_z + p + g} \quad (5)$$

$$\frac{dg}{dt} = \gamma p - (\mu + \delta_g)g - \frac{v_z g}{K_z + p + g} \quad (6)$$

$$z = z_0 + \frac{g}{\ominus} \quad (7)$$

Recent Experimental Data



- ▶ Single cell fluorescence time series data (Jaramillo Lab, unpublished).
- ▶ Data records $z = z_0 + \frac{g}{\Theta}$

Method for Parameter Estimation in ODE Model

Goal


Estimate unknown parameters in model using time series data.

- ▶ Least squares error minimization approach


$$\arg \min_{\theta} \sum_{i=1}^n (z_{\text{exp,mean}}(t_i) - z(t_i, \theta))^2.$$

- ▶ Used evolutionary algorithm, CMA-ES, to perform minimisation.

Initial Estimation Results




Initial Estimation Results




► Many parameters poorly estimated.

Sensitivity Analysis



- Sensitivity analysis suggests perturbations of several parameters all give similar effects on model output - **hard to resolve**

Model output for all state variables



Full model again

$$\frac{ds}{dt} = \frac{N_{\alpha T}}{f_T} y(t) - (\mu + \delta_s)s - k_{\text{on}}sm + k_{\text{off}}s : m$$

$$\frac{dm}{dt} = \frac{N_{\alpha L}}{f_L} x(t) - (\mu + \delta_m)m - k_{\text{on}}sm + k_{\text{off}}s : m$$

$$\frac{ds : m}{dt} = k_{\text{on}}sm - (k_{\text{off}} + k_{\text{hyb}})s : m - (\mu + \delta_m)s : m$$

$$\frac{dc}{dt} = k_{\text{hyb}}s : m - (\mu + \delta_m)c$$

$$\frac{dp}{dt} = \beta m + f_s \beta c - (\gamma + \mu + \delta_g)p - \frac{v_z p}{K_z + p + g}$$

$$\frac{dg}{dt} = \gamma p - (\mu + \delta_g)g - \frac{v_z g}{K_z + p + g}$$

$$z = z_0 + \frac{g}{\ominus}$$

Simplified Model

$$\frac{ds}{dt} = \frac{N\alpha_T}{f_T} y(t) - (\mu + \delta_s)s - k_{\text{on}}sm + k_{\text{off}}s : m$$

$$\frac{dm}{dt} = \frac{N\alpha_L}{f_L} x(t) - (\mu + \delta_m)m - k_{\text{on}}sm + k_{\text{off}}s : m$$

$$\frac{ds : m}{dt} = k_{\text{on}}sm - (k_{\text{off}} + k_{\text{hyb}})s : m - (\mu + \delta_{sm})s : m$$

$$\frac{dc}{dt} = k_{\text{hyb}}s : m - (\mu + \delta_c)c$$

$$\frac{dp}{dt} = Fy(t) - (\gamma + \mu + \delta_g)p - \frac{v_z p}{K_z + p + g}$$

$$\frac{dg}{dt} = \gamma p - (\mu + \delta_g)g - \frac{v_z g}{K_z + p + g}$$

$$z = z_0 + \frac{g}{\Theta}$$

Simplified Model


$$\frac{dp}{dt} = Fy(t) - (\gamma + \mu + \delta_g)p - \frac{v_z p}{K_z + p + g} \quad (8)$$

$$\frac{dg}{dt} = \gamma p - (\mu + \delta_g)g - \frac{v_z g}{K_z + p + g} \quad (9)$$

$$z = z_0 + \frac{g}{\Theta} \quad (10)$$


- ▶ Model only rate limiting steps - force translation term directly.
- ▶ 3 unknown parameters

Simplified Model Results (1)



- ▶ Error values as low as full model
- ▶ Clearer parameter estimation results
- ▶ Two peak structure due to local minima

Simplified Model Results (2)



- Error landscape, $\Theta = 5400$.

Future Work

- ▶ Fluorescence data **not enough** to estimate all unknown parameters
 - ▶ Further experiments ?
 - ▶ Bounds on unknown parameter values.
- ▶ Improve modelling of translation step
- ▶ Change methodology - Bayesian Methods?