

Preconditioning of Iterative Methods for the Transport Equation

Jack Blake

28th October, 2011

Neutron Transport Equation in Nuclear Fission

Diffusion Synthetic Acceleration (DSA)

An Example

Neutron Transport Equation in Nuclear Fission

Diffusion Synthetic Acceleration (DSA)

An Example

Neutron Transport Equation

Neutron Transport Equation

The mono-energetic, steady-state, linear 3D Boltzmann transport equation:

$$\begin{aligned}\boldsymbol{\Omega} \cdot \nabla \Psi(\mathbf{r}, \boldsymbol{\Omega}) + \sigma(\mathbf{r})\Psi(\mathbf{r}, \boldsymbol{\Omega}) = \\ \frac{1}{4\pi}\sigma_s(\mathbf{r}) \int_{\mathbb{S}^2} \Psi(\mathbf{r}, \boldsymbol{\Omega}') \, d\boldsymbol{\Omega}' \\ + \frac{1}{4\pi}\nu(\mathbf{r})\sigma_f(\mathbf{r}) \int_{\mathbb{S}^2} \Psi(\mathbf{r}, \boldsymbol{\Omega}') \, d\boldsymbol{\Omega}' + \mathcal{Q}(\mathbf{r}, \boldsymbol{\Omega}).\end{aligned}$$

Neutron Transport Equation

The mono-energetic, steady-state, linear 3D Boltzmann transport equation:

$$\begin{aligned}\boldsymbol{\Omega} \cdot \nabla \Psi(\mathbf{r}, \boldsymbol{\Omega}) + \sigma(\mathbf{r})\Psi(\mathbf{r}, \boldsymbol{\Omega}) = \\ \frac{1}{4\pi}\sigma_s(\mathbf{r}) \int_{\mathbb{S}^2} \Psi(\mathbf{r}, \boldsymbol{\Omega}') \, d\boldsymbol{\Omega}' \\ + \frac{1}{4\pi}\nu(\mathbf{r})\sigma_f(\mathbf{r}) \int_{\mathbb{S}^2} \Psi(\mathbf{r}, \boldsymbol{\Omega}') \, d\boldsymbol{\Omega}' + \mathcal{Q}(\mathbf{r}, \boldsymbol{\Omega}).\end{aligned}$$

In 1D:

$$\mu \frac{\partial \Psi}{\partial x} + \sigma \Psi = \frac{1}{2}\sigma_s \int_{-1}^1 \Psi \, d\mu' + \frac{1}{2}\nu\sigma_f \int_{-1}^1 \Psi \, d\mu' + \mathcal{Q}.$$

where σ , σ_s , σ_f and ν are functions of x , the spatial variable, and Ψ and \mathcal{Q} are functions of both x and μ , the angular variable.

Neutron Transport Equation

Neutron Transport Equation

Define:

- ▶ $\mathcal{T}(\cdot) \equiv \mu \frac{\partial(\cdot)}{\partial x} + \sigma(x)(\cdot),$
- ▶ $\mathcal{S}(\cdot) \equiv \frac{1}{2} \sigma_s(x) \int_{-1}^1 (\cdot) \, d\mu',$
- ▶ $\mathcal{F}(\cdot) \equiv \frac{1}{2} \nu(x) \sigma_f(x) \int_{-1}^1 (\cdot) \, d\mu'.$

Neutron Transport Equation

Define:

- ▶ $\mathcal{T}(\cdot) \equiv \mu \frac{\partial(\cdot)}{\partial x} + \sigma(x)(\cdot),$
- ▶ $\mathcal{S}(\cdot) \equiv \frac{1}{2} \sigma_s(x) \int_{-1}^1 (\cdot) d\mu',$
- ▶ $\mathcal{F}(\cdot) \equiv \frac{1}{2} \nu(x) \sigma_f(x) \int_{-1}^1 (\cdot) d\mu'.$

In operator form:

$$\mathcal{T}\Psi = \mathcal{S}\Psi + \mathcal{F}\Psi,$$

where \mathcal{T} , \mathcal{S} and \mathcal{F} are known as the *transport*, *scatter* and *fission* operators respectively.

Neutron Transport Equation

Define:

- ▶ $\mathcal{T}(\cdot) \equiv \mu \frac{\partial(\cdot)}{\partial x} + \sigma(x)(\cdot),$
- ▶ $\mathcal{S}(\cdot) \equiv \frac{1}{2} \sigma_s(x) \int_{-1}^1 (\cdot) d\mu',$
- ▶ $\mathcal{F}(\cdot) \equiv \frac{1}{2} \nu(x) \sigma_f(x) \int_{-1}^1 (\cdot) d\mu'.$

In operator form:

$$\mathcal{T}\Psi = \mathcal{S}\Psi + \mathcal{F}\Psi,$$

where \mathcal{T} , \mathcal{S} and \mathcal{F} are known as the *transport*, *scatter* and *fission* operators respectively.

As an eigenvalue problem:

$$(\mathcal{T} - \mathcal{S})\Psi = \lambda \mathcal{F}\Psi. \quad \textit{Criticality Problem}$$

Inverse Iteration

Inverse Iteration

Let A, B be n -by- n matrices, \mathbf{x} an n -dimensional vector.
Eigenvalue problem:

$$A\mathbf{x} = \lambda B\mathbf{x},$$

Inverse Iteration

Let A, B be n -by- n matrices, \mathbf{x} an n -dimensional vector.
Eigenvalue problem:

$$A\mathbf{x} = \lambda B\mathbf{x}, \quad \Longleftrightarrow \quad (A - \gamma B)\mathbf{x} = (\lambda - \gamma)B\mathbf{x}.$$

Inverse Iteration

Let A, B be n -by- n matrices, \mathbf{x} an n -dimensional vector.
Eigenvalue problem:

$$A\mathbf{x} = \lambda B\mathbf{x}, \quad \Longleftrightarrow \quad (A - \gamma B)\mathbf{x} = (\lambda - \gamma)B\mathbf{x}.$$

Thus

$$(A - \gamma B)^{-1} B\mathbf{x} = \frac{1}{\lambda - \gamma} \mathbf{x}.$$

Inverse Iteration

Let A , B be n -by- n matrices, \mathbf{x} an n -dimensional vector.
Eigenvalue problem:

$$A\mathbf{x} = \lambda B\mathbf{x}, \quad \Longleftrightarrow \quad (A - \gamma B)\mathbf{x} = (\lambda - \gamma)B\mathbf{x}.$$

Thus

$$(A - \gamma B)^{-1} B\mathbf{x} = \frac{1}{\lambda - \gamma} \mathbf{x}.$$

Iteratively we solve a sequence of inverse systems of the form,

$$(A - \gamma B)\mathbf{x} = \text{RHS}.$$

Inverse Iteration

Let A , B be n -by- n matrices, \mathbf{x} an n -dimensional vector.
Eigenvalue problem:

$$A\mathbf{x} = \lambda B\mathbf{x}, \quad \Longleftrightarrow \quad (A - \gamma B)\mathbf{x} = (\lambda - \gamma)B\mathbf{x}.$$

Thus

$$(A - \gamma B)^{-1} B\mathbf{x} = \frac{1}{\lambda - \gamma} \mathbf{x}.$$

Iteratively we solve a sequence of inverse systems of the form,

$$(A - \gamma B)\mathbf{x} = \text{RHS}.$$

In the previous operator form

$$(\mathcal{T} - \mathcal{S} - \gamma \mathcal{F})\Psi = \mathcal{Q}. \quad \text{Source Problem}$$

The Problem

The Problem

The focus of this talk will be efficient iterative solves of

$$(\mathcal{T} - \mathcal{S} - \gamma \mathcal{F})\Psi = \mathcal{Q}.$$

The Problem

The focus of this talk will be efficient iterative solves of

$$(\mathcal{T} - \mathcal{S} - \gamma \mathcal{F})\Psi = \mathcal{Q}.$$

$$(\mathcal{T} - \mathcal{S})\Psi = \mathcal{Q}.$$

Source Problem

The Problem

The focus of this talk will be efficient iterative solves of

$$\begin{aligned}(\mathcal{T} - \mathcal{S} - \gamma \mathcal{F})\Psi &= \mathcal{Q}. \\ (\mathcal{T} - \mathcal{S})\Psi &= \mathcal{Q}. \end{aligned} \quad \textit{Source Problem}$$

- Splitting methods (\rightarrow *Source Iteration*)

The Problem

The focus of this talk will be efficient iterative solves of

$$\begin{aligned}(\mathcal{T} - \mathcal{S} - \gamma \mathcal{F})\Psi &= \mathcal{Q}. \\ (\mathcal{T} - \mathcal{S})\Psi &= \mathcal{Q}. \quad \textit{Source Problem}\end{aligned}$$

- ▶ Splitting methods (\rightarrow *Source Iteration*)
- ▶ Synthetic Acceleration methods (\rightarrow *Diffusion Synthetic Acceleration (DSA)*)


Neutron Transport Equation in Nuclear Fission

Diffusion Synthetic Acceleration (DSA)

An Example

The Structure of a Diffusion Synthetic Acceleration Scheme

$$\phi_0^{(i)} \equiv \frac{1}{2} \int_{-1}^1 \Psi^{(i)} d\mu'$$



Source Iteration

'Splitting' Applied to our problem of interest:

$$A\Psi = Q$$

Source Iteration

'Splitting' Applied to our problem of interest:

$$A\Psi = Q$$

$$(\mathcal{T} - \mathcal{S})\Psi = Q$$

Source Iteration

'Splitting' Applied to our problem of interest:

$$A\Psi = Q$$

$$(\mathcal{T} - \mathcal{S})\Psi = Q$$

$$\mathcal{T}\Psi^{(i+1/2)} = \mathcal{S}\Psi^{(i)} + Q. \quad \textit{Source Iteration}$$

Source Iteration

'Splitting' Applied to our problem of interest:

$$A\Psi = \mathcal{Q}$$

$$(\mathcal{T} - \mathcal{S})\Psi = \mathcal{Q}$$


$$\mathcal{T}\Psi^{(i+1/2)} = \mathcal{S}\Psi^{(i)} + \mathcal{Q}. \quad \textit{Source Iteration}$$

Recall:

- ▶ $\mathcal{T}(\cdot) \equiv \mu \frac{\partial(\cdot)}{\partial x} + \sigma(x)(\cdot),$
- ▶ $\mathcal{S}(\cdot) \equiv \frac{1}{2}(\sigma_s(x) + \nu(x)\sigma_f(x)) \int_{-1}^1 (\cdot) \, d\mu'.$

The Structure of a Diffusion Synthetic Acceleration Scheme

$$\phi_0^{(i)} \equiv \frac{1}{2} \int_{-1}^1 \Psi^{(i)} d\mu'$$



General Synthetic Acceleration

From the Source Iteration we have:

$$\begin{aligned}\mathcal{T}\Psi^{(i+1/2)} &= \mathcal{S}\Psi^{(i)} + \mathcal{Q}. \\ \text{and } \mathcal{T}\Psi &= \mathcal{S}\Psi + \mathcal{Q}.\end{aligned}$$

General Synthetic Acceleration

From the Source Iteration we have:

$$\begin{aligned}\mathcal{T}\Psi^{(i+1/2)} &= \mathcal{S}\Psi^{(i)} + \mathcal{Q}. \\ \text{and } \mathcal{T}\Psi &= \mathcal{S}\Psi + \mathcal{Q}.\end{aligned}$$

Subtracting we obtain an expression for a *linear correction* term

$$\begin{aligned}\mathcal{T}(\Psi - \Psi^{(i+1/2)}) &= \mathcal{S}(\Psi - \Psi^{(i)}), \\ &= \mathcal{S}(\Psi - \Psi^{(i+1/2)}) + \mathcal{S}(\Psi^{(i+1/2)} - \Psi^{(i)}),\end{aligned}$$

General Synthetic Acceleration

From the Source Iteration we have:

$$\begin{aligned}\mathcal{T}\Psi^{(i+1/2)} &= \mathcal{S}\Psi^{(i)} + \mathcal{Q}. \\ \text{and } \mathcal{T}\Psi &= \mathcal{S}\Psi + \mathcal{Q}.\end{aligned}$$

Subtracting we obtain an expression for a *linear correction* term

$$\begin{aligned}\mathcal{T}(\Psi - \Psi^{(i+1/2)}) &= \mathcal{S}(\Psi - \Psi^{(i)}), \\ &= \mathcal{S}(\Psi - \Psi^{(i+1/2)}) + \mathcal{S}(\Psi^{(i+1/2)} - \Psi^{(i)}),\end{aligned}$$

Thus

$$\begin{aligned}(\mathcal{T} - \mathcal{S})(\Psi - \Psi^{(i+1/2)}) &= \mathcal{S}(\Psi^{(i+1/2)} - \Psi^{(i)}), \\ \Rightarrow \Psi &= \Psi^{(i+1/2)} + (\mathcal{T} - \mathcal{S})^{-1}\mathcal{S}(\Psi^{(i+1/2)} - \Psi^{(i)}).\end{aligned}$$

General Synthetic Acceleration

From the Source Iteration we have:

$$\begin{aligned}\mathcal{T}\Psi^{(i+1/2)} &= \mathcal{S}\Psi^{(i)} + \mathcal{Q}. \\ \text{and } \mathcal{T}\Psi &= \mathcal{S}\Psi + \mathcal{Q}.\end{aligned}$$

Subtracting we obtain an expression for a *linear correction* term

$$\begin{aligned}\mathcal{T}(\Psi - \Psi^{(i+1/2)}) &= \mathcal{S}(\Psi - \Psi^{(i)}), \\ &= \mathcal{S}(\Psi - \Psi^{(i+1/2)}) + \mathcal{S}(\Psi^{(i+1/2)} - \Psi^{(i)}),\end{aligned}$$

Thus

$$\begin{aligned}(\mathcal{T} - \mathcal{S})(\Psi - \Psi^{(i+1/2)}) &= \mathcal{S}(\Psi^{(i+1/2)} - \Psi^{(i)}), \\ \Rightarrow \Psi &= \Psi^{(i+1/2)} + \underbrace{(\mathcal{T} - \mathcal{S})^{-1}\mathcal{S}}_M(\Psi^{(i+1/2)} - \Psi^{(i)}).\end{aligned}$$

Diffusion Synthetic Acceleration

Synthetic Acceleration:

$$\Psi = \Psi^{(i+1/2)} + M\mathcal{S}(\Psi^{(i+1/2)} - \Psi^{(i)}).$$

Diffusion Synthetic Acceleration

Synthetic Acceleration:

$$\Psi = \Psi^{(i+1/2)} + M\mathcal{S}(\Psi^{(i+1/2)} - \Psi^{(i)}).$$

Synthetic Acceleration is equivalent to preconditioning...

Diffusion Synthetic Acceleration

Synthetic Acceleration:

$$\Psi = \Psi^{(i+1/2)} + M\mathcal{S}(\Psi^{(i+1/2)} - \Psi^{(i)}).$$

Synthetic Acceleration is equivalent to preconditioning...

DSA is a Synthetic Acceleration method that uses the P_1 diffusion approximation as the choice for M :

$$-\frac{d}{dx} \left(\frac{1}{3\sigma} \frac{d\phi_0}{dx} \right) + \sigma_c \phi_0 = q_0 - \frac{d}{dx} \left(\frac{q_1}{\sigma} \right).$$

The P_1 Diffusion Approximation

The P_1 Diffusion Approximation

Legendre polynomials:

$$P_0(\mu) = 1$$

$$P_n(\mu) = \frac{1}{2^n n!} \frac{d^n}{d\mu^n} (\mu^2 - 1)^n, \quad n = 1, 2, \dots$$

The P_1 Diffusion Approximation

Legendre polynomials:

$$P_0(\mu) = 1$$

$$P_n(\mu) = \frac{1}{2^n n!} \frac{d^n}{d\mu^n} (\mu^2 - 1)^n, \quad n = 1, 2, \dots$$

Orthogonality:

$$\int_{-1}^1 P_n(\mu) P_{\hat{n}}(\mu) d\mu = \frac{2\delta_{n,\hat{n}}}{2n+1}.$$

The P_1 Diffusion Approximation

Legendre polynomials:

$$P_0(\mu) = 1$$
$$P_n(\mu) = \frac{1}{2^n n!} \frac{d^n}{d\mu^n} (\mu^2 - 1)^n, \quad n = 1, 2, \dots$$

Orthogonality:

$$\int_{-1}^1 P_n(\mu) P_{\hat{n}}(\mu) d\mu = \frac{2\delta_{n,\hat{n}}}{2n+1}.$$

Once normalised they form a complete orthonormal sequence, so we can write the neutron flux as an expansion:

$$\Psi(x, \mu) = \sum_{n=0}^{\infty} (2n+1) \phi_n(x) P_n(\mu),$$

where

$$\phi_n(x) \equiv \frac{1}{2} \int_{-1}^1 \Psi(x, \mu') P_n(\mu') d\mu'.$$

The P_1 Diffusion Approximation

The P_N approximation in slab geometry for the Transport equation, consisting of $N + 1$ coupled differential equations is given by

$$\frac{n}{2n+1} \frac{d\phi_{n-1}}{dx}(x) + \frac{n+1}{2n+1} \frac{d\phi_{n+1}}{dx}(x) + (\sigma(x) - \sigma_{s,n}(x)) \phi_n(x) = q_n$$

for $n = 0, \dots, N-1$,

and

$$\frac{N}{2N+1} \frac{d\phi_{N-1}}{dx}(x) + (\sigma - \sigma_{s,N}) \phi_N = q_N.$$

where $q_n \equiv \frac{1}{2} \int_{-1}^1 P_n(\mu) q(x, \mu) d\mu$

and $\sigma_{s,n} \equiv 2\pi \int_{-1}^1 \sigma_s(x, \hat{\mu}) P_n(\hat{\mu}) d\hat{\mu}$

The P_1 Diffusion Approximation

The P_1 Diffusion Approximation

To move from P_N to P_1 set $N = 1$, to obtain:

$$\begin{aligned}\frac{d\phi_1}{dx} + (\sigma - \sigma_{s,0})\phi_0 &= q_0, \\ \frac{1}{3}\frac{d\phi_0}{dx} + (\sigma - \sigma_{s,1})\phi_1 &= q_1.\end{aligned}$$

The P_1 Diffusion Approximation

To move from P_N to P_1 set $N = 1$, to obtain:

$$\begin{aligned}\frac{d\phi_1}{dx} + (\sigma - \sigma_{s,0})\phi_0 &= q_0, \\ \frac{1}{3} \frac{d\phi_0}{dx} + (\sigma - \sigma_{s,1})\phi_1 &= q_1.\end{aligned}$$

Combining these and rearranging yields:

$$-\frac{d}{dx} \left(\frac{1}{3\sigma} \frac{d\phi_0}{dx} \right) + \sigma_c \phi_0 = q_0 - \frac{d}{dx} \left(\frac{q_1}{\sigma} \right).$$

The P_1 Diffusion Approximation

To move from P_N to P_1 set $N = 1$, to obtain:

$$\begin{aligned}\frac{d\phi_1}{dx} + (\sigma - \sigma_{s,0})\phi_0 &= q_0, \\ \frac{1}{3} \frac{d\phi_0}{dx} + (\sigma - \sigma_{s,1})\phi_1 &= q_1.\end{aligned}$$

Combining these and rearranging yields:

$$-\frac{d}{dx} \left(\frac{1}{3\sigma} \frac{d\phi_0}{dx} \right) + \sigma_c \phi_0 = q_0 - \frac{d}{dx} \left(\frac{q_1}{\sigma} \right).$$

Note: ϕ_0 depends only upon x , the spatial variable.

Diffusion Synthetic Acceleration

Linear correction:

$$\mathbf{e}^{(i+1)}(x) = \Psi(x, \mu) - \Psi^{(i+1/2)}(x, \mu),$$

Diffusion Synthetic Acceleration

Linear correction:

$$\mathbf{e}^{(i+1)}(x) = \Psi(x, \mu) - \Psi^{(i+1/2)}(x, \mu),$$

DSA finds an approximation for the integral of this error:

$$\begin{aligned} F^{(i+1)}(x) &\approx \int_{-1}^1 \Psi(x, \mu') - \Psi^{(i+1/2)}(x, \mu') \, d\mu', \\ &= 2\phi_0(x) - 2\phi_0^{(i+1/2)}(x), \end{aligned}$$

Diffusion Synthetic Acceleration

Linear correction:

$$\mathbf{e}^{(i+1)}(x) = \Psi(x, \mu) - \Psi^{(i+1/2)}(x, \mu),$$

DSA finds an approximation for the integral of this error:

$$\begin{aligned} F^{(i+1)}(x) &\approx \int_{-1}^1 \Psi(x, \mu') - \Psi^{(i+1/2)}(x, \mu') \, d\mu', \\ &= 2\phi_0(x) - 2\phi_0^{(i+1/2)}(x), \end{aligned}$$

hence

$$\phi_0^{(i+1)}(x) = \frac{1}{2} F^{(i+1)}(x) - \phi_0^{(i+1/2)}(x).$$

Neutron Transport Equation in Nuclear Fission

Diffusion Synthetic Acceleration (DSA)

An Example

Example Problem


[A. Greenbaum *Iterative methods for solving linear systems*]

water	water	iron	water
$0 \leq x \leq 12$	$12 \leq x \leq 15$	$15 \leq x \leq 21$	$21 \leq x \leq 30$
$\sigma_t = 3.3333$	$\sigma_t = 3.3333$	$\sigma_t = 1.3333$	$\sigma_t = 3.3333$
$\sigma_s = 3.3136$	$\sigma_s = 3.3136$	$\sigma_s = 1.1077$	$\sigma_s = 3.3136$
$f = 1$	$f = 0$	$f = 0$	$f = 0$

Example Problem

[A. Greenbaum *Iterative methods for solving linear systems*]


water	water	iron	water
$0 \leq x \leq 12$	$12 \leq x \leq 15$	$15 \leq x \leq 21$	$21 \leq x \leq 30$
$\sigma_t = 3.3333$	$\sigma_t = 3.3333$	$\sigma_t = 1.3333$	$\sigma_t = 3.3333$
$\sigma_s = 3.3136$	$\sigma_s = 3.3136$	$\sigma_s = 1.1077$	$\sigma_s = 3.3136$
$f = 1$	$f = 0$	$f = 0$	$f = 0$



Example Problem

[A. Greenbaum *Iterative methods for solving linear systems*]

water	water	iron	water
$0 \leq x \leq 12$	$12 \leq x \leq 15$	$15 \leq x \leq 21$	$21 \leq x \leq 30$
$\sigma_t = 3.3333$	$\sigma_t = 3.3333$	$\sigma_t = 1.3333$	$\sigma_t = 3.3333$
$\sigma_s = 3.3136$	$\sigma_s = 3.3136$	$\sigma_s = 1.1077$	$\sigma_s = 3.3136$
$f = 1$	$f = 0$	$f = 0$	$f = 0$



Thank You!