

Does Motivation Matter?

A systematic review and meta-analysis of outcomes following intentional ingestion of foreign objects.

Jack Galbraith-Edge
190662350

Queen Mary, University of London

MSc Emergency & Resuscitation Medicine

July 2025

Abstract Word Count:

Paper Word Count: 3538

Supplement Word Count: 1412

CONTENTS

I	Abstract	2
I-A	Background	2
I-B	Objectives	2
I-C	Methods	2
I-D	Results	2
I-E	Conclusion	2
I-F	Abbreviations	2
I-G	Symbols	2
II	Introduction	2
II-A	Rationale	2
II-B	Objectives	3
II-C	Eligibility Criteria	3
II-D	Information Sources	3
II-E	Data Extraction	3
II-F	Risk of Bias Assessment	3
II-G	Synthesis Methods	4
III	Results	5
III-A	Study Selection	5
III-B	Risk of Bias	5
III-C	Study Characteristics	5
III-C1	Case Reports	5
III-C2	Case Series	5
III-D	Synthesis	7
III-D1	Univariate Association Testing	7
III-D2	Meta-Analysis of Proportions	8
III-D3	Meta-Regression	10
III-E	Discussion	10
III-E1	Intention is poorly reported	10
III-E2	Protest motivation, and intent-to-harm in detention, could modulate outcomes	10
III-E3	Middle-age with differing motivations - reduced surgery but increased mortality	10
III-F	Limitations	11
III-F1	Publication bias, extraordinary cases and countertransference anger	11
III-F2	Selection Bias	11
III-F3	Unmeasured confounding	11
III-F4	Information Bias	11
III-G	Conclusion	11
Appendix A: Methods		18
A-A	Eligibility Criteria	18
A-B	Search Strategy	19
A-B1	PubMed Query	22
A-B2	Embase Query (All Fields)	22
A-B3	CENTRAL (Cochrane) Query (All Fields)	22
A-B4	Web of Science Query	22
A-B5	Scopus Query	22
A-B6	PsycINFO Query	23
A-C	Grey Literature	23
A-D	Screening Process	24
A-E	Data Extraction	24
A-E1	Definitions	24
A-E2	Process	25
A-F	Computational Risk of Bias Assessment	25
A-F1	Case Reports	25
A-F2	Case Series	25

I. ABSTRACT

A. Background

Intentional ingestion of foreign objects (IIFO) is a clinically distinct form of self-harm with increasing prevalence. Whether patient motivation modifies clinical outcomes has not been systematically evaluated.

B. Objectives

To synthesise available evidence on outcomes following IIFO, and determine whether patient motivation, object characteristics, or demographics influence the need for intervention, complications, or mortality.

C. Methods

A comprehensive search of PubMed, Embase, CENTRAL, Web of Science, Scopus, PsycINFO, and Google Scholar identified studies reporting intentional ingestion of non-nutritive foreign bodies in humans.

Studies of any design and age were included if they reported outcomes of endoscopy, surgery, conservative management, complications, or mortality. Exclusions were accidental or substance ingestion, non-English full texts, animal studies, and studies lacking motivation, object, or outcome data.

Screening was conducted by a single reviewer, with 10% checked by a second. Data were extracted at study and case level. Meta-analysis of proportions pooled outcome rates. χ^2 testing and univariate meta-regression explored associations.

D. Results

Seventy-one case reports ($n = 71$) and three case series ($n = 90$) met inclusion. Pooled outcome rates: endoscopy 50%, surgery 30%, conservative management 20%, complications 30%, mortality <1%. Heterogeneity was low to substantial.

Case series populations were exclusively male and detained, with pooled protest motivation $\approx 80\%$ and sharp object ingestion $\approx 90\%$. Meta-regression showed protest motivation significantly reduced odds of surgery (OR = 0.98, 95% CI 0.98–0.98, $p = 0.003$).

In case reports, intent-to-harm was associated with a six-fold increase in surgical intervention (OR = 5.68, 95% CI: 1.43–22.64, $p = 0.020$). Protest and other motivations reduced odds. Adults aged 40–64 had lower surgery rates (OR = 0.19, 95% CI: 0.04–0.78, $p = 0.020$), but two of three deaths occurred in this group, suggesting vulnerability with psychiatric comorbidity. Sharp object ingestion was paradoxically associated with reduced endoscopy, possibly due to bias.

Over 200 studies were excluded for missing motivation or outcome data, highlighting widespread under-reporting.

E. Conclusion

Motivation influences outcomes after IIFO. Protest ingestion in detention may be managed conservatively; intent-to-harm and psychiatric comorbidity increase surgical risk. Standardised prospective reporting is needed.

DATA AVAILABILITY

Data collection, manipulation and analysis in this review were conducted using *Python* [1] in *Visual Studio Code* [2] and *Jupyter Notebooks* [3]. The manuscript was compiled using *LaTeX* [4]. Specific *Python* packages used include: *Pandas* [5], *scikit-learn* [6], *statsmodel* [7], *seaborn* [8], *matplotlib* [9], *Cartopy* [10], *Forestplot* [11].

The data and code used in this systematic review are available on Github at http://github.com/jackgedge/iifo_systematic_review.

ABBREVIATIONS AND SYMBOLS

F. Abbreviations

- BB – Button Battery
- CENTRAL - Cochrane Central Register of Controlled Trials
- CI – Confidence Interval
- HK – Hartung–Knapp adjustment
- IIFO – Intentional Ingestion of Foreign Object(s)
- IQR – Interquartile Range
- REML – Restricted Maximum Likelihood
- OR – Odds Ratio

G. Symbols

- χ^2 – Chi-squared statistic (Cochran's Q)
- τ^2 – Between-study variance

II. INTRODUCTION

A. Rationale

As of May 2024, over 100 million individuals were forcibly displaced worldwide [12]. Refugees and asylum seekers often endure extreme hardship—including violence, trauma, and detention—leading to elevated rates of mental health disorders [13–18].

Among the most alarming manifestations is self-harm, which is up to 216 times more common in offshore detention settings than in the general population [19–21]. Methods vary and include cutting, poisoning, hanging, self-immolation, and intentional ingestion of foreign objects (IIFO) [20, 22, 23].

IIFO – defined as the non-accidental true ingestion of non-nutritive items – is a serious clinical issue, with 10–20% of cases requiring endoscopy and up to 1% needing surgery [24–26]. In displaced populations, delayed access to care increases risks [27].

Rates of intentional ingestion of foreign objects (IIFO) are rising globally and vary across populations. In the United States, incidence nearly doubled between 1997 and 2017, from 3 to 5.3 cases per 100,000 people [28]. Among adults from lower socio-economic backgrounds, up to 92% of ingestions are intentional [28, 29]. In the UK, a forensic mental health study reported 133 IIFO incidents in a single year, involving just 27 patients – equating to one episode every 2.7 days [30].

Complications vary widely. While most involve perforation, gastro-intestinal bleeding, obstruction and mucosal erosion [25], rare incidents of aortoesophageal perforation, fistula and haemothorax [31], cardiac injury [32], liver [33] and pancreatic injury [34] are reported in the literature.

While techniques for foreign body removal have advanced—from early gastroscopy to modern endoscopy—clinical outcomes depend on multiple factors including object characteristics, patient co-morbidities, and timeliness of intervention [25, 35–41].

Category	Details
Population	Any human; any age group.
Interventions or exposures	Non-accidental ingestion of a true foreign body (non-nutritive items).
Comparators / Control group	<p>Demographics: Gender, age, detained person, psychiatric inpatient, displaced person, under influence of alcohol, psychiatric history, severely disabled, previous ingestion.</p> <p>Motivation: Intent to harm, psychiatric, psychosocial, protest, other.</p> <p>Object characteristics: Button battery, magnet, long (>5 cm), large diameter (>2.5 cm), multiple, blunt objects, sharp-pointed objects.</p>
Outcomes of interest	Endoscopic intervention, surgical intervention, conservative management, complication rates, mortality.
Setting	Any setting.
Study designs	Any design.

TABLE I: Inclusion criteria structured using the PICOS framework.

IIFO motivations vary widely. In detention, it may be a form of protest or communication [42, 43]; in psychiatric contexts, it may stem from conditions like psychosis, personality disorders, pica, or malingering [30, 43–48]. In borderline personality disorder, it may function as emotional regulation rather than a suicide attempt [44]. Rare cases of treatment-resistant recurrent IIFO have prompted palliative approaches to care [49].

Despite this, little research explores how these differing motivations affect clinical management and outcomes across different populations [50–52]. Understanding motivation is crucial, as it may influence decisions around intervention [53, 54].

This review aims to examine how motivation shapes IIFO outcomes – specifically rates of endoscopic and surgical intervention, conservative management, complications, and mortality – to better inform treatment in this vulnerable patient group.

B. Objectives

The primary object of this systematic review was to quantify the rates of endoscopy, surgery, death, complication and conservative management following intentional ingestion of foreign objects in human populations. The review sought to examine how individual factors such as demographic/population characteristics, object characteristics and motivations for ingestion influence the likelihood of these outcomes.

Ethical approval was not required as all analysis was based on published data. Eligibility criteria were structured using the PICOS (Population, Intervention, Comparator, Outcome, Studies) framework.

C. Eligibility Criteria

A full list of eligibility criteria is shown in Table I. This is reproduced in a larger format for clarity in Appendix A-A in Table VII. A full list of exclusion criteria is available in Table VIII in Appendix A-A and in the PRISMA diagram shown in Figure 1.

D. Information Sources

Relevant articles were identified through a systematic search of PubMed, Web of Science, Embase, Scopus, PsycINFO, CENTRAL and Google Scholar during January 2025, with the assistance of a librarian. Included articles then had their bibliography's searched by the primary author (JGE) on 14th May 2025 to identify any potential additional literature not uncovered in the primary search. The search was conducted using keywords and MeSH terms based on the concepts underpinning this review. The search queries, keywords and MeSH terms used can be found in Appendix A-B.

All identified articles were collated and duplicate articles were identified and removed. Remaining articles underwent independent title and abstract screening conducted by the first author (JGE). A randomly selected 10% sample of these articles underwent independent screening by a second author (MS). Any discrepancies identified between these two reviewers were resolved by a third reviewer (GC). Inter-reviewer agreement was calculated using Cohen's kappa [55]. Remaining articles proceeded to full text review, where the same independent screening process was repeated on full text articles.

E. Data Extraction

Data were extracted by a single reviewer (JGE) into *Microsoft Excel* [56] and processed in *Python* [1] using *Pandas* [5]. This process is outlined in Appendix A-E.

Data was first extracted from case reports. Predictors were grouped into five subgroups: gender; age group; demographic characteristics, motivation; object characteristics. Ages were grouped into age groups based on clinical relevance. Outcome data were extracted for rates of endoscopy, surgery, conservative management, mortality, and complications. All outcomes were binary and coded per event, rather than per individual. Predictor variables and outcome variables were not mutually exclusive, nor were outcomes. For example, patients, or ingesters – hereafter referred to as the latter – could have multiple outcomes (e.g. endoscopy and surgery) and multiple predictors from each group (e.g. intent-to-harm and psychiatric, and detained and displaced person).

After case report data extraction, data was collapsed and aggregated to form a series. This data was used as a template for case series data extraction to homogenise data and reduce heterogeneity.

Full definitions of all variables (predictors and outcomes) are provided in Appendix A-E. The full dataset of extracted case-level and series-level data (including bias assessments), is available on Github.

F. Risk of Bias Assessment

Risk of bias was assessed manually for all included studies by a single reviewer (JGE), using the *Joanna Briggs Institute (JBI) Critical Appraisal Checklists for Case Reports and Case Series* [57]. Studies were first classified as either case reports or case series based on the level of granularity in the data. Each study was then evaluated using the corresponding JBI tool. A novel computational risk of bias filter was then applied in *Pandas* [5]. That process is outlined in Appendix A-F.

PRISMA 2020 flow diagram for new systematic reviews which included searches of databases, registers and other sources.




Fig. 1: PRISMA flow diagram summarising the study selection process.

G. Synthesis Methods

For case-level associations between binary predictors and outcomes were assessed using χ^2 Test and Fisher's exact test, reporting odds ratios (ORs), 95% confidence intervals, and p-values. Where appropriate, further χ^2 tests were also used to evaluate differences in outcome proportions between groups [55].

Univariate logistic regression was considered but ultimately not used, as the primary aim was to explore associations without assuming a specific functional relationship between predictors and outcomes. Given the binary nature of most predictors and the categorical outcomes, χ^2 tests offered a more transparent and assumption-light approach. This choice also avoided complications from sparse data and convergence issues that can arise with logistic models in small exploratory datasets.

Multivariate logistic regression was not performed due to the limited sample size, high co-linearity between predictors (e.g., overlapping motivations), and the exploratory nature of the analysis.

For series-level data, univariate meta-regression will be conducted to assess associations between binary series-level predictors and pooled outcome proportions, where sufficient data are available. Each predictor will be entered separately to account for incomplete reporting across studies and to avoid over-fitting. Effect estimates were reported as odds ratios (ORs) with REML

estimation [58].

Initially, a meta-analysis of outcome proportions was conducted using a random-effects model to estimate pooled outcome rates across included studies. The random-effects approach was chosen due to the anticipated heterogeneity in study populations, motivations, and object types to analyse the effect of pooling case reports.

REML estimation was used to compute between-study variance (τ^2), while HK adjustments were applied to produce more accurate confidence intervals, particularly in the presence of small sample sizes.

Heterogeneity was quantified using the I^2 statistic, which describes the percentage of total variation across studies that is due to true between-study differences rather than chance. Following the Cochrane Handbook's guidelines, I^2 values were interpreted as follows: 0–40% may not be important; 30–60% may indicate moderate heterogeneity; 50–90% may represent substantial heterogeneity; and 75–100% may reflect considerable heterogeneity [59]. These thresholds are intended as general guidance rather than strict rules and were interpreted in the context of the number of studies, consistency of effect sizes, and confidence interval overlap.

First, meta-analysis was undertaken on case series alone, and then on case series with pooled case report data. This method was given in anticipation of the case reports introducing heterogeneity

into the case series meta-analysis.

Due to the inclusion of primarily case reports and small case series, formal assessment of reporting bias (e.g., via funnel plots or statistical tests for asymmetry) was not feasible.

Confidence in the body of evidence was not formally graded but was considered low to very low due to reliance on uncontrolled observational designs, small sample sizes, and incomplete reporting.

III. RESULTS

A. Study Selection

Details on the screening and selection process are demonstrated in Figure 1. Detail on independent inter-reviewer agreement and third author review can be found in Appendix A-D. Significantly, 135 reports were excluded for failing to explicitly state whether the ingestion report was intentional. Another 91 were excluded for not reporting motivation.

B. Risk of Bias

Case Reports: 75 cases from 67 studies [43, 46, 53, 60–123] were evaluated using the *JBI Checklist for Case Reports* [57]. 3 cases were excluded. Cases were excluded at this stage if they failed to describe the following domains: patient history and timeline (1 case) [96], current patient condition (2 cases) [96], interventions and treatments (1 case) [108], patient post-intervention condition (2 cases) [96], harms (2 cases) [96], and takeaway lessons (2 cases) [96]. The excluded cases came from the following studies: [96, 108]. Of the remaining 71 cases, all reported interventions and treatments (71 cases, 100%) [43, 46, 53, 60–95, 97–107, 109–115, 117–123]. Most clearly described patient history and timeline (70 cases, 99%) [43, 46, 53, 60–73, 75–95, 97–107, 109–115, 117–123], patient post-intervention condition (69 cases, 97%) [43, 46, 53, 60–94, 97, 99–107, 109–115, 117–123], takeaway lessons (69 cases, 97%) [43, 46, 53, 60–73, 75, 77–95, 97–107, 109–115, 117–123], patient demographic (68 cases, 96%) [46, 53, 60–68, 70–95, 97–107, 109–115, 117, 118, 120–123], and current patient condition (68 cases, 96%) [43, 46, 53, 60–79, 81–95, 97–107, 109–115, 117–123]. Reporting was also strong for harms (38 cases, 93%) [63, 64, 68–73, 75, 77, 81, 83, 84, 90, 91, 93, 94, 98, 99, 101–103, 110–115, 117–119, 121, 122], and diagnostic assessments (65 cases, 92%) [43, 46, 53, 60–73, 75–79, 81–85, 87–95, 97–107, 109–115, 117–123].

Case Series: Separately, 3 studies [124–126] were evaluated using the *JBI Checklist for Case Series* [57]. Reporting quality was generally high across all JBI domains. All included case series fully reported clear inclusion criteria, standard condition measurements, valid patient identification methods, complete inclusion, clear demographic information, clear clinical information, clear outcome and follow-up, and appropriate statistical analysis [124–126]. However, fewer studies (2) reported consecutive inclusion, and clear site demographic information [124, 125].

C. Study Characteristics

1) Case Reports: A total of 71 cases were reported 33 countries [43, 46, 53, 60–95, 97–107, 109–115, 117–123].

The top three countries represented were the United States of America ($n = 12$), India ($n = 7$), and the United Kingdom ($n =$




Fig. 2: Meta-Analysis of case series and pooled case reports. Meta-analysis of case series. Prop = Pooled Proportion; Tau² = τ^2 ; N = n; I² = I^2 ; REML and HK estimated 95% Confidence Intervals.

7). The median number of case reports per country was 1 (IQR = 1.0, range 1-12).

Cases were present from a wide age range (7 to 100 years) with a median age of 28 year (IQR = 18). The majority of cases were reported in males (60% vs 39%), with 1 case of unknown gender.

Half of cases had a psychiatric history and over a quarter (26%) had ingested previously. 17% were detained, 6% were psychiatric inpatients, 10% had a severe disability and 3% ($n = 2$) were displaced persons.

Psychiatric motivation was reported in nearly half of cases (48%), with intent-to-harm and psychosocial reported in 29% and 23% of cases respectively. Protest was reported in 11% of cases and other in 12%.

Most ingestions involved large (> 2.5cm) diameter (72%), sharp (62%) objects and multiple object ingestion (62%). Almost 50% of cases (48%) involved sharp and sharp (45%) object ingestion. Fewer ingestions involved magnets (12%) and button batteries (2%).

Complication rates were high (66%), as were rates of endoscopy and surgery (61% and 44% respectively). Cases were only managed conservatively 10% of the time. The mortality rate in case reports was 2.8%.

A table of case-level characteristics is shown in Table II

2) Case Series: 3 studies were case series, yielding 90 cases [124–126]. Case series were present from the United States of America [125] ($n = 19$), South Korea [126] ($n = 52$) and Tunisia [124] ($n = 19$).

Values reported herein are mean averages across all case series. Unreported variables are treated as 0.

All cases were male, aged 17–50 years and detained 33% had a psychiatric history (range 82–95%) - no psychiatric inpatients.

Demographic predictors were poorly recorded. Previous ingestion rates were only reported in one series at 11%. There was no

TABLE II: Case-level summary statistics.

Variable	Count	Percentage	References
<i>Age</i>			
Max	100		[97]
Mean	30		
Median	28		
IQR	18		
Min	7		[98, 103]
<i>Age Group</i>			
Adult (25–39)	25	35%	[53, 62, 65, 66, 69, 71, 76, 77, 79, 82–84, 86, 89, 95, 99, 101, 105–107, 111, 115, 119, 121, 122]
Young Adult (18–24)	17	24%	[43, 60, 63, 67, 72, 78, 90, 91, 93, 99, 100, 104, 109, 120]
Child/Adolescent (<18)	13	18%	[61, 64, 75, 80, 85, 98, 102, 103, 112, 113, 118]
Middle-Aged (40–64)	11	16%	[46, 73, 74, 81, 87, 88, 94, 110, 114, 117, 123]
Older Adult (65+)	3	4%	[68, 92, 97]
<i>Gender</i>			
Male	43	61%	[43, 46, 60, 62, 63, 65–67, 69, 70, 73, 74, 78, 81–84, 86, 88, 89, 93–95, 98–101, 105, 106, 109, 111–115, 117, 120, 123]
Female	27	38%	[53, 61, 64, 68, 71, 72, 75–77, 79, 80, 85, 87, 90–92, 97, 102–104, 107, 110, 118, 119, 121]
Unknown	1	1%	[122]
<i>Demographic</i>			
Psychiatric History	36	51%	[53, 61, 62, 64, 66–68, 74, 76, 79, 80, 82, 83, 86–90, 92–95, 98, 100, 101, 103, 104, 107, 110–112, 121–123]
Prior Ingestion	19	27%	[62, 66, 70, 72, 78, 80, 81, 86, 88, 89, 98, 107, 112, 114, 121–123]
Detained	12	17%	[43, 62, 65, 66, 99, 105, 111, 115]
Severe Disability	7	10%	[67, 92, 98, 103, 104, 121, 123]
Psychiatric Inpatient	4	6%	[80, 122, 123]
Alcohol Influence	3	4%	[69, 78, 114]
Displaced Person	2	3%	[60, 84]
<i>Motivation</i>			
Psychiatric	34	48%	[46, 53, 62, 64, 66, 67, 71, 72, 79–82, 86–95, 97, 98, 101, 103, 107, 110–112, 120, 123]
Intent-to-Harm	21	30%	[43, 46, 61–63, 74, 76–78, 83, 97, 99, 100, 107, 111, 112, 122]
Psychosocial	16	22%	[60, 69, 72, 75, 85, 87, 93, 97, 102, 105, 106, 109, 113, 114, 118, 119]
Other	9	13%	[64, 65, 81, 86, 104, 107, 115, 117, 121]
Protest	8	11%	[43, 73, 84, 99]
<i>Object</i>			
Large (>2.5cm) Diameter	51	72%	[43, 46, 60–63, 65–67, 70, 71, 74–78, 80, 81, 84, 86, 88, 89, 91–95, 99–107, 110, 112, 114, 115, 119, 121–123]
Multiple	44	62%	[53, 64, 66–68, 72–75, 81–83, 85–91, 93–95, 97–103, 109–114, 117, 118, 120, 122, 123]
Sharp	34	48%	[43, 53, 61, 62, 66, 69, 71, 72, 74, 78–83, 86–89, 91, 93, 95, 99–101, 109, 120, 123]
Long (>5cm)	32	45%	[46, 61, 63, 65, 67, 71, 74, 76–78, 80, 81, 83, 84, 89, 91–93, 95, 100, 101, 103, 105, 107, 110, 114, 115, 120, 121, 123]
Magnet	9	13%	[64, 73, 75, 98, 102, 103, 112, 113, 118]
Button Battery	2	3%	[70, 73]
<i>Outcome</i>			
Complication	47	66%	
Surgery	43	61%	
Endoscopy	31	44%	
Conservative	7	10%	
Death	2	3%	

severe disability (not reporting in two series). Data on displaced person and alcohol influence were not reported at all.

Motivations were predominantly protest (78%, range 16–97%); psychiatric in 13% (63.2% in one series, 0% in the other two); intent-to-harm in 7% (range 0–21%). There were no reports of

other motivation or psychosocial motivation.

One series only report sharp object ingestion and no other object characteristics [125]. In the other two series, overall ingestion involved a sharp object in most cases (76%, range 64–100%). Long objects were ingested 37% of cases (36%, 0–67%),

multiple objects were ingested in (27.8%, range 0-67%)

In terms of outcomes, complications were only reported in one series [126], data was absent from the other two. Endoscopy occurring in 52% (range 5-89%), although it was unreported in one series [125]. Surgery occurred following 17% of ingestions (range 12-26%), conservative management following 32% (0% in one case and above 70% in the other two).

A full list of grouped series-level characteristics and outcomes is available in Table IV.




Fig. 3: Heat map of all cases per country.

D. Synthesis

1) Univariate Association Testing: In case reports, middle-age (40–64 years) was associated with significantly reduced odds of surgery ($OR = 0.19$, 95% CI: 0.04–0.78, $p = 0.020$). There were non-significant trends towards increased of endoscopy ($OR = 2.62$, 95% CI: 0.69–9.95, $p=0.192$) and conservative management ($OR = 2.44$, 95% CI: 0.41–14.57, $p=0.295$)

Further analysis of this subgroup ($n = 11$ vs $n = 60$) revealed a significantly higher proportion of males and individuals ingesting multiple objects (+25% and +8%, respectively). There was a non-significant tendency toward higher rates of psychiatric motivation (+13.5%) and history of prior ingestion (+9.3%). Conversely, fewer individuals in this age group were detained, and intent-to-harm motivation appeared less common.

Two deaths occurred in this age group, outside of a detention context: one from metal ingestion in the context of pica in schizophrenia [94] and another from drug-induced acuphagia [81].

There were no other observed statistical relationships in any of the age groups.

Intent-to-harm motivation was associated with increased odds of surgery ($OR = 5.68$, 95% CI: 1.43–22.64, $p = 0.020$). This subgroup ($n = 21$ vs $n = 50$) included significantly more young adults (+20%), males (+16%), and individuals ingesting large-diameter objects (+6%) and sharp objects (+4%). A significant reduction in psychosocial co-motivation was observed in this group (-25%).

The presence of other motivation was associated with significantly reduced odds of surgery ($OR = 0.15$, 95% CI: 0.03–0.77, $p = 0.024$). There were non significant increases in the odds of conservative management ($OR = 1.17$, 95% CI: 0.12, 10.99, $p=1.000$), endoscopy ($OR = 2.96$, 95% CI: 0.68–12.95, $p=0.165$), death ($OR = 7.62$, 95% CI: 0.43–134.24, $p=0.239$)

TABLE III: Case counts per country.

Country	Case Count	%	References
South Korea	52	32.3%	[126]
United States of America	31	19.3%	[53, 62, 73, 83, 86–88, 92, 94, 98, 111, 113, 125]
Tunisia	19	11.8%	[124]
United Kingdom	7	4.3%	[68, 70, 75, 77, 84, 105]
India	7	4.3%	[71, 72, 90, 91, 95, 101, 117]
Bulgaria	6	3.7%	[43, 99]
Iran	5	3.1%	[80–82]
Turkey	4	2.5%	[60, 67, 112, 121]
Spain	2	1.2%	[74, 122]
Poland	2	1.2%	[93, 119]
China	2	1.2%	[89, 97]
Pakistan	1	0.6%	[120]
Switzerland	1	0.6%	[118]
Taiwan	1	0.6%	[76]
South Africa	1	0.6%	[109]
Saudi Arabia	1	0.6%	[110]
Qatar	1	0.6%	[63]
Portugal	1	0.6%	[104]
Sweden	1	0.6%	[102]
Australia	1	0.6%	[66]
Oman	1	0.6%	[61]
Netherlands	1	0.6%	[69]
United Arab Emirates	1	0.6%	[64]
Japan	1	0.6%	[103]
Italy	1	0.6%	[106]
Israel	1	0.6%	[85]
Iraq	1	0.6%	[46]
Hungary	1	0.6%	[78]
Greece	1	0.6%	[107]
Germany	1	0.6%	[123]
Ethiopia	1	0.6%	[100]
Egypt	1	0.6%	[65]
Ecuador	1	0.6%	[79]
Croatia	1	0.6%	[115]
Nepal	1	0.6%	[114]

Note: % rounded to one decimal place.

One death occurred in this subgroup [81], from 3,4-methylenedioxymethamphetamine (MDMA) induced acuphagia as the other motivation.

Although no statistically significant demographic differences were observed in this subgroup ($n = 9$ vs $n = 62$), there was a non-significant tendency toward more adults (+22%), individuals with prior ingestion history (+9%), those ingesting long objects (+9%), and those with severe disability (+8%). This group also showed lower prevalence of intent-to-harm motivation (-20%), fewer sharp object ingestions (-10%), and reduced psychiatric history and psychiatric motivation. This suggests a diverse subgroup.

Sharp object ingestion was associated with decreased odds of endoscopy. Subgroup analysis ($n = 34$ vs $n = 37$) revealed a significant reduction in large-diameter object ingestion (-11%) and

TABLE IV: Grouped series-level summary.

Variable	Average	Karp <i>et al.</i> (1991) [125]	Lee <i>et al.</i> (2007) [126]	Elghali <i>et al.</i> (2016) [124]
Total Cases	90	19	52	19
Gender				
Male	90 (100%)	19 (100%)	52 (100%)	19 (100%)
Female	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Unknown	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Demographic				
Detained	90 (100%)	19 (100%)	52 (100%)	19 (100%)
Psychiatric History	30 (33%)	18 (95%)	9 (18%)	2 (12%)
Prior Ingestion	2 (2%)	—	—	2 (10%)
Psychiatric Inpatient	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Displaced Person	0 (0%)	—	—	—
Alcohol Influence	0 (0%)	—	—	—
Severe Disability	0 (0%)	—	0 (0%)	—
Motivation				
Protest	70 (78%)	3 (16%)	50 (97%)	17 (90%)
Psychiatric	12 (13%)	12 (63%)	0 (0%)	0 (0%)
Intent-to-Harm	6 (7%)	4 (21%)	0 (0%)	2 (10%)
Psychosocial	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Other	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Object				
Sharp	68 (76%)	19 (100%)	33 (64%)	16 (84%)
Long (>5cm)	32 (36%)	—	32 (62%)	0 (0%)
Multiple	25 (28%)	—	24 (46%)	1 (5%)
Button Battery	0 (0%)	—	0 (0%)	0 (0%)
Magnet	0 (0%)	—	0 (0%)	0 (0%)
Large (>2.5cm) Diameter	0 (0%)	—	—	—
Outcome				
Endoscopy	47 (52%)	—	46 (88%)	1 (5%)
Conservative	29 (32%)	14 (74%)	0 (0%)	15 (79%)
Surgery	15 (17%)	5 (26%)	6 (12%)	4 (21%)
Complication	6 (7%)	—	6 (12%)	—
Death	1 (1%)	0 (0%)	0 (0%)	1 (5%)
Age				
Mean	24	24	—	24
Min	17	17	25	19
Median	35	—	35	—
Max	50	40	50	27

Key: n (%). Average = predictor and outcome mean average rates across series.

magnet ingestion (~13%). There was a non-significant tendency to towards more conservative management (OR 1.51, 95% CI: 0.31-7.30, $p=0.703$) and surgery (OR 2.27 95% CI: 0.85-6.06, $p=0.157$).

All subgroup comparisons were conducted within the overall case-level dataset ($n = 71$). While observed differences may reflect genuine patterns in clinical behavior, they should be interpreted with caution due to the small sample sizes and potential for unmeasured confounding.

Subgroup analysis of predictors significantly associated with outcomes are shown graphically in Figure 5.

2) *Meta-Analysis of Proportions:* Pooled predictors and outcomes rates for the three included case series were first examined using meta-analysis of proportions, employing REML estimation with HK-adjusted 95% CIs. Results are presented in Figure 4.

All patients in the included case series were male and detained,

resulting in a demographically homogenous cohort with respect to gender and detention status.

The pooled prevalence of psychiatric history was ≈40% (range: 0.1-99.8% across three studies), with substantial heterogeneity ($I^2 = 68\%$, $\tau^2 = 5.0$). Motivational subgroups showed moderate to high heterogeneity ($I^2 = 33\%-69\%$). Protest motivation was reported in ≈78% of patients (range: 0-100%), with moderate heterogeneity ($I^2 = 32.7\%$, $\tau^2 = 4.3$). Intent-to-harm motivation was present in ≈4% of cases (range: 0-98%), with moderate heterogeneity ($I^2 = 40\%$, $\tau^2 = 4.0$). Psychiatric motivation was reported in ≈2% of cases (range: 0-100%), with high heterogeneity ($I^2 = 69\%$, $\tau^2: 3.6$).

Meta-analysis was not feasible for several predictors—namely, displaced person status, alcohol use, severe disability, history of prior ingestion, large-diameter objects, and complications—as each was reported in only a single study. Similarly, meta-analysis

TABLE V: Univariate associations between predictors and outcomes.

Variable	Conservative	Endoscopy	Surgery	Death	Complication
<i>Gender</i>					
Male	0.22 [0.04, 1.25] (p=0.104)	1.34 [0.51, 3.53] (p=0.722)	0.99 [0.37, 2.62] (p=1.000)	—	1.49 [0.55, 4.06] (p=0.595)
Female	4.77 [0.86, 26.63] (p=0.097)	0.82 [0.31, 2.18] (p=0.887)	0.92 [0.34, 2.44] (p=1.000)	—	0.79 [0.29, 2.17] (p=0.847)
<i>Age Group</i>					
Child/Adolescent (<18)	1.93 [0.33, 11.24] (p=0.604)	0.32 [0.08, 1.29] (p=0.178)	2.53 [0.63, 10.15] (p=0.307)	—	1.89 [0.47, 7.65] (p=0.521)
Young Adult (18–24)	1.31 [0.23, 7.44] (p=0.670)	0.31 [0.09, 1.06] (p=0.101)	2.60 [0.75, 9.01] (p=0.210)	—	1.30 [0.40, 4.25] (p=0.885)
Adult (25–39)	0.28 [0.03, 2.45] (p=0.409)	2.17 [0.81, 5.85] (p=0.195)	0.96 [0.36, 2.61] (p=1.000)	—	0.86 [0.31, 2.39] (p=0.979)
Middle-Aged (40–64)	2.44 [0.41, 14.57] (p=0.295)	2.62 [0.69, 9.95] (p=0.192)	0.19 [0.04, 0.78] (p=0.020)*	—	0.56 [0.15, 2.05] (p=0.490)
Older Adult (65+)	—	0.63 [0.05, 7.32] (p=1.000)	1.32 [0.11, 15.25] (p=1.000)	—	1.02 [0.09, 11.88] (p=1.000)
<i>Demographic</i>					
Detained	—	0.96 [0.27, 3.42] (p=1.000)	0.89 [0.25, 3.18] (p=1.000)	—	0.64 [0.18, 2.32] (p=0.515)
Psychiatric Inpatient	—	0.43 [0.04, 4.40] (p=0.630)	2.32 [0.23, 23.75] (p=0.630)	—	0.15 [0.01, 1.54] (p=0.108)
Displaced Person	—	—	0.67 [0.03, 12.84] (p=1.000)	—	0.50 [0.03, 9.77] (p=1.000)
Alcohol Influence	—	0.83 [0.07, 10.20] (p=1.000)	1.05 [0.09, 12.88] (p=1.000)	—	—
Psychiatric History	0.92 [0.19, 4.51] (p=1.000)	0.74 [0.27, 2.06] (p=0.749)	1.45 [0.52, 4.07] (p=0.657)	0.69 [0.04, 11.51] (p=1.000)	0.74 [0.25, 2.17] (p=0.780)
Severe Disability	—	4.02 [0.72, 22.47] (p=0.120)	0.83 [0.17, 4.04] (p=1.000)	—	1.35 [0.24, 7.54] (p=1.000)
Prior Ingestion	1.62 [0.29, 9.05] (p=0.669)	0.78 [0.24, 2.50] (p=0.902)	0.78 [0.24, 2.51] (p=0.911)	1.56 [0.09, 26.48] (p=1.000)	0.36 [0.10, 1.29] (p=0.203)
<i>Motivation</i>					
Protest	—	0.39 [0.07, 2.09] (p=0.447)	4.81 [0.55, 41.92] (p=0.239)	—	4.12 [0.47, 35.95] (p=0.247)
Psychiatric	6.86 [0.78, 60.48] (p=0.105)	1.37 [0.52, 3.61] (p=0.699)	0.49 [0.18, 1.33] (p=0.248)	—	0.70 [0.25, 1.94] (p=0.670)
Psychosocial	1.29 [0.22, 7.37] (p=1.000)	0.99 [0.32, 3.08] (p=1.000)	0.56 [0.18, 1.75] (p=0.482)	—	0.86 [0.27, 2.76] (p=1.000)
Intent-to-Harm	—	0.47 [0.15, 1.48] (p=0.307)	5.68 [1.43, 22.64] (p=0.020)*	—	0.88 [0.29, 2.67] (p=1.000)
Other	1.17 [0.12, 10.99] (p=1.000)	2.96 [0.68, 12.95] (p=0.165)	0.15 [0.03, 0.77] (p=0.024)*	7.62 [0.43, 134.24] (p=0.239)	0.60 [0.14, 2.46] (p=0.475)
<i>Object</i>					
Sharp	1.51 [0.31, 7.30] (p=0.703)	0.32 [0.12, 0.85] (p=0.037)*	2.27 [0.85, 6.06] (p=0.157)	1.09 [0.07, 18.15] (p=1.000)	1.13 [0.42, 3.04] (p=1.000)
Long (>5cm)	0.44 [0.08, 2.44] (p=0.442)	0.76 [0.29, 1.97] (p=0.746)	2.56 [0.94, 6.94] (p=0.106)	1.19 [0.07, 19.88] (p=1.000)	1.96 [0.70, 5.49] (p=0.304)
Multiple	4.11 [0.47, 36.14] (p=0.240)	0.46 [0.17, 1.21] (p=0.181)	1.09 [0.41, 2.91] (p=1.000)	—	2.79 [1.01, 7.71] (p=0.081)
Button Battery	—	—	—	—	—
Magnet	—	1.04 [0.25, 4.24] (p=1.000)	2.53 [0.49, 13.17] (p=0.467)	—	—
Large (>2.5cm) Diameter	0.22 [0.04, 1.10] (p=0.070)	1.29 [0.43, 3.86] (p=0.857)	1.83 [0.62, 5.44] (p=0.413)	—	1.00 [0.32, 3.13] (p=1.000)

OR: Odds Ratio; CI: Confidence Interval; p: p-value. * indicates $p < 0.05$. Bold = statistically significant. — = missing or unstable estimate.

of complication rates was not possible due to insufficient data.

Fig 2.

Cases from case reports were collapsed and pooled for meta-analysis alongside the case series data. The results are shown in

Substantial heterogeneity was introduced in this process. Heterogeneity in detention status rose ($I^2 = 69\%$, $\tau^2 = 5$), is did

TABLE VI: Univariate meta-regression results (series-level).

Variable	Surgery
<i>Demographic</i>	
Detained	0.97 [0.96, 0.99] ($p=0.157$)
Psychiatric History	1.03 [0.89, 1.18] ($p=0.771$)
<i>Gender</i>	
Male	0.97 [0.96, 0.99] ($p=0.157$)
<i>Motivation</i>	
Intent-to-Harm	1.29 [1.16, 1.44] ($p=0.133$)
Protest	0.98 [0.98, 0.98] ($p=0.003$)*
Psychiatric	1.06 [0.96, 1.18] ($p=0.448$)
<i>Object</i>	
Long (>5cm)	—
Multiple	—
Sharp	0.95 [0.91, 0.99] ($p=0.250$)

OR: Odds Ratio; CI = 95% Confidence Interval.
* indicates $p < 0.05$; — = missing or unstable estimate.

gender heterogeneity.

Heterogeneity was only low in; psychiatric inpatient pooled proportions between 0–10% ($I^2 = 10$, $\tau^2 = 3$); other motivation pooled proportion between 0–30% ($I^2 = 13$, $\tau^2 = 4.8$); and surgery pooled proportion of 10–60% (($I^2 = 17.5$, $\tau^2 = 0.8$).

These findings should be interpreted cautiously, as the wide confidence intervals and substantial heterogeneity likely reflect limited and inconsistent reporting across the small number of included studies.

3) *Meta-Regression:* Univariate meta-regression revealed exploratory associations between several predictors and the likelihood of surgical intervention across the three included studies. With this heterogeneity in mind, it was decided not to pool case series for univariate meta-regression, to examine detained case series as a subgroup against the more heterogeneous case reports.

In detained populations, protest motivation was associated with a significantly lower likelihood of surgery (OR = 0.98, 95% CI: 0.98–0.98, $p = 0.003$). However, this result must be interpreted with caution due to the extremely small standard error and the limited number of studies. It is possible that this reflects quasi-complete separation (i.e., the outcome is nearly perfectly predicted by one variable) or instability in the regression estimates due to sparse data or model instability rather than a reliable effect.

Intent-to-harm motivation showed a positive association with surgery (OR = 1.29, 95% CI: 1.16–1.44, $p = 0.133$), though this was not statistically significant.

Male gender and prisoner status were both associated with a non-significant trend toward lower odds of surgery (OR = 0.97, 95% CI: 0.96–0.99, $p = 0.157$ for both).

Presence of a sharp object also trended toward reduced odds of surgery (OR = 0.95, 95% CI: 0.91–0.99, $p = 0.250$), but this was not statistically significant.

Neither psychiatric motivation (OR = 1.06, 95% CI: 0.96–1.18, $p = 0.448$) nor a history of psychiatric illness (OR = 1.03, 95% CI: 0.89–1.18, $p = 0.771$) showed a meaningful association with surgical intervention.

Given the limited number of series ($n = 3$), these findings should be regarded as exploratory, and interpreted with caution.

E. Discussion

1) *Intention is poorly reported:* A total of 135 reports were excluded for not reporting intention and 91 were excluded for not reporting outcomes. The absence of these key variables meant that these studies could not be meaningfully integrated into the analysis. Collectively, these exclusions represent a substantial portion of the available literature and could have more than doubled the total sample size of this review. Their omission not only limits statistical power but also reflects a broader issue in the literature – namely, inconsistent or incomplete documentation of psychosocial factors and clinical outcomes. This under-reporting may introduce bias and restrict the generalisability of findings, particularly regarding the relationship between patient motivations and clinical decision-making.

2) *Protest motivation, and intent-to-harm in detention, could modulate outcomes:* Outcomes among detained individuals differed markedly across motivations. In case reports involving lesser-detained ingestors ($n = 71$), the pooled surgery rate was high (58%), with intent-to-harm as the dominant motivation (80%) and protest present in 55%. Sharp object ingestion was common (67%). Intent-to-harm was associated with nearly a six-fold increase in odds of surgery.

However, case series involving exclusively detained men ($n = 90$) in three countries revealed a contrasting trend. Despite pooled sharp object ingestion rates of approximately 90%, protest motivation was associated with an 80% reduction in odds of surgery. This inverse association emerged despite moderate heterogeneity and wide confidence intervals.

In the largest series ($n = 52$) [126], protest motivation was nearly universal (97%), intent-to-harm was absent, and 64% of objects were sharp, yet the surgery rate was only 12%. In contrast, an older series ($n = 19$) [125] reported lower protest motivation (16%) and higher intent-to-harm (12%), with a surgery rate of 26%. Endoscopy rates were not reported in that series. A third series published in 2016 [124]—falling chronologically between the two—reported protest in 90%, sharp object ingestion in 84%, and an intent-to-harm rate of 10%, with a surgery rate of 21%, 79% managed conservatively, and one death.

These findings suggest that motivational context may shape clinical decisions, even when object risk is high. However, publication bias in case reports cohort likely inflates surgery rates. For example, a Bulgarian series [43, 99] – reported in this review as cases due to the level of detail – described five inmates who ingested a total of 20 metal crosses – all required laparotomy for perforation.

3) *Middle-age with differing motivations - reduced surgery but increased mortality:* In case reports, individuals in the middle-aged group demonstrated a non-significant tendency towards reduced ingestion of sharp and large-diameter objects (~5% for both) and lower rates of intent-to-harm motivation. As intent-to-harm was associated with increased odds of surgery in this review, its lower prevalence may partly explain reduced surgical intervention in this group.

However, this apparent protective trend was offset by increased mortality. Two deaths occurred among middle-aged individuals, both outside custodial settings but with psychiatric motivations and history. One case involved a 54-year-old man with a 15-year history of repeated ingestions, who ultimately died after refusing surgery multiple times [94]. The other involved a patient who died

after ingesting metal as a result of acuphagia linked to MDMA use [81].

This duality suggests middle age may represent a complex and vulnerable subgroup. On one side is a cohort with significant psychiatric comorbidities—reflected in higher recurrence—who may present late or decline treatment, thereby increasing mortality risk. On the other, a lower prevalence of intent-to-harm may contribute to reduced rates of surgical intervention.




Fig. 4: Meta-Analysis of case series. Prop = Pooled Proportion; Tau² = τ^2 ; N = n; I² = I^2 ; REML and HK estimated 95% Confidence Intervals.

F. Limitations

1) *Publication bias, extraordinary cases and countertransference anger:* Both of the above effects could be explained by publication bias, were deaths was published as they were exemplary and extraordinary cases.

This is demonstrated in the case reports. Authors appear to enjoy publishing extraordinary cases with sarcastic and sometimes offensive titles. Examples include: “Loose Screws”, in reference to a female with an extensive psychiatric history who (“as advised by her rastafarian”) ingested nails in a milkshake [87]; *Ingested Magnets: the force within* [113]; *Now You See It, Endo You Don’t: Case of the Disappearing Knife* [127]; and *A Jackass And A Fish* [69]. Extreme cases of IIFO are remarkable, publication-worthy and interesting to read, likely leading to publication bias.

Furthermore, extreme cases of recurrent intentional ingestion cause “countertransference anger” [128] among clinicians who have to endure sometimes years of repeat encounters with prolific recurrent ingesters, who are often complex patients with complex mental health needs. This is well described by Gitlin *et al.* [44] who describes how clinicians are “held hostage” by ingesters “in the midst of their self-harming process” after an initiating an “ongoing injurious sequence”. This anger perhaps finds an outlet in publishing literature.

2) *Selection Bias:* This review was conducted by a single reviewer (JGE), which introduces the potential for selection bias, motivation miss-classification and inconsistent application of eligibility criteria. Although inclusion decisions were guided by a

predefined protocol and supervision by an experienced reviewer (GC), dual independent screening was not performed due to resource limitations. This may have affected reproducibility and sensitivity of the search and screening processes.

3) *Unmeasured confounding:* The review did not record or analyse item location. This could have introduced an unmeasured confounding factor, perhaps reducing rates of perceived high risk ingestion, modulating outcome odds.

Time to presentation was not recorded. Ingesters that present early have increased successful endoscopy. This may have skewed endoscopy rates in some cases reports and case series, but the effect cannot be quantified in this review. It was attempted, but was difficult to standardise.

Also, this review did not record any alteration of objects that ingesters may have undertaken. Although, in clinical practice, this is likely difficult to assess. Sharp objects may wrapped to appear radiopaque and sharp on imaging, but in reality not be sharp or high risk, reducing the odds of morbidity [53, 54]. The opposite is also observed, as demonstrated in Bulgaria by the of home-made “gastrointestinal crosses”. Paperclips were wrapped with paper and elastic and designed to spring open in the distal gastrointestinal tract. This lead to five perforations in 5 patients in one week in one prison [99]. Intention and motivation in both examples playing a clear role.

4) *Information Bias:* Information bias from missing data in studies likely distorted true effects in the case series analysis. This is particularly in age group and object characteristics analysis, but also endoscopic outcomes and demographic characteristics.

G. Conclusion

This review represents the first systematic synthesis of outcomes following intentional ingestion of foreign objects (IIFO) with a focus on patient motivation. Across both case reports and case series, the nature of the ingestion—particularly protest versus intent-to-harm—appears to modulate clinical decision-making, especially in detained populations. While protest-motivated ingestion was often managed conservatively, intent-to-harm cases were associated with significantly increased rates of surgical intervention.

Middle-aged adults (40–64 years) were significantly less likely to undergo surgery with a non-significant tendency to increased surgical and conservative management. Subgroup analysis suggests that psychiatric history, prior ingestion, and multiple object ingestion may contribute to more conservative management strategies and refusal of treatment, modulating this effect. Sharp object ingestion, traditionally viewed as high-risk, was not uniformly associated with surgical outcomes, particularly in the context of protest motivations and detention settings. However, outside of protest motivations in this age group, those with psychiatric comorbidities had a tendency towards increased mortality.

Findings must be interpreted in light of limitations including small sample sizes, publication bias, selection bias, missing data, and unmeasured confounding factors such as object location, modification, and timing of presentation. Nonetheless, this review highlights the complexity of IIFO and the influence of both psychosocial context and clinical framing on treatment decisions.

Future research should focus on prospective, standardised reporting of IIFO cases to better understand how patient characteristics and motivations influence management. Greater clarity

around risk stratification may help clinicians provide safer, more consistent care for this often-complex population.

Population Differences in Key Subgroups




Fig. 5: Significant associated predictors from univariate analysis.

REFERENCES

- [1] Python Software Foundation. *Python*. May 2025.
- [2] Microsoft Corporation. *Visual Studio Code*. Microsoft Corporation. 2025. (Visited on 06/24/2025).
- [3] Kluyver, T., Ragan-Kelley, B., Peres, F., Granger, B., Bussonnier, M., Frederic, J., Kelley, K., Hamrick, J., Grout, J., Corlay, S., Ivanov, P., Avila, D., Abdalla, S., and Willing, C. *Jupyter Notebooks - a publishing format for reproducible computational workflows*. Positioning and Power in Academic Publishing: Players, Agents and Agendas. IOS Press, 2016.
- [4] Lamport, L. *LaTeX*. LaTeX Project Team. 1984.
- [5] The Pandas Development Team. *pandas-dev/pandas: Pandas*. Zenodo. Mar. 2020. doi: 10.5281/zenodo.13819579. (Visited on 04/03/2025).
- [6] Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, É. “Scikit-learn: Machine Learning in Python”. In: *Journal of Machine Learning Research* 12.85 (2011), pp. 2825–2830. ISSN: 1533-7928. (Visited on 04/04/2025).
- [7] Seabold, S. and Perktold, J. “Statsmodels: Econometric and statistical modeling with Python”. In: *Proceedings of the 9th Python in Science Conference*. Vol. 57. SciPy, 2010, pp. 92–96.
- [8] Waskom, M. L. “seaborn: statistical data visualization”. In: *Journal of Open Source Software* 6.60 (Apr. 2021), p. 3021. ISSN: 2475-9066. doi: 10.21105/joss.03021. (Visited on 06/24/2025).
- [9] Hunter, J. D. “Matplotlib: A 2D Graphics Environment”. In: *Computing in Science & Engineering* 9.3 (2007), pp. 90–95. ISSN: 1521-9615. doi: 10.1109/MCSE.2007.55. (Visited on 01/12/2025).
- [10] Met Office. “Cartopy: a cartographic Python library with a Matplotlib interface”. In: (2010).
- [11] Shen, L. *Forestplot*. 2022. (Visited on 07/04/2025).
- [12] UNHCR. *UNHCR: A record 100 million people forcibly displaced worldwide | UN News*. <https://news.un.org/en/story/2022/05/1118772>. May 2022. (Visited on 10/29/2024).
- [13] UNHCR. *Convention and Protocol Relating to the Status of Refugees*. <https://www.unhcr.org/media/convention-and-protocol-relating-status-refugees>. 2010. (Visited on 10/29/2024).
- [14] Amnesty International. *Refugees, Asylum Seekers and Migrants - Amnesty International*. <https://www.amnesty.org/en/what-we-do/refugees-asylum-seekers-and-migrants/>. 2024. (Visited on 10/29/2024).
- [15] Athwal, H. “‘I don’t have a life to live’: deaths and UK detention”. In: *Race & Class* 56.3 (Jan. 2015), pp. 50–68. ISSN: 0306-3968. doi: 10.1177/0306396814556224. (Visited on 10/29/2024).
- [16] Sundvall, M., Tidemalm, D. H., Titelman, D. E., Runeson, B., and Bäärnhielm, S. “Assessment and treatment of asylum seekers after a suicide attempt: a comparative study of people registered at mental health services in a Swedish location”. In: *BMC Psychiatry* 15.1 (Dec. 2015),

- p. 235. ISSN: 1471-244X. DOI: 10.1186/s12888-015-0613-8. (Visited on 10/29/2024).
- [17] Nickerson, A., Byrow, Y., O'Donnell, M., Mau, V., McMahon, T., Pajak, R., Li, S., Hamilton, A., Minihan, S., Liu, C., Bryant, R. A., Berle, D., and Liddell, B. J. "The association between visa insecurity and mental health, disability and social engagement in refugees living in Australia". In: *European Journal of Psychotraumatology* (Dec. 2019). ISSN: 2000-8198. (Visited on 10/29/2024).
- [18] Bevione, F., Panero, M., Abbate-Daga, G., Cossu, G., Carta, M. G., and Preti, A. "Risk of suicide and suicidal behavior in refugees. A meta-review of current systematic reviews and meta-analyses". In: *Journal of Psychiatric Research* 177 (Sept. 2024), pp. 287–298. ISSN: 0022-3956. DOI: 10.1016/j.jpsychires.2024.07.024. (Visited on 10/29/2024).
- [19] von Werthern, M., Robjant, K., Chui, Z., Schon, R., Ottisova, L., Mason, C., and Katona, C. "The impact of immigration detention on mental health: a systematic review". In: *BMC Psychiatry* 18.1 (Dec. 2018), p. 382. ISSN: 1471-244X. DOI: 10.1186/s12888-018-1945-y. (Visited on 10/29/2024).
- [20] Hedrick, K., Armstrong, G., Coffey, G., and Borschmann, R. "Self-harm in the Australian asylum seeker population: A national records-based study". In: *SSM - Population Health* 8 (Aug. 2019), p. 100452. ISSN: 23528273. DOI: 10.1016/j.ssmph.2019.100452. (Visited on 10/29/2024).
- [21] Global Detention Project. *United Kingdom Immigration Detention Profile*. <https://www.globaldetentionproject.org/countries/europe/united-kingdom>. 2024. (Visited on 11/02/2024).
- [22] Davidson, H. and Doherty, B. "Iranian refugee critically ill after setting himself on fire on Nauru during UN visit". In: *The Guardian* (Apr. 2016). ISSN: 0261-3077. (Visited on 06/26/2025).
- [23] Ajdacic-Gross, V., Weiss, M. G., Ring, M., Hepp, U., Bopp, M., Gutzwiler, F., and Rössler, W. "Methods of suicide: international suicide patterns derived from the WHO mortality database". In: *Bulletin of the World Health Organization* 86.9 (June 2008), p. 726. DOI: 10.2471/BLT.07.043489. (Visited on 10/29/2024).
- [24] Becq, A., Camus, M., and Dray, X. "Foreign body ingestion: dos and don'ts". In: *Frontline Gastroenterology* 12.7 (Dec. 2021), pp. 664–670. ISSN: 2041-4137, 2041-4145. DOI: 10.1136/flgastro-2020-101450. (Visited on 05/15/2025).
- [25] Ikenberry, S. O., Jue, T. L., Anderson, M. A., Appalaneni, V., Banerjee, S., Ben-Menachem, T., Decker, G. A., Fanelli, R. D., Fisher, L. R., Fukami, N., Harrison, M. E., Jain, R., Khan, K. M., Krinsky, M. L., Maple, J. T., Sharaf, R., Strohmeyer, L., and Dominitz, J. A. "Management of ingested foreign bodies and food impactions". In: *Gastrointestinal Endoscopy* 73.6 (June 2011), pp. 1085–1091. ISSN: 00165107. DOI: 10.1016/j.gie.2010.11.010. (Visited on 11/18/2024).
- [26] Birk, M., Bauerfeind, P., Deprez, P., Häfner, M., Hartmann, D., Hassan, C., Hucl, T., Lesur, G., Aabakken, L., and Meining, A. "Removal of foreign bodies in the upper gastrointestinal tract in adults: European Society of Gastrointestinal Endoscopy (ESGE) Clinical Guideline". In: *Endoscopy* 48.05 (Feb. 2016), pp. 489–496. ISSN: 0013-726X, 1438-8812. DOI: 10.1055/s-0042-100456. (Visited on 09/28/2024).
- [27] Vujkovic, M. "The suffering of Omid: How a burns victim died despite injuries that were 'very survivable'". In: *ABC News* (Mar. 2019). (Visited on 10/29/2024).
- [28] Hsieh, A., Hsiehchen, D., Layne, S., Ginsberg, G. G., and Keo, T. "Trends and clinical features of intentional and accidental adult foreign body ingestions in the United States, 2000 to 2017". In: *Gastrointestinal Endoscopy* 91.2 (Feb. 2020), 350–357.e1. ISSN: 0016-5107, 1097-6779. DOI: 10.1016/j.gie.2019.09.010. (Visited on 11/02/2024).
- [29] Palta, R., Sahota, A., Bemarki, A., Salama, P., Simpson, N., and Laine, L. "Foreign-body ingestion: characteristics and outcomes in a lower socioeconomic population with predominantly intentional ingestion". In: *Gastrointestinal endoscopy* 69.3 Pt 1 (Mar. 2009), pp. 426–433. ISSN: 1097-6779 0016-5107. DOI: 10.1016/j.gie.2008.05.072.
- [30] Tromans, S., Chester, V., Wells, H., and Alexander, R. "Deliberate ingestion of foreign bodies as a form of self-harm among inpatients within forensic mental health and intellectual disability services". In: *Journal of Forensic Psychiatry & Psychology* 30.2 (Apr. 2019), pp. 189–202. ISSN: 1478-9949. DOI: 10.1080/14789949.2018.1530287.
- [31] Hunt, I., Hartley, S., Alwahab, Y., and Birkill, G. J. "Aortoesophageal perforation following ingestion of razorblades with massive haemothorax". In: *European journal of cardio-thoracic surgery : official journal of the European Association for Cardio-thoracic Surgery* 31.5 (May 2007), pp. 946–948. ISSN: 1010-7940. DOI: 10.1016/j.ejcts.2007.01.073.
- [32] Udemgba, C., Missov, E., Percy, R., and Sattiraju, S. "A case report of an unusual left atrial mass". In: *European heart journal. Case reports* 5.1 (Jan. 2021), ytaa500. ISSN: 2514-2119. DOI: 10.1093/euher/ytaa500.
- [33] Zhao, G. "Journal Of Forensic Sciences". In: (2022). DOI: 10.1111/1556-4029.15079.
- [34] Bozorgmehr, R., Bahadorinia, M., Pouyanfar, S., Ahmadinejad, M., Bahri, M. H., and Bagherpour, J. Z. "A rare case of abdominal foreign bodies; laparoscopic removal of a sewing needle". In: *Annals of medicine and surgery* (2012) 82 (Oct. 2022), p. 104747. ISSN: 2049-0801. DOI: 10.1016/j.amsu.2022.104747.
- [35] Moehlau, F. G. "Gastrostomy in the Seventeenth Century". In: *Buffalo Medical Journal* 35.5 (Dec. 1895), pp. 395–397. ISSN: 1040-3817.
- [36] Saint, J. H. "Surgery Of The Esophagus". In: *Archives of Surgery* 19.1 (July 1929), pp. 53–128. ISSN: 0272-5533. DOI: 10.1001/archsurg.1929.01150010056003. (Visited on 04/15/2025).
- [37] Barros, J. L., Caballero, A., Rueda, J. C., and Monturiol, J. M. "Foreign body ingestion: management of 167 cases". In: *World Journal of Surgery* 15.6 (1991), pp. 783–788. ISSN: 0364-2313. DOI: 10.1007/BF01665320.
- [38] Lerche, W. "The Esophagoscope In Removing Sharp Foreign Bodies From The Esophagus". In: *Journal of the American Medical Association* LVI.9 (Mar. 1911), p. 634. ISSN: 0002-9955. DOI: 10.1001/jama.1911.02560090004002. (Visited on 04/15/2025).

- [39] Jackson, C. L. "Foreign bodies in the esophagus". In: *The American Journal of Surgery* 93.2 (Feb. 1957), pp. 308–312. ISSN: 00029610. DOI: 10.1016/0002-9610(57)90783-3. (Visited on 04/15/2025).
- [40] Ricote, G. C., Torre, L. R., De Ayala, V. P., Castellanos, D., Menchen, P., Senent, C., Velo, J. L., Robles, J., and Alcalá-Santaella, R. "Fiberendoscopic removal of foreign bodies of the upper part of the gastrointestinal tract". In: *Surgery, Gynecology & Obstetrics* 160.6 (June 1985), pp. 499–504. ISSN: 0039-6087.
- [41] Chalk, S. G. "Foreign Bodies In The Stomach: Report Of A Case In Which More Than Two Thousand Five Hundred Foreign Bodies Were Found". In: *Archives of Surgery* 16.2 (Feb. 1928), p. 494. ISSN: 0272-5533. DOI: 10.1001/archsurg.1928.01140020045003. (Visited on 04/15/2025).
- [42] Puggioni, R. "Speaking through the body: detention and bodily resistance in Italy". In: *Citizenship Studies* 18.5 (July 2014), pp. 562–577. ISSN: 1362-1025. DOI: 10.1080/13621025.2014.923707. (Visited on 10/29/2024).
- [43] Losanoff, J. E., Kjossev, K. T., and Losanoff, H. E. "Oesophageal "cross"—a sinister foreign body." In: *Journal of accident & emergency medicine* 14.1 (Jan. 1997), pp. 54–55. ISSN: 1351-0622. DOI: 10.1136/emj.14.1.54.
- [44] Gitlin, D. F., Caplan, J. P., Rogers, M. P., Avni-Barron, O., Braun, I., and Barsky, A. J. "Foreign-body ingestion in patients with personality disorders". In: *Psychosomatics* 48.2 (2007), pp. 162–166. ISSN: 0033-3182. DOI: 10.1176/appi.psy.48.2.162.
- [45] Poynter, B. A., Hunter, J. J., Coverdale, J. H., and Kempinsky, C. A. "Hard to swallow: a systematic review of deliberate foreign body ingestion". In: *General Hospital Psychiatry* 33.5 (Sept. 2011), pp. 518–524. ISSN: 0163-8343. DOI: 10.1016/j.genhosppsych.2011.06.011. (Visited on 09/24/2024).
- [46] Al-Faham, F. S. M. and Al-Hakkak, S. M. M. "The largest esophageal foreign body in adults: A case report". In: *Annals of Medicine and Surgery* (2012) 54 (June 2020), pp. 82–84. ISSN: 2049-0801. DOI: 10.1016/j.amsu.2020.04.039.
- [47] Pantazopoulos, I., Mavrovounis, G., Mermiri, M., Adamou, A., and Gourgoulianis, K. "Intentional ingestion of batteries and razor blades by a prisoner: A true emergency?" In: *International Journal of Prisoner Health* 18.3 (2022), pp. 316–322. ISSN: 1744-9200. DOI: 10.1108/IJPH-06-2021-0054.
- [48] Aitchison, G. and Essex, R. "Self-harm in immigration detention: political, not (just) medical". In: *Journal of Medical Ethics* 50.11 (Nov. 2024), pp. 786–793. ISSN: 0306-6800, 1473-4257. DOI: 10.1136/jme-2022-108366. (Visited on 10/29/2024).
- [49] Jaini, P. A., Haliburton, J., and Rush, A. J. "Management Challenges of Recurrent Foreign Body Ingestions in a Psychiatric Patient: A Case Report." In: *Journal of psychiatric practice* 29.2 (Mar. 2023), pp. 167–173. ISSN: 1538-1145 1527-4160. DOI: 10.1097/PRA.0000000000000694.
- [50] Bhugra, D., Craig, T. K. J., and Bhui, K. *Mental Health of Refugees and Asylum Seekers*. OUP Oxford, Aug. 2010. ISBN: 978-0-19-955722-6.
- [51] Haase, E., Schönfelder, A., Nesterko, Y., and Glaesmer, H. "Prevalence of suicidal ideation and suicide attempts among refugees: a meta-analysis". In: *BMC Public Health* 22.1 (Apr. 2022), p. 635. ISSN: 1471-2458. DOI: 10.1186/s12889-022-13029-8. (Visited on 10/29/2024).
- [52] Hedrick, K. and Borschmann, R. "Self-harm among unaccompanied asylum seekers and refugee minors: protocol for a global systematic review of prevalence, methods and characteristics". In: *BMJ Open* 13.6 (June 2023), e069237. ISSN: 2044-6055, 2044-6055. DOI: 10.1136/bmjopen-2022-069237. (Visited on 10/03/2024).
- [53] Ataya, A., Alraiyes, A., and Alraies, M. "Razor blades in the stomach". In: *QJM: An International Journal of Medicine* 106.8 (Aug. 2013), pp. 783–784. ISSN: 1460-2725. DOI: 10.1093/qjmed/hcs165. (Visited on 04/14/2025).
- [54] Albeldawi, M. and Birgisson, S. "Conservative management of razor blade ingestion". In: *Gastroenterology Report* 2.2 (May 2014), pp. 158–159. ISSN: 2052-0034. DOI: 10.1093/gastro/gou002. (Visited on 04/14/2025).
- [55] Machin, D., Campbell, M. J., and Walters, S. J. K. *Medical statistics: a textbook for the health sciences*. 4. ed., reprinted. Chichester: Wiley, 2008. ISBN: 978-0-470-02519-2.
- [56] Microsoft Corporation. *Microsoft Excel*. 2025.
- [57] Moola, S., Munn, Z., Tufanaru, C., Aromataris, E., Sears, K., Sfetcu, R., Sfetcu, M., Lisy, K., Qureshi, R., Mattis, P., and Mu, P. "Chapter 7: Systematic reviews of etiology and risk". In: *JBI Manual for Evidence Synthesis*. JBI, 2020. ISBN: 978-0-6488488-0-6. DOI: 10.46658/JBIMES-20-08. (Visited on 05/07/2025).
- [58] Tanriver-Ayder, E., Faes, C., van de Castele, T., McCann, S. K., and Macleod, M. R. "Comparison of commonly used methods in random effects meta-analysis: application to preclinical data in drug discovery research". In: *BMJ open science* 5.1 (2021), e100074. ISSN: 2398-8703. DOI: 10.1136/bmjos-2020-100074.
- [59] Sterne, J. A. C., Egger, M., Moher, D., and Boutron. "Cochrane Handbook for Systematic Reviews of Interventions". In: *Chapter 10*. Vol. 5.2.0. Cochrane, 2017. (Visited on 06/30/2025).
- [60] Akay, S., Günay, S., Binicier, Ö. B., Paköz, Z. B., and Akar, H. "A deliberately swallowed foreign body: money package." In: *Endoscopy* 47 Suppl 1 (2015), E602–603. ISSN: 1438-8812 0013-726X. DOI: 10.1055/s-0035-1569668.
- [61] Al Shaaibi, R. and Al Waili, I. "Laparoscopic Retrieval of Ingested Foreign Body." In: *Oman medical journal* 36.3 (May 2021), e264. ISSN: 1999-768X 2070-5204. DOI: 10.5001/omj.2021.35.
- [62] Alao, A. O. and Abraham, B. "Foreign body ingestions in a schizophrenic patient." In: *West African journal of medicine* 25.3 (Sept. 2006), pp. 239–241. ISSN: 0189-160X. DOI: 10.4314/wajm.v25i3.28286.
- [63] Ali, S. M. "Duodenal Perforation by Swallowed Toothbrush: Case Report and Review of Literature". In: *Open Access Journal of Surgery* 4.2 (May 2017). ISSN: 24761346. DOI: 10.19080/OAJS.2017.04.555632. (Visited on 04/14/2025).

- [64] Ali, A. and Alhindi, S. "A Child With a Gastrocolic Fistula After Ingesting Magnets: An Unusual Complication." In: *Cureus* 12.7 (July 2020), e9336. ISSN: 2168-8184. DOI: 10.7759/cureus.9336.
- [65] Ali, A., Mahgoub, A. M., Emad, S., and Abdelfattah, A. H. "Endoscopic Retrieval of an Ingested Mobile Phone From the Stomach of a Prisoner: When Gastroenterologists Answer the Call." In: *Cureus* 14.12 (Dec. 2022), e33053. ISSN: 2168-8184. DOI: 10.7759/cureus.33053.
- [66] Apikotoa, S., Ballal, H., and Wijesuriya, R. "Endoscopic foreign body retrieval from the caecum - A case report and push for intervention guidelines." In: *International journal of surgery case reports* 90 (Jan. 2022), p. 106755. ISSN: 2210-2612. DOI: 10.1016/j.ijscr.2022.106755.
- [67] Atayan, Y., Cagin, Y. F., Erdogan, M. A., Bilgic, Y., Bestas, R., Harputluoglu, M., and Seckin, Y. "Lighter Ingestion as an Uncommon Cause of Severe Vomiting in a Schizophrenia Patient". In: *Case Reports in Gastrointestinal Medicine* 2016 (2016), p. 6301302. ISSN: 2090-6528. DOI: 10.1155/2016/6301302.
- [68] Beecroft, N., Bach, L., Tunstall, N., and Howard, R. "An unusual case of pica". In: *International Journal of Geriatric Psychiatry* 13.9 (Sept. 1998), pp. 638–641. ISSN: 0885-6230. DOI: 10.1002/(sici)1099-1166(199809)13:9<638::aid-gps837>3.0.co;2-n.
- [69] Benoist, L. "Acta Oto-Laryngologica Case Reports". In: (2019). DOI: 10.1080/23772484.2018.1555436.
- [70] Berry, P. and Kotha, S. "Crying wolf: the danger of recurrent intentional foreign body ingestion." In: *Frontline gastroenterology* 13.3 (2022), p. 266. ISSN: 2041-4137 2041-4145. DOI: 10.1136/flgastro-2021-101888.
- [71] Bhasin, S. K., Kachroo, S. L., Kumar, V., Kumar, R., and Chandail, V. S. "7" long knife for 7 years in the duodenum: a rare case report and review of literature". In: *International Surgery Journal* 1.1 (2014), pp. 29–32. ISSN: 2349-2902. (Visited on 04/14/2025).
- [72] Bhattacharjee, P. and Singh, O. "Repeated ingestion of sharp-pointed metallic objects". In: *Archives of Iranian medicine* 11 (Oct. 2008), pp. 563–5.
- [73] Bhumi, S., Mago, S., Mavilia-Scranton, M. G., Birk, J. W., and Rezaizadeh, H. "Esophageal Button Battery Retrieval: Time-In May Not Be Everything." In: *Cureus* 16.4 (Apr. 2024), e58327. ISSN: 2168-8184. DOI: 10.7759/cureus.58327.
- [74] Camacho Dorado, C., Sánchez Gallego, A., Miota de Llama, J. I., and González Masiá, J. A. "Metallic bezoar after suicide attempt". In: *Cirugia Espanola* 96.8 (Oct. 2018), p. 515. ISSN: 2173-5077. DOI: 10.1016/j.ciresp.2018.02.015.
- [75] Cauchi, J. A. and Shawis, R. N. "Multiple magnet ingestion and gastrointestinal morbidity". In: *Archives of Disease in Childhood* 87.6 (Dec. 2002), pp. 539–540. ISSN: 1468-2044. DOI: 10.1136/adc.87.6.539.
- [76] Chang, W.-J. and Chiu, W.-Y. "Gastric foreign body: a comb." In: *Clinical case reports* 5.6 (June 2017), pp. 1036–1037. ISSN: 2050-0904. DOI: 10.1002/ccr3.957.
- [77] Cox, D., Donohue, P., and Costa, V. "A swallowed toothbrush causing perforation 2 years after ingestion". In: *British Journal of Hospital Medicine (London, England)* 2005) 68.10 (Oct. 2007), p. 559. ISSN: 1750-8460. DOI: 10.12968/hmed.2007.68.10.27330.
- [78] Csáky, G., Szederkényi, I., Botos, A., and Kiss, I. "Laparoscopic removal of a foreign body from the jejunum." In: *Surgical laparoscopy & endoscopy* 8.1 (Feb. 1998), pp. 68–70. ISSN: 1051-7200.
- [79] Salazar, J., Pacheco, N., Jaramillo, P., Yépez, S., Loza, V., Alvarado, C., Ayala, B., and Molina, G. "Ingestion of razor blades, a rare event: A case report in a psychiatric patient". In: *Journal of Surgical Case Reports* 2020.5 (2020). ISSN: 2042-8812. DOI: 10.1093/JSCR/RJAA094.
- [80] Divsalar, P. "International Journal Of High Risk Behaviors And Addiction". In: (2023). DOI: 10.5812/ijhrba-134720.
- [81] Emamhadi, M. A., Najari, F., Hedayatshode, M. J., and Sharif, S. "Sudden Death Following Oral Intake of Metal Objects (Acuphagia): a Case Report". In: *Emergency (Tehran, Iran)* 6.1 (2018), e16. ISSN: 2345-4563.
- [82] Farhadi, F., Mohtadi, A., Pakmehr, M., Ghaedamini, H., Shafieian, F., and Aminifar, S. A. "This is a successful removal of more than 450 pieces of metal objects from a patient's stomach: a case report". In: *Journal of Medical Case Reports* 18.1 (Aug. 2024), p. 381. ISSN: 1752-1947. DOI: 10.1186/s13256-024-04672-3. (Visited on 12/10/2024).
- [83] Fry, E. and Counselman, F. L. "A right scrotal abscess and foreign body ingestion in a schizophrenic patient". In: *The Journal of Emergency Medicine* 38.5 (June 2010), pp. 587–592. ISSN: 0736-4679. DOI: 10.1016/j.jemermed.2007.07.018.
- [84] Gardner, A., Radwan, R., Allison, M., and Codd, R. "Double duodenal perforation following foreign body ingestion". In: *BMJ Case Reports* 2017 (2017). ISSN: 1757790X (ISSN). DOI: 10.1136/bcr-2017-223182.
- [85] Goldman, R. D., Schachter, P., Katz, M., Bilik, R., and Avigad, I. "A bizarre bezoar: case report and review of the literature." In: *Pediatric surgery international* 14.3 (Dec. 1998), pp. 218–219. ISSN: 0179-0358. DOI: 10.1007/s003830050492.
- [86] Guinan, D., Drvar, T., Brubaker, D., Ang-Rabanes, M., Kupec, J., and Marshalek, P. "Intentional Foreign Body Ingestion: A Complex Case of Pica." In: *Case reports in gastrointestinal medicine* 2019 (2019), p. 7026815. ISSN: 2090-6528 2090-6536. DOI: 10.1155/2019/7026815.
- [87] Hardy, J. C., Ashcroft, C., Kay, C., Liane, B.-J., and Horn, C. "Loose Screws: Removal of Foreign Bodies From the Lower Gastrointestinal Tract." In: *Cureus* 15.8 (Aug. 2023), e43093. ISSN: 2168-8184. DOI: 10.7759/cureus.43093.
- [88] Jehangir, M., Mallory, C., and Medverd, J. R. "Digital tomosynthesis for detection of ingested foreign objects in the emergency department: a case of razor blade ingestion." In: *Emergency radiology* 26.2 (Apr. 2019), pp. 249–252. ISSN: 1438-1435 1070-3004. DOI: 10.1007/s10140-018-01664-x.
- [89] Jin, S., Horiguchi, T., Ma, X., Yuan, S., and Liu, Q. "Metallic foreign bodies ingestion by schizophrenic patient: a case report". In: *Annals of Medicine and Surgery* 85.4 (Apr. 2023), p. 1270. ISSN: 2049-0801. DOI: 10.1097/MS9.000000000000497. (Visited on 03/30/2025).

- [90] Kar, S. K., Kamboj, A., and Kumar, R. "Pica and psychosis - clinical attributes and correlations: a case report". In: *Journal of Family Medicine and Primary Care* 4.1 (2015), pp. 149–150. ISSN: 2249-4863. DOI: 10.4103/2249-4863.152277.
- [91] Kariholu, P. L., Jakareddy, R., Hemanthkumar, M., Paramesh, K. N., and Pavankumar, N. P. "Pica - a case of acuphagia or hyalophagia?" In: *The Indian Journal of Surgery* 70.3 (June 2008), pp. 144–146. ISSN: 0972-2068. DOI: 10.1007/s12262-008-0040-x.
- [92] Kerestes, T. and Smith, J. "Paper or Plastic? A Foreign Body Ingestion Leading to Small Bowel Obstruction. A Case Report". In: *ARC Journal of Clinical Case Reports* 5.2 (2019). ISSN: 24559806. DOI: 10.20431/2455-9806.0502002. (Visited on 04/14/2025).
- [93] Kobiela, J., Mittlener, S., Gorycki, T., Lachinski, A., and Adrych, K. "Vast collection of foreign bodies in the stomach presenting as acute gastrointestinal bleeding in a patient with schizophrenia". In: *Endoscopy* 47.S 01 (July 2015), E356–E357. ISSN: 0013-726X, 1438-8812. DOI: 10.1055/s-0034-1392611. (Visited on 04/14/2025).
- [94] Kumar, A. and Jazieh, A. R. "Case report of sideroblastic anemia caused by ingestion of coins". In: *American Journal of Hematology* 66.2 (Feb. 2001), pp. 126–129. ISSN: 0361-8609. DOI: 10.1002/1096-8652(200102)66:2<126::AID-AJH1029>3.0.CO;2-J.
- [95] Kumar, R., Soni, N., Bhardwaj, S., Bhoil, R., and Gupta, S. C. "Intentional foreign body ingestion." In: *Internal and emergency medicine* 14.8 (Nov. 2019), pp. 1331–1333. ISSN: 1970-9366 1828-0447. DOI: 10.1007/s11739-019-02183-4.
- [96] Lee, J. H., Kim, H. C., Yang, D. M., Kim, S. W., Jin, W., Park, S. J., and Kim, H. J. "What is the role of plain radiography in patients with foreign bodies in the gastrointestinal tract?" In: *Clinical imaging* 36.5 (Oct. 2012), pp. 447–454. ISSN: 1873-4499 0899-7071. DOI: 10.1016/j.clinimag.2011.11.017.
- [97] Li, Q.-P., Ge, X.-X., Ji, G.-Z., Fan, Z.-N., Zhang, F.-M., Wang, Y., and Miao, L. "Endoscopic retrieval of 28 foreign bodies in a 100-year-old female after attempted suicide". In: *World Journal of Gastroenterology* 19.25 (July 2013), pp. 4091–4093. ISSN: 2219-2840. DOI: 10.3748/wjg.v19.i25.4091.
- [98] Liu, S., de Blacam, C., Lim, F.-Y., Mattei, P., and Mamula, P. "Magnetic foreign body ingestions leading to duodenocolonic fistula". In: *Journal of Pediatric Gastroenterology and Nutrition* 41.5 (Nov. 2005), pp. 670–672. ISSN: 0277-2116. DOI: 10.1097/01.mpg.0000177703.99786.c9.
- [99] Losanoff, J. E. and Kjossev, K. T. "Gastrointestinal 'crosses'. A new shade from an old palette." In: *Archives of surgery (Chicago, Ill. : 1960)* 131.2 (Feb. 1996), pp. 166–169. ISSN: 0004-0010. DOI: 10.1001/archsurg.1996.01430140056015.
- [100] Mesfin, T., Tekalegn, Y., Aman, M., Geta, G., Ketema, A., Defere, F., Girma, D., Tsegaye, M., Mengistu, T., and Seyoum, K. "Ingestion of Metallic Materials Found in the Stomach and First Part of the Duodenum of a Schizophrenic Man: Case Report". In: *International Medical Case Reports Journal* 15 (2022), pp. 681–684. ISSN: 1179-142X. DOI: 10.2147/IMCRJ.S386883.
- [101] Misra, S., Jain, V., Ahmad, F., Kumar, R., and Kishore, N. "Metallic sewing needle ingestion presenting as acute abdomen". In: *Nigerian Journal of Clinical Practice* 16.4 (2013), pp. 540–543. ISSN: 1119-3077. DOI: 10.4103/1119-3077.116879.
- [102] Naji, H., Isacson, D., Svensson, J. F., and Wester, T. "Bowel injuries caused by ingestion of multiple magnets in children: a growing hazard." In: *Pediatric surgery international* 28.4 (Apr. 2012), pp. 367–374. ISSN: 1437-9813 0179-0358. DOI: 10.1007/s00383-011-3026-x.
- [103] Ohno, Y., Yoneda, A., Enjoji, A., Furui, J., and Kanematsu, T. "Gastroduodenal fistula caused by ingested magnets". In: *Gastrointestinal Endoscopy* 61.1 (Jan. 2005), pp. 109–110. ISSN: 0016-5107. DOI: 10.1016/s0016-5107(04)02387-9.
- [104] Peixoto, A., Pereira, P., and Macedo, G. "Gastrointestinal: Voluntary padlock ingestion." In: *Journal of gastroenterology and hepatology* 32.12 (Dec. 2017), p. 1910. ISSN: 1440-1746 0815-9319. DOI: 10.1111/jgh.13828.
- [105] Qureshi, N. A., Cherian, N., Ben-Hamida, A., and Solkar, M. H. "Endoscopic Retrieval of an Intentionally Ingested Mobile Phone in an Adult: First Case Report of its Kind". In: *Annals of Clinical Case Reports* 1 (2016).
- [106] Riva, C. G., Toti, F. A. T., Siboni, S., and Bonavina, L. "Unusual foreign body impacted in the upper oesophagus: original technique for transoral extraction." In: *BMJ case reports* 2018 (June 2018), bcr-2018-225241. ISSN: 1757-790X. DOI: 10.1136/bcr-2018-225241.
- [107] Sakellaridis, T. "Annals Of Thoracic Surgery". In: (2008). DOI: 10.1016/j.athoracsur.2007.07.027.
- [108] Sharma, J. "Journal Of The Academy Of Consultation-Liaison Psychiatry". In: (2022). DOI: 10.1016/j.jaclp.2022.10.025.
- [109] Sobnach, S., Castillo, F., Blanco Vinent, R., Kahn, D., and Bhyat, A. "Penetrating cardiac injury following sewing needle ingestion." In: *Heart, lung & circulation* 20.7 (July 2011), pp. 479–481. ISSN: 1444-2892 1443-9506. DOI: 10.1016/j.hlc.2011.01.006.
- [110] Sultan, N., Attar, H., Sembawa, H., and Alharthi, H. "A plastic bezoar causing bowel obstruction: A case of table cover ingestion." In: *International journal of surgery case reports* 117 (Apr. 2024), p. 109506. ISSN: 2210-2612. DOI: 10.1016/j.ijscr.2024.109506.
- [111] Tammana, V. S., Valluru, N., and Sanderson, A. "All the wrong places: an unusual case of foreign body ingestion and inhalation." In: *Case reports in gastroenterology* 6.3 (Sept. 2012), pp. 778–783. ISSN: 1662-0631. DOI: 10.1159/000346287.
- [112] Tanrikulu, Y. "Hong Kong Journal Of Emergency Medicine". In: (2015). DOI: 10.1177/102490791502200107.
- [113] Tay, E. T., Weinberg, G., and Levin, T. L. "Ingested magnets: the force within". In: *Pediatric Emergency Care* 20.7 (July 2004), pp. 466–467. ISSN: 1535-1815. DOI: 10.1097/01.pec.0000134926.03030.a7.
- [114] Thapa, N., Basnyat, S., and Maharjan, M. "Ingestion of Bell Clappers by a Shaman in Jumla, Nepal: A Case Report." In: *JNMA; journal of the Nepal Medical Association* 57.215 (Feb. 2019), pp. 56–58. ISSN: 1815-672X 0028-2715. DOI: 10.31729/jnma.4055.

- [115] Trgo, G., Tonkic, A., Simunic, M., and Puljiz, Z. "Successful endoscopic removal of a lighter swallowed 17 months before." In: *Case reports in gastroenterology* 6.2 (May 2012), pp. 238–242. ISSN: 1662-0631. DOI: 10.1159/000338839.
- [116] Tupesis, J. P., Kaminski, A., Patel, H., and Howes, D. "A penny for your thoughts: small bowel obstruction secondary to coin ingestion." In: *The Journal of emergency medicine* 27.3 (Oct. 2004), pp. 249–252. ISSN: 0736-4679. DOI: 10.1016/j.jemermed.2004.03.013.
- [117] Wadhwa, C. "Journal Of Digestive Endoscopy". In: (2015). DOI: 10.4103/0976-5042.159247.
- [118] Wildhaber, B. E., Le Coultr, C., and Genin, B. "Ingestion of magnets: innocent in solitude, harmful in groups". In: *Journal of Pediatric Surgery* 40.10 (Oct. 2005), e33–35. ISSN: 1531-5037. DOI: 10.1016/j.jpedsurg.2005.06.022.
- [119] Wnęk, B., Łożyńska-Nelke, A., and Karoń, J. "Foreign body in the gastrointestinal tract leading to small bowel obstruction—case report and literature review." In: *Polski przeglad chirurgiczny* 86.12 (Mar. 2015), pp. 594–597. ISSN: 2299-2847 0032-373X. DOI: 10.1515/pjs-2015-0006.
- [120] Yasin, M. A., Malik, G. N., Malik, S. A., and Rathore, F. A. "Metal in stomach: a rare cause of gastric bezoar". In: *BMJ Case Reports* 2009 (Feb. 2009), bcr06.2008.0278. ISSN: 1757-790X. DOI: 10.1136/bcr.06.2008.0278. (Visited on 04/14/2025).
- [121] Yıldız, İ., Koca, Y. S., Avşar, G., and Barut, İ. "Tendency to Ingest Foreign Bodies in Mentally Retarded Patients: A Case with Ileal Perforation Caused by the Ingestion of a Teaspoon." In: *Case reports in surgery* 2016 (2016), p. 8075432. ISSN: 2090-6900 2090-6919. DOI: 10.1155/2016/8075432.
- [122] Fj Buils. "Repeated Behavior Of Deliberate Foreign Body Ingestion In A Patient With Psychiatric Disorder". In: *A Case Report. Clin Surg* (2024). DOI: 10.52916/jmrs244144.
- [123] te Wildt, B. T., Tettenborn, C., Schneider, U., Ohlmeier, M. D., Zedler, M., Zakhalev, R., and Krueger, M. "Swallowing Foreign Bodies as an Example of Impulse Control Disorder in a Patient With Intellectual Disabilities". In: *Psychiatry (Edgmont)* 7.9 (Sept. 2010), pp. 34–37. ISSN: 1550-5952. (Visited on 04/14/2025).
- [124] Elghali, M. amine, Ghrissi, R., Fadhl, H., Mahjoub, M., Jarrar, M. S., Jedidi, M., Letaief, R., and Hamila, F. "The Management of Voluntary Ingestion of Razor Blades by Inmates". In: *International Surgery* 105.1-3 (Nov. 2016), pp. 129–133. ISSN: 0020-8868. DOI: 10.9738/INTSURG-D-16-00204.1. (Visited on 04/14/2025).
- [125] Karp, J. G., Whitman, L., and Convit, A. "Intentional Ingestion of Foreign Objects by Male Prison Inmates". In: *Psychiatric Services* 42.5 (May 1991), pp. 533–535. ISSN: 1075-2730, 1557-9700. DOI: 10.1176/ps.42.5.533. (Visited on 09/27/2025).
- [126] Lee, T. H., Kang, Y. W., Kim, H. J., Kim, S. M., Im, E. H., Huh, K. C., Choi, Y. W., Kim, T. H., Lee, O. J., and Jung, U. T. "Foreign Objects in Korean Prisoners". In: *The Korean Journal of Internal Medicine* 22.4 (Dec. 2007), pp. 275–278. ISSN: 1226-3303, 2005-6648. DOI: 10.3904/kjim.2007.22.4.275. (Visited on 04/14/2025).
- [127] Fine, S., Watson, J. B., and Habr, F. "Now You See It, Endo You Don't: Case of the Disappearing Knife". In: *Gastroenterology* 144.7 (June 2013), e6–e7. ISSN: 0016-5085, 1528-0012. DOI: 10.1053/j.gastro.2013.01.059. (Visited on 04/14/2025).
- [128] Palese, C. and Al-Kawas, F. H. "Repeat intentional foreign body ingestion: the importance of a multidisciplinary approach". In: *Gastroenterology & Hepatology* 8.7 (July 2012), pp. 485–486. ISSN: 1554-7914.
- [129] American Psychiatric Association. *Diagnostic and statistical manual of mental disorders (5th ed.)* 2013.
- [130] UN General Assembly. *Protocol Relating to the Status of Refugees*. Jan. 1967. (Visited on 05/27/2025).
- [131] Deng, F. M., Persons, U. R. o. t. S.-G. o. I. D., and Affairs, U. O. f. t. C. o. H. *Guiding Principles on Internal Displacement*. Tech. rep. UN Doc E/CN.4/1998/53/Add.2. United Nations Commission on Human Rights, Feb. 1998. (Visited on 03/31/2025).
- [132] United Nations High Commissioner for Refugees. *Asylum-seekers*. <https://www.unhcr.org/us/about-unhcr/who-we-protect/asylum-seekers>. 2025. (Visited on 05/27/2025).

APPENDIX A
METHODS

A. Eligibility Criteria

TABLE VII: Inclusion criteria structured using the PICOS framework.

Category	Details
Population	Any human. Any age group.
Interventions or exposures	Humans that have: – Non-accidental ingestion – Ingestion of a true foreign body (non-nutritive items)
Comparators / Control group	Demographic: – Gender – Age – Detained Person – Psychiatric Inpatient – Displaced Person – Under Influence of Alcohol – Psychiatric History – Severely Disabled – Previous Ingestion Motivation: – Intent to harm – Psychiatric – Psychosocial – Protest – Other Object characteristics: – Button battery – Magnet – Long (>5 cm) – Large diameter (>2.5 cm) – Multiple – Blunt objects – Sharp-pointed objects
Outcomes of interest	– Endoscopic intervention – Surgical intervention – Conservative management – Complication rates – Mortality rates
Setting	Any setting.
Study designs	Any design.

TABLE VIII: Exclusion criteria for study exclusion.

#	Exclusion Criterion
1	Full text not available in English.
2	Studies not focusing on intentional self-ingestion (into the gastrointestinal tract) of a foreign object via the oral cavity (mouth), or where it is unclear if ingestion occurred.
3	Studies focusing solely on accidental ingestion.
4	Non-human or animal studies.
5	Reviews, editorials, commentaries, and opinion pieces without original empirical data.
6	Duplicate publications or studies with overlapping datasets (only the most comprehensive or recent study was included).
7	Studies focusing on ingestion or co-ingestion of substances (e.g., poisons, medications) rather than physical foreign objects.
8	Ingestions undertaken in controlled environments as part of a voluntary study.
9	Ingestions not explicitly stated to be intentional and not suggestive of deliberate ingestion.
10	Does not meet inclusion criteria.
11	Ingestions where death resulted from other means (e.g., suicide by other method).
12	Studies published before the advent of endoscopy (1906).
13	Outcomes not reported.
14	Motivation not reported.
15	Object characteristics not reported

B. Search Strategy

TABLE IX: Concepts with associated keywords and MeSH terms used in PubMed search strategy.

Concept	Keywords	MeSH Terms
Foreign Bodies	"foreign obj*" "foreign bod*"	Foreign Bodies [MeSH]
Intentional Ingestion / Self-harm	"intent*" "deliberate*" "purpose*" "self-injur*" "selfharm*" "self-harm*"	Self-Injurious Behavior [MeSH]
Ingestion Behavior	"ingest*" "swallow*"	-
Interventions	"surg*" "endoscop*" "EGD" "OGD" "Esophagogastroduodenoscopy" "Oesophagogastroduodenoscopy" "manag*"	Endoscopy [MeSH] Surgical Procedures, Operative [MeSH] Conservative Treatment [MeSH] Drug Therapy [MeSH]

TABLE X: Concepts with associated keywords and EMTREE terms used in Embase search strategy.

Concept	Keywords	EMTREE Terms
Foreign Bodies	"foreign obj*" "foreign bod*"	"foreign body"/exp
Intentional Ingestion / Self-harm	"intent*" "deliberate*" "purpose*" "self-injur*" "selfharm*" "self-harm*"	"automutilation"/exp
Ingestion Behavior	"ingest*" "swallow*"	"swallowing"/exp
Interventions	"surg*" "endoscop*" "EGD" "OGD" "Esophagogastroduodenoscopy" "Oesophagogastroduodenoscopy" "manag*"	"endoscopy"/exp "surgery"/exp "conservative treatment"/exp "drug therapy"/exp

TABLE XI: Concepts with associated keywords and Cochrane MeSH terms used in CENTRAL search strategy.

Concept	Keywords	Cochrane MeSH Terms
Foreign Bodies	"foreign obj*" "foreign bod*" (foreign NEXT obj*) (foreign NEXT bod*) intent* deliberate*	[mh foreign bodies]
Intentional Ingestion / Self-harm	purpose* (self NEXT injur*) (self NEXT harm*) ingest* swallow* surg* endoscop*	[mh self-injurious behavior]
Ingestion Behavior	EGD Esophagogastrroduodenoscopy Oesophagogastrroduodenoscopy manag*	-
Interventions		[mh endoscopy] [mh surgical procedures, operative] [mh conservative treatment] [mh drug therapy]

TABLE XII: Concepts with associated keywords and Web of Science fields used in the search strategy.

Concept	Keywords	Search Field
Foreign Bodies	foreign obj* foreign bod* aut mutilation intent* deliberate*	ALL=
Intentional Ingestion / Self-harm	purpose* self-injur* selfharm* self-harm* swallowing	ALL=
Ingestion Behavior	ingest* swallow* endoscopy surgery conservative treatment drug therapy	ALL=
Interventions	surg* endoscop* EGD Esophagogastrroduodenoscopy Oesophagogastrroduodenoscopy manag*	ALL=

TABLE XIII: Concepts with associated keywords and Scopus syntax used in the search strategy.

Concept	Keywords	Search Field / Syntax
Foreign Bodies	foreign PRE/0 obj* foreign PRE/0 bod* intent* deliberate*	ALL()
Intentional Ingestion / Self-harm	purpose* self PRE/0 injur* self PRE/0 harm* ingest* swallow* endoscopy surgery 'conservative' 'treatment' 'drug' 'therapy'	ALL()
Ingestion Behavior	surg* endoscop* egd esophagogastrroduodenoscopy oesophagogastrroduodenoscopy manag*	ALL()
Interventions		ALL()

TABLE XIV: Concepts with associated keywords and controlled vocabulary (Descriptors) used in PsycINFO search strategy.

Concept	Keywords	PsycINFO Descriptors
Foreign Bodies	foreign obj* foreign bod* automutilation intent* deliberate*	—
Intentional Ingestion / Self-harm	purpose* self injur* self harm*	DE "Nonsuicidal Self-Injury"
Ingestion Behavior	ingest* swallow* endoscop* conservative treatment drug therapy	DE "Ingestion"
Interventions	surg* egd esophagogastroduodenoscopy oesophagogastroduodenoscopy manag*	DE "Surgery"

TABLE XV: Concepts with associated keywords used in Google Scholar search strategy.

Concept	Keywords	Search Field
Foreign Bodies	"foreign obj*" "foreign bod*" "intenti*" "deliberate*" "purpose*" "self-injur*" "selfharm*" "self-harm*"	—
Intentional Ingestion / Self-harm	"ingest*" "swallow*"	—
Ingestion Behavior		—

1) PubMed Query:

(“Foreign Bodies”[MeSH] OR “foreign obj*” OR “foreign bod*”)
AND
(“Self-Injurious Behavior”[MeSH] OR “intent*” OR “deliberate*” OR “purpose*” OR “self-injur*” OR “selfharm*” OR “self-harm*”)
AND
(“ingest*” OR “swallow*”)
AND
(“Endoscopy”[MeSH] OR “Surgical Procedures, Operative”[MeSH] OR “Conservative Treatment”[MeSH] OR “Drug Therapy”[MeSH] OR “surg*” OR “endoscop*” OR “EGD” OR “OGD” OR “Esophagogastroduodenoscopy” OR “Oesophagogastroduodenoscopy” OR “manag*”)

2) Embase Query (All Fields):

(‘foreign body’/exp OR “foreign obj*” OR “foreign bod*”)
AND
(‘automutilation’/exp OR “intent*” OR “deliberate*” OR “purpose*” OR “self-injur*” OR “selfharm*” OR “self-harm*”)
AND
(‘swallowing’/exp OR “ingest*” OR “swallow*”)
AND
(‘endoscopy’/exp OR ‘surgery’/exp OR ‘conservative treatment’/exp OR ‘drug therapy’/exp OR “surg*” OR “endoscop*” OR “EGD” OR “OGD” OR “Esophagogastroduodenoscopy” OR “Oesophagogastroduodenoscopy” OR “manag*”)

3) CENTRAL (Cochrane) Query (All Fields):

([mh “foreign bodies”] OR (“foreign” NEXT “obj*”) OR (“foreign” NEXT “bod*”))
AND
([mh “self-injurious behavior”] OR “intent*” OR “deliberate*” OR “purpose*” OR (“self” NEXT “injur*”) OR (“self” NEXT “harm*”))
AND
(“ingest*” OR “swallow*”)
AND
([mh “endoscopy”] OR [mh “surgical procedures, operative”] OR [mh “conservative treatment”] OR [mh “drug therapy”] OR “surg*” OR “endoscop*” OR “EGD” OR “Esophagogastroduodenoscopy” OR “Oesophagogastroduodenoscopy” OR “manag*”)

4) Web of Science Query: Link: <https://www.webofscience.com/wos/woscc/summary/4da44d48-3e09-4a94-a3bd-ff8139e94859-01387ccd63/relevance/1>

ALL=(“foreign obj*” OR “foreign bod*”)
AND
ALL=(“automutilation” OR “intent*” OR “deliberate*” OR “purpose*” OR “self-injur*” OR “selfharm*” OR “self-harm*”)
AND
ALL=(“swallowing” OR “ingest*” OR “swallow*”)
AND
ALL=(“endoscopy” OR “surgery” OR “conservative treatment” OR “drug therapy” OR “surg*” OR “endoscop*” OR “EGD” OR “Esophagogastroduodenoscopy” OR “Oesophagogastroduodenoscopy” OR “manag*”)

5) Scopus Query:

ALL (“foreign PRE/0 obj*” OR “foreign PRE/0 bod*”)
AND
ALL (“intent*” OR “deliberate*” OR “purpose*” OR “self PRE/0 injur*” OR “self PRE/0 harm*”)
AND
ALL (“ingest*” OR “swallow*”)
AND
ALL (“endoscopy” OR “surgery” OR “conservative” OR “treatment” OR “drug” OR “therapy” OR “surg*” OR “endoscop*” OR “EGD” OR “Esophagogastroduodenoscopy” OR “Oesophagogastroduodenoscopy” OR “manag*”)

6) *PsycINFO Query:* **Link:** [https://search-ebscohost-com.ezproxy.library.qmul.ac.uk/login.aspx?direct=true&db=psyh&bquery=\(foreign+obj*+OR+foreign+bod*\)+AND+\(DE+%26quot%3bNonsuicidal+Self-Injury%26quot%3b+OR+automutilation+OR+intent*+OR+deliberate*+OR+purpose*+OR+self+injur*+OR+self+harm*\)+AND+\(DE+%26quot%3bIngestion%26quot%3b+OR+ingest*+OR+swallow*\)+AND+\(DE+%26quot%3bSurgery%26quot%3b+OR+endoscop*+OR+conservative+treatment+OR+drug+therapy+surg*+OR+endoscop*+OR+egd+OR+esophagogastroduodenoscopy+OR+oesophagogastroduodenoscopy+OR+manag*\)&type=0&searchMode=Standard&site=ehost-live](https://search-ebscohost-com.ezproxy.library.qmul.ac.uk/login.aspx?direct=true&db=psyh&bquery=(foreign+obj*+OR+foreign+bod*)+AND+(DE+%26quot%3bNonsuicidal+Self-Injury%26quot%3b+OR+automutilation+OR+intent*+OR+deliberate*+OR+purpose*+OR+self+injur*+OR+self+harm*)+AND+(DE+%26quot%3bIngestion%26quot%3b+OR+ingest*+OR+swallow*)+AND+(DE+%26quot%3bSurgery%26quot%3b+OR+endoscop*+OR+conservative+treatment+OR+drug+therapy+surg*+OR+endoscop*+OR+egd+OR+esophagogastroduodenoscopy+OR+oesophagogastroduodenoscopy+OR+manag*)&type=0&searchMode=Standard&site=ehost-live)

(“foreign obj*” OR “foreign bod*”)

AND

(DE “Nonsuicidal Self-Injury” OR “automutilation” OR “intent*” OR “deliberate*” OR “purpose*” OR “self injur*” OR “self harm*”)

AND

(DE “Ingestion” OR “ingest*” OR “swallow*”)

AND

(DE “Surgery” OR “endoscop*” OR “conservative treatment” OR “drug therapy” OR “surg*” OR “endoscop*” OR “EGD” OR “Esophagogastroduodenoscopy” OR “Oesophagogastroduodenoscopy” OR “manag*”)

C. Grey Literature

Google Scholar:

(“foreign obj*” OR “foreign bod*”)

AND

(“intent*” OR “deliberate*” OR “purpose*” OR “self-injur*” OR “selfharm*” OR “self-harm*”)

AND

(“ingest*” OR “swallow*”)

D. Screening Process

A total of 808 records were identified through initial database searches: PubMed (317), Web of Science (277), Google Scholar (135), Embase (25), SCOPUS (24), PsycINFO (16), and Cochrane (14). 316 duplicates were identified and removed.

Title and abstract screening was undertaken, with JGE reviewing all 492 records. A random sample of 50 records was generated for independent screening MS. Cohen's Kappa was calculated for inter-reviewer agreement between JGE and MS, yielding a value of 0.38, indicating fair agreement. Where JGE and MS disagreed, 16 records were reviewed by GC. In total, 176 records were excluded, leaving 316 for full text review.

During full text review, JGE reviewed all 316 records. A random sample of 32 records was generated for independent review by MS. Inter-reviewer agreement was again calculated using Cohen's Kappa, yielding a value of 0.21, indicating fair agreement. Where JGE and MS disagreed, 5 records were reviewed by GC. In total, 276 records were excluded during full text review. 40 records were included and proceeded to bibliography search.

The bibliographies of the 40 included papers were searched by manually JGE. Relevant bibliography items were identified, collated, and evaluated against the eligibility criteria, yielding 194 results.

These 194 results were reviewed by JGE. 164 bibliography search records were excluded, leaving 30 for inclusion.

Therefore, a total of 70 records were included in this study and proceeded to bias assessment. This process is illustrated in Figure 1.

E. Data Extraction

TABLE XVI: Variables used for case report data extraction. Aggregates of which where used to create Variable_Rate and Variable_Cases.

Variable	Definition
Is_Prisoner	Documented in prison, police custody, or detained (including immigration detention) at the time of the encounter; 'N' if not detained; 'UK' if unknown.
Psych_Hx	Documented DSM-V mental disorder (including substance-related disorders) [129]; 'N' if no diagnosis; 'UK' if data unavailable.
Is_Displaced_Person	'Y' if: meets the UN General Assembly [130] definition of 'Refugee'; or meets UNHCR [131] definition of an 'internally displaced person'; or meets the UNHCR [132] definition for 'asylum seeker'; 'N' if not displaced; 'UK' if unknown.
Under_Influence_Alcohol	Evidence, suspicion, or self-report of alcohol influence at presentation; 'N' if no indication; 'UK' if unknown.
Is_Psych_Inpat	Admitted (voluntarily or involuntarily) to a psychiatric facility/ward at encounter; 'N' if not admitted; 'UK' if unknown.
Severe_Disability_Hx	History of severe learning disability or impaired consciousness; 'N' if absent; 'UK' if unknown.
Previous_Ingestions	Prior episode of foreign-body ingestion documented; 'N' if first ingestion; 'UK' if history unknown.
Motivation_Intent_To_Harm	Ingestion intended for self-harm, self-injury, or suicide; 'N' if other motive; 'UK' if unclear.
Motivation_Protest	Ingestion as protest, demonstration, or manipulation (e.g., objection to detention conditions); 'N' if not protest-related; 'UK' if unclear.
Motivation_Psychiatric	Ingestion driven primarily by an underlying psychiatric condition (psychosis, impulsivity, etc.); 'N' if not psychiatric; 'UK' if unclear.
Motivation_Psychosocial	Ingestion motivated by social or interpersonal factors (imitative acts, shock value, body-image, safekeeping, etc.); 'N' if not psychosocial; 'UK' if unclear.
Motivation_Uncertain	No clear motivation identified in documentation; 'N' if specific motive recorded; 'UK' if ambiguous.
Object_Button_Battery	Button battery ingested; 'N' if not; 'UK' if object type not recorded.
Object_Magnet	Magnet ingested; 'N' if none; 'UK' if unknown.
Object_Long	Ingested object length > 5 cm; 'N' if ≤ 5 cm; 'UK' if dimensions unknown.
Object_Sharp	Object described as sharp or pointed (e.g., blades, nails, needles); 'N' if not sharp; 'UK' if unclear.
Object_Multiple	More than one object ingested in same episode; 'N' for single object; 'UK' if number unspecified.
Object_Uncertain	Where object characteristics are unknown. 'N' if known; 'UK' if Unknown.
Outcome_Endoscopy	Endoscopic intervention performed during episode; 'N' if not; 'UK' if unavailable.
Outcome_Surgery	Surgical intervention performed (operative procedure under anaesthesia); 'N' if not; 'UK' if not documented.
Outcome_Conservative	'Y' if managed without endoscopy or surgery; 'N' if either procedure performed.
Outcome_Death	Death causally related to ingestion complications; 'N' if survived; 'UK' if outcome unknown.
Outcome_Complication	'Y' if any complication directly related to ingestion or resulting from management strategy; 'N' if no complication; 'UK' if unknown.
Outcome_Uncertain	Where no outcome identified; 'N' if outcome identified; 'UK' if Unknown.

1) Definitions: Full variable definitions are shown in XVI.

For the purposes of this study, "surgery" was defined as "any operative intervention performed in a sterile operating theatre under general or regional anaesthesia, involving incision or surgical access to body cavities (including laparotomy, laparoscopy, thoracotomy, or cervical exploration) for the purpose of removing an ingested object or managing complications of ingestion". Procedures performed "solely via flexible or rigid endoscopy through natural orifices" were categorised as "endoscopy" and not considered surgical interventions.

2) *Process*: Data were initially extracted by a single reviewer (JGE) into *Microsoft Excel* [56]. Variables for extraction were developed iteratively through engagement with the literature and analysis of consistent reporting patterns. A preliminary review of the first 30 case reports informed the development of additional data categories, which were subsequently applied to the remaining reports.

Following initial extraction, data were imported into *Python* [1] for further processing and analysis. The Python-based pipeline included data cleaning, validation, and transformation to ensure consistency across heterogeneous study formats. These structured data were then used to guide the extraction of aggregate data from case series. Studies were grouped for extraction based on their classification as case reports or case series. Where case series contained sufficiently granular data, cases were extracted individually and treated as case reports; otherwise, data were extracted at the aggregate level. Case grouping for analysis followed the criteria for inclusion as individual case reports or case series, as defined above. Relevant data from reviews and other literature types were recorded under the case report category.

F. Computational Risk of Bias Assessment

To reduce bias dilution of intentionality effect, a novel computation risk of bias assessment was undertaken, using a combination of human review followed by computational risk of bias assessment. First, the author (JGE) extracted data into Microsoft Excel [56]. Then, a computation risk of bias filter was applied to extracted case report and case series data. That process is outlined in this appendix.

1) *Case Reports*: For case reports, the JBI Checklist for Case Reports was used. This tool assesses eight domains of reporting quality, including whether patient demographics were clearly described, a timeline of clinical history was provided, the presenting condition and diagnostic assessment were outlined, and whether the intervention, post-intervention condition, and any adverse events were reported. The final domain evaluates whether the case provides meaningful takeaway lessons.

In addition to manual JBI appraisal, a logic-based validation filter was applied to all case reports using *Python Pandas* [5]. This secondary filter assessed whether key variables — specifically, outcomes, object characteristics, and motivation — were completely unreported. For each domain, a binary flag was generated:

- *Outcome_Unknown* was marked 1 if all outcome-related fields were either missing or marked as unknown.
- *Object_Unknown* was marked 1 if all object-related fields (excluding *Object_Other_Long*) were missing or unknown.
- *Motivation_Unknown* was predefined in the dataset and indicated absence of motivational information.

If any of these flags were triggered, the corresponding JBI item most affected by the missing domain was marked as not reported (e.g., *Post_Intervention_Condition_Described* or *History_Timeline* set to N). Finally, an *Overall_Appraisal* score of *Exclude* was assigned, indicating high risk of bias and exclusion from analysis. This ensured that only case reports with sufficient information to meaningfully contribute to the review question were retained.

2) *Case Series*: For case series, the JBI Checklist for Case Series was applied. The JBI Checklist for Case Series assesses 10 domains of methodological and reporting quality. These include whether the case series defined clear inclusion criteria, applied valid and consistent methods to identify the condition, and included participants consecutively and completely. The checklist also evaluates whether participant demographics and clinical information were clearly reported, whether outcomes or follow-up results were adequately described, and whether the study setting was detailed. Finally, it considers whether the statistical analysis used was appropriate for the data presented.

In addition to manual JBI appraisal, a logic-based exclusion filter was applied using *Python Pandas* [5]. This filter assessed whether key variables — specifically, motivation, object characteristics, and outcomes — were unreported for the entire study population. For each of these domains, a derived rate variable was calculated:

- *Outcome_Unknown_Rate* was marked as 1 if all outcome-related fields were missing or marked as unknown (i.e. the entire population had an had an unknown outcome).
- *Motivation_Unknown_Rate* indicated whether motivation was absent or only partially reported across cases within the study.
- *Object_Unknown_Rate* was derived if all object-related fields were missing or unknown.

If any of these indicators were flagged, the corresponding JBI checklist item (e.g., *Clear_Outcome_Followup_Reported*, *Clear_Demographic_Reported*, or *Clear_Clinical_Info_Reported*) was marked as N, and the study received an *Overall_Appraisal* of *Exclude*. This logic-based validation ensured that case series lacking essential variables could be systematically excluded from the final analysis, maintaining consistency with the review question and minimising risk of bias in the dataset.