

# Lecture 2: Basics of Probability Theory

Professor Ilias Bilonis



## The obvious rule of probability

# The obvious rule

$$p(A \cup \bar{A}) = 1$$

The obvious rule:

$$\underline{p(A | I)} + \underline{p(\neg A | I)} = \underline{1}$$





Venn  
Diagram

# Example: Drawing balls from a box without replacement

Consider the following example of prior information I:


*We are given a box with 10 balls 6 of which are red and 4 of which are blue. The box is sufficiently mixed so that when we get a ball from it, we don't know which one we pick. When we take a ball out of the box, we do not put it back.*



# Example: Drawing balls from a box without replacement

Consider the following example of prior information I:

*We are given a box with 10 balls 6 of which are red and 4 of which are blue. The box is sufficiently mixed so that when we get a ball from it, we don't know which one we pick. When we take a ball out of the box, we do not put it back.*



Let  $B_1$  be the sentence:

*The first ball we draw is blue.*

Let's compute a probability:


$$p(B_1 | I) = \frac{4}{10}$$

*Insufficient Reason  
Laplace.*

# Example: Drawing balls from a box without replacement

Consider the following example of prior information I:

*We are given a box with 10 balls 6 of which are red and 4 of which are blue. The box is sufficiently mixed so that when we get a ball from it, we don't know which one we pick. When we take a ball out of the box, we do not put it back.*



$$p(B_1 | I) = 0.4$$

What is the probability that we get a red ball ( $\neg B_1 \equiv R_1$  is true)?

$$p(R_1 | I) = p(\neg B_1 | I) \stackrel{\text{ob.}}{=} 1 - p(B_1 | I)$$

$$= 1 - 0.4 = 0.6$$

$$p(R_1 | I) = \frac{6}{10}$$