Sequences and Series

Jack Krebsbach

Oct 4

0.1 Sequences and Series

Definition 0.1.1: Sequence

A sequence is a function from $f: \mathbb{N} \to \mathbb{R}$. Examples:

- 1. (a_n)
- 2. $(a_1, a_2, a_3, \ldots, a_n)$

Definition 0.1.2: Convergence

A sequence, (a_n) , converges to a point, x, if for all $\epsilon > 0$ there exist $N \in \mathbb{N}$ such that for all n > N, $|a_n - x| < \epsilon$.

Theorem 0.1.1 Uniqueness of Limits.

The limit of a sequence, when it exists, must me unique.

Proof: Let (x_n) be a convergent series that converges to x. By way of contradiction, suppose that $(x_n) \to y$ where $x \neq y$ and x < y. Let $\epsilon = \frac{1}{3}(y - x)$. Since (x_n) converges to x there exists $N_x \in \mathbb{N}$ such that for all $n > N_x$, $|x_n - x| < \epsilon$. Similarly, since (x_n) converges to y there exists $N_Y \in \mathbb{N}$ such that for all $n > N_y$, $|x_n - y| < \epsilon$.

Let $N = \max\{N_x, N_y\}$. Then $x_{N+2} \in \mathcal{B}(x, \epsilon) \cap \mathcal{B}(y, \epsilon)$. This is a contradiction, $x_{N+2} \notin \mathcal{B}(x, \epsilon) \cap \mathcal{B}(y, \epsilon)$. Thus, x = y and limits are unique!

Theorem 0.1.2 Convergent Sequences are Bounded

Proof: Let (x_n) be a convergent sequence converging to x. Let $\epsilon = 1$. Since (x_n) converges there exists $N \in \mathbb{N}$ such that for all n > N, $|x_n - x| < \epsilon$. By the triangle inequality theorem, $|x_n| - |x| \le |x_n - x| < 1$. So $|x_n| < |x| + 1$ for all n > N.

Now consider the set $\{|x_1|, |x_2|, |x_3|, \dots, |x_N|\}$. All, the elements outside the ball of convergence. Let $B = \{|x_1|, |x_2|, |x_3|, \dots, |x_N|, |x_1| + 1\}$. Thus, $|x_n| \le B$ for all $n \in \mathbb{N}$ and (x_n) is bounded.

☺

Theorem 0.1.3 Algebraic Limit Theorem

Let $\lim a_n = a$, and $\lim b_n = b$. Then,

- 1. $\lim (ca_n) = ca$, for all $c \in \mathbb{R}$;
- 2. $\lim (a_n + b_n) = a + b$;
- 3. $\lim (a_n b_n) = ab$;
- 4. $\lim (a_n/b_n) = a/b$, provided $b \neq 0$.