

[13.32] Show that every finite group \mathbf{G} has a faithful representation in $GL(n)$ where n is the order of \mathbf{G} .

Solution

A **representation** is a function $T: \mathbf{G} \rightarrow GL(n)$ that preserves the group structure; i.e., for all $g_i, g_j \in \mathbf{G}$, $T(g_i)T(g_j) = T(g_i g_j)$. T is **faithful** if it is one-to-one; i.e., if $T(g_i) = T(g_j) \Rightarrow g_i = g_j$.

Part A. Show T is a representation

I do this 2 ways. I do it my way first, then I repeat it using Robin's method which is very slick. For motivation, I use Penrose's hint to label the matrix for $T(g_i)$.

$$T(g_i) = g_j \begin{matrix} & & & g_k & & \\ \begin{bmatrix} T(g_i)_1^1 & T(g_i)_2^1 & \cdots & T(g_i)_k^1 & \cdots & T(g_i)_n^1 \\ \vdots & \vdots & & \vdots & & \vdots \\ T(g_i)_1^j & T(g_i)_2^j & \cdots & T(g_i)_k^j & \cdots & T(g_i)_n^j \\ \vdots & \vdots & & \vdots & & \vdots \\ T(g_i)_1^n & T(g_i)_2^n & \cdots & T(g_i)_k^n & \cdots & T(g_i)_n^n \end{bmatrix} \end{matrix}$$

Given k there is a unique j such that $g_k = g_i g_j$ (namely, $g_j = g_i^{-1} g_k$). So I think of this matrix as g_i taking g_j to g_k . (Robin had g_i take g_k to g_j , which also works, as does right multiplication by g_i .) Thus I make the definition

$$(1) \quad T(g_i) \equiv \begin{cases} 1 & \text{if } i, j, \text{ and } k \text{ are such that } g_k = g_i g_j \\ 0 & \text{Otherwise} \end{cases}$$

which in matrix notation is

$$T(g_i) \begin{bmatrix} 0 \\ \vdots \\ 1_j \\ \vdots \\ 0 \end{bmatrix} = \begin{bmatrix} T(g_i)_j^1 \\ \vdots \\ T(g_i)_j^k \\ \vdots \\ T(g_i)_j^n \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 1_k \\ \vdots \\ 0 \end{bmatrix}$$

It suffices to show that $T(g_i)T(g_j)$ agrees with $T(g_i g_j)$ on each basis element

$$\begin{bmatrix} 0 \\ \vdots \\ 1_r \\ \vdots \\ 0 \end{bmatrix}.$$

From definition (1) we get that

$$(2) \quad T(g_j)_r^s = \begin{cases} 1 & \text{if } j, r, \text{ and } s \text{ are such that } g_s = g_j g_r \\ 0 & \text{Otherwise} \end{cases}, \text{ or } T(g_j) \begin{bmatrix} 0 \\ \vdots \\ 1_r \\ \vdots \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 1_s \\ \vdots \\ 0 \end{bmatrix}$$

and

$$(3) \quad T(g_i)_s^t = \begin{cases} 1 & \text{if } i, s, \text{ and } t \text{ are such that } g_t = g_i g_s \\ 0 & \text{Otherwise} \end{cases}, \text{ or } T(g_i) \begin{bmatrix} 0 \\ \vdots \\ 1_s \\ \vdots \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 1_t \\ \vdots \\ 0 \end{bmatrix}.$$

Thus,

$$[T(g_i)T(g_j)] \begin{bmatrix} 0 \\ \vdots \\ 1_r \\ \vdots \\ 0 \end{bmatrix} = T(g_i) \left(T(g_j) \begin{bmatrix} 0 \\ \vdots \\ 1_r \\ \vdots \\ 0 \end{bmatrix} \right) \stackrel{(2)}{=} T(g_i) \begin{bmatrix} 0 \\ \vdots \\ 1_s \\ \vdots \\ 0 \end{bmatrix} \stackrel{(3)}{=} \begin{bmatrix} 0 \\ \vdots \\ 1_t \\ \vdots \\ 0 \end{bmatrix}.$$

Again, from definition (1), we get that

$$(4) \quad T(g_k)_r^p = \begin{cases} 1 & \text{if } k, r, \text{ and } p \text{ are such that } g_p = g_k g_r \\ 0 & \text{Otherwise} \end{cases}, \text{ or}$$

$$T(g_k) \begin{bmatrix} 0 \\ \vdots \\ 1_r \\ \vdots \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 1_p \\ \vdots \\ 0 \end{bmatrix}. \text{ So } g_p = g_k g_r = (g_i g_j) g_r = g_i (g_j g_r) = g_i g_s = g_t; \text{ i.e., } p = t.$$

$$\text{So, } T(g_i g_j) \begin{bmatrix} 0 \\ \vdots \\ 1_r \\ \vdots \\ 0 \end{bmatrix} = T(g_k) \begin{bmatrix} 0 \\ \vdots \\ 1_r \\ \vdots \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 1_p \\ \vdots \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 1_t \\ \vdots \\ 0 \end{bmatrix} = [T(g_i) T(g_j)] \begin{bmatrix} 0 \\ \vdots \\ 1_r \\ \vdots \\ 0 \end{bmatrix}.$$

That is, $T(g_i) T(g_j) = T(g_i g_j)$. ✓

Alternate proof of Part A

$$[T(g_i) T(g_j)]_b^a = T(g_i)_c^a T(g_j)_b^c = \begin{cases} 1 & \text{if } i, c, a, j, \text{ and } b \text{ satisfy} \\ & g_a = g_i g_c \text{ and } g_c = g_j g_b \\ 0 & \text{Otherwise} \end{cases}.$$

$g_a = g_i g_c$ and $g_c = g_j g_b \Leftrightarrow g_i^{-1} g_a = g_j g_b \Leftrightarrow g_a g_b^{-1} = g_i g_j$. Therefore

$$[T(g_i) T(g_j)]_b^a = \begin{cases} 1 & \text{if } a, b, i, \text{ and } j \text{ satisfy } g_a g_b^{-1} = g_i g_j \\ 0 & \text{Otherwise} \end{cases}. \text{ But,}$$

$$T(g_i g_j)_b^a = \begin{cases} 1 & \text{if } a, b, i, \text{ and } j \text{ satisfy } g_a = (g_i g_j) g_b \Leftrightarrow g_a g_b^{-1} = g_i g_j \\ 0 & \text{Otherwise} \end{cases} \quad \checkmark$$

Part B Show T is faithful

We suppose $T(g_i) = T(g_j)$ and we must show that $g_i = g_j$. Since $T(g_i) = T(g_j)$,

$$\forall a, b \quad T(g_i)_b^a = T(g_j)_b^a \Leftrightarrow T(g_i)_b^a = 1 \text{ if and only if } T(g_j)_b^a = 1.$$

$T(g_i)_b^a = 1$ iff $g_a = g_i g_b$ and $T(g_j)_b^a = 1$ iff $g_a = g_j g_b$. So for all a and b we have $g_a = g_i g_b$ iff $g_a = g_j g_b$. Letting $b = e$ (the identity of \mathbf{G}) we get

$$g_i = g_a \text{ iff } g_j = g_a \Leftrightarrow g_i = g_j. \quad \checkmark$$