


## Chapter 11: Linear Models and Estimation by Least Squares

- 11.1 Using the hint,  $\hat{y}(\bar{x}) = \hat{\beta}_0 + \hat{\beta}_1 \bar{x} = (\bar{y} - \hat{\beta}_1 \bar{x}) + \hat{\beta}_1 \bar{x} = \bar{y}$ .
- 11.2
- a. slope = 0, intercept = 1. SSE = 6.
  - b. The line with a negative slope should exhibit a better fit.
  - c. SSE decreases when the slope changes from .8 to .7. The line is pivoting around the point (0, 1), and this is consistent with  $(\bar{x}, \bar{y})$  from part Ex. 11.1.
  - d. The best fit is:  $y = 1.000 + 0.700x$ .
- 11.3 The summary statistics are:  $\bar{x} = 0$ ,  $\bar{y} = 1.5$ ,  $S_{xy} = -6$ ,  $S_{xx} = 10$ . Thus,  $\hat{y} = 1.5 - .6x$ .




The graph is above.

- 11.4 The summary statistics are:  $\bar{x} = 72$ ,  $\bar{y} = 72.1$ ,  $S_{xy} = 54,243$ ,  $S_{xx} = 54,714$ . Thus,  $\hat{y} = 0.72 + 0.99x$ . When  $x = 100$ , the best estimate of  $y$  is  $\hat{y} = 0.72 + 0.99(100) = 99.72$ .
- 11.5 The summary statistics are:  $\bar{x} = 4.5$ ,  $\bar{y} = 43.3625$ ,  $S_{xy} = 203.35$ ,  $S_{xx} = 42$ . Thus,  $\hat{y} = 21.575 + 4.842x$ . Since the slope is positive, this suggests an increase in median prices over time. Also, the expected annual increase is \$4,842.
- 11.6
- a. intercept = 43.362, SSE = 1002.839.
  - b. the data show an increasing trend, so a line with a negative slope would not fit well.
  - c. Answers vary.
  - d. Answers vary.
  - e. (4.5, 43.3625)
  - f. The sum of the areas is the SSE.
- 11.7
- a. The relationship appears to be proportional to  $x^2$ .
  - b. No.
  - c. No, it is the best *linear* model.

**11.8** The summary statistics are:  $\bar{x} = 15.505$ ,  $\bar{y} = 9.448$ ,  $S_{xy} = 1546.459$ ,  $S_{xx} = 2359.929$ . Thus,  $\hat{y} = -0.712 + 0.655x$ . When  $x = 12$ , the best estimate of  $y$  is  $\hat{y} = -0.712 + 0.655(12) = 7.148$ .

**11.9** a. See part c.  
b.  $\hat{y} = -15.45 + 65.17x$ .



c. The graph is above.


d. When  $x = 1.9$ , the best estimate of  $y$  is  $\hat{y} = -15.45 + 65.17(1.9) = 108.373$ .

**11.10**  $\frac{dSSE}{d\beta_1} = -2\sum_{i=1}^n (y_i - \beta_1 x_i)x_i = -2\sum_{i=1}^n (x_i y_i - \beta_1 x_i^2) = 0$ , so  $\hat{\beta}_1 = \frac{\sum_{i=1}^n x_i y_i}{\sum_{i=1}^n x_i^2}$ .

**11.11** Since  $\sum_{i=1}^n x_i y_i = 134,542$  and  $\sum_{i=1}^n x_i^2 = 53,514$ ,  $\hat{\beta}_1 = 2.514$ .

**11.12** The summary statistics are:  $\bar{x} = 20.4$ ,  $\bar{y} = 12.94$ ,  $S_{xy} = -425.571$ ,  $S_{xx} = 1859.2$ .


a. The least squares line is:  $\hat{y} = 17.609 - 0.229x$ .



- b. The line provides a reasonable fit.  
c. When  $x = 20$ , the best estimate of  $y$  is  $\hat{y} = 17.609 - 0.229(20) = 13.029$  lbs.

**11.13** The summary statistics are:  $\bar{x} = 6.177$ ,  $\bar{y} = 270.5$ ,  $S_{xy} = -5830.04$ ,  $S_{xx} = 198.29$ .


- a. The least squares line is:  $\hat{y} = 452.119 - 29.402x$ .



- b. The graph is above.

**11.14** The summary statistics are:  $\bar{x} = .325$ ,  $\bar{y} = .755$ ,  $S_{xy} = -.27125$ ,  $S_{xx} = .20625$

- a. The least squares line is:  $\hat{y} = 1.182 - 1.315x$ .



b. The graph is above. The line provides a reasonable fit to the data.


$$\begin{aligned}
 11.15 \quad \text{a. } SSE &= \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2 = \sum_{i=1}^n [y_i - \bar{y} - \hat{\beta}_1 (x_i - \bar{x})]^2 = \sum_{i=1}^n (y_i - \bar{y})^2 \\
 &\quad + \hat{\beta}_1^2 \sum_{i=1}^n (x_i - \bar{x})^2 - 2\hat{\beta}_1 \sum_{i=1}^n (y_i - \bar{y})(x_i - \bar{x}) \\
 &= \sum_{i=1}^n (y_i - \bar{y})^2 + \hat{\beta}_1 S_{xy} - 2\hat{\beta}_1 S_{xy} = S_{yy} - \hat{\beta}_1 S_{xy}.
 \end{aligned}$$

b. Since  $SSE = S_{yy} - \hat{\beta}_1 S_{xy}$ ,

$$\begin{aligned}
 S_{yy} &= SSE + \hat{\beta}_1 S_{xy} = SSE + (S_{xy})^2 / S_{xx}. \quad \text{But, } S_{xx} > 0 \text{ and } (S_{xy})^2 \geq 0. \text{ So,} \\
 S_{yy} &\geq SSE.
 \end{aligned}$$

11.16 The summary statistics are:  $\bar{x} = 60$ ,  $\bar{y} = 27$ ,  $S_{xy} = -1900$ ,  $S_{xx} = 6000$ .

a. The least squares line is:  $\hat{y} = 46.0 - .31667x$ .



b. The graph is above.

c. Using the result in Ex. 11.15(a),  $SSE = S_{yy} - \hat{\beta}_1 S_{xy} = 792 - (-.31667)(-1900) = 190.327$ . So,  $s^2 = 190.327/10 = 19.033$ .

**11.17** a. With  $S_{yy} = 1002.8388$  and  $S_{xy} = 203.35$ ,  $SSE = 1002.8388 - 4.842(203.35) = 18.286$ . So,  $s^2 = 18.286/6 = 3.048$ .


b. The fitted line is  $\hat{y} = 43.35 + 2.42x^*$ . The same answer for SSE (and thus  $s^2$ ) is found.

**11.18** a. For Ex. 11.8,  $S_{yy} = 1101.1686$  and  $S_{xy} = 1546.459$ ,  $SSE = 1101.1686 - .6552528(1546.459) = 87.84701$ . So,  $s^2 = 87.84701/8 = 10.98$ .

b. Using the coding  $x_i^* = x_i - \bar{x}$ , the fitted line is  $\hat{y} = 9.448 + .655x^*$ . The same answer for  $s^2$  is found.

**11.19** The summary statistics are:  $\bar{x} = 16$ ,  $\bar{y} = 10.6$ ,  $S_{xy} = 152.0$ ,  $S_{xx} = 320$ .

a. The least squares line is:  $\hat{y} = 3.00 + 4.75x$ .



b. The graph is above.

c.  $s^2 = 5.025$ .

**11.20** The likelihood function is given by,  $K = (\sigma\sqrt{2\pi})^n$ ,

$$L(\beta_0, \beta_1) = K \exp\left[-\frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2\right], \text{ so that}$$

$$\ln L(\beta_0, \beta_1) = \ln K - \frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2.$$

Note that maximizing the likelihood (or equivalently the log-likelihood) with respect to  $\beta_0$  and  $\beta_1$  is identical to minimizing the positive quantity  $\sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2$ . This is the least-squares criterion, so the estimators will be the same.

**11.21** Using the results of this section and Theorem 5.12,

$$\text{Cov}(\hat{\beta}_0, \hat{\beta}_1) = \text{Cov}(\bar{Y} - \hat{\beta}_1 \bar{x}, \hat{\beta}_1) = \text{Cov}(\bar{Y}, \hat{\beta}_1) - \text{Cov}(\hat{\beta}_1 \bar{x}, \hat{\beta}_1) = 0 - \bar{x}V(\hat{\beta}_1).$$

Thus,  $\text{Cov}(\hat{\beta}_0, \hat{\beta}_1) = -\bar{x}\sigma^2 / S_{xx}$ . Note that if  $\sum_{i=1}^n x_i = 0$ ,  $\bar{x} = 0$  so  $\text{Cov}(\hat{\beta}_0, \hat{\beta}_1) = 0$ .

**11.22** From Ex. 11.20, let  $\theta = \sigma^2$  so that the log-likelihood is

$$\ln L(\theta) = -\frac{n}{2} \ln(2\pi) - \frac{n}{2} \ln \theta - \frac{1}{2\theta} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2.$$

Thus,

$$\frac{d}{d\theta} \ln L(\theta) = -\frac{n}{2\theta} + \frac{1}{2\theta^2} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2.$$

The MLE is  $\hat{\theta} = \hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2$ , but since  $\beta_0$  and  $\beta_1$  are unknown, we can insert their MLEs from Ex. 11.20 to obtain:

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2 = \frac{1}{n} \text{SSE}.$$

- 11.23** From Ex. 11.3, it is found that  $S_{yy} = 4.0$
- Since  $SSE = 4 - (-.6)(-.6) = .4$ ,  $s^2 = .4/3 = .1333$ . To test  $H_0: \beta_1 = 0$  vs.  $H_a: \beta_1 \neq 0$ ,  $|t| = \frac{|-.6|}{\sqrt{.1333(.1)}} = 5.20$  with 3 degrees of freedom. Since  $t_{.025} = 3.182$ , we can reject  $H_0$ .
  - Since  $t_{.005} = 5.841$  and  $t_{.01} = 4.541$ ,  $.01 < p\text{-value} < .02$ .  
Using the Applet,  $2P(T > 5.20) = 2(.00691) = .01382$ .
  - $-.6 \pm 3.182\sqrt{.1333}\sqrt{.1} = -.6 \pm .367$  or  $(-.967, -.233)$ .
- 11.24** To test  $H_0: \beta_1 = 0$  vs.  $H_a: \beta_1 \neq 0$ ,  $SSE = 61,667.66$  and  $s^2 = 5138.97$ . Then,  
 $|t| = \frac{|-29.402|}{\sqrt{5138.97(.005043)}} = 5.775$  with 12 degrees of freedom.
- From Table 5,  $P(|T| > 3.055) = 2(.005) = .01 > p\text{-value}$ .
  - Using the Applet,  $2P(T > 5.775) = .00008$ .
  - Reject  $H_0$ .
- 11.25** From Ex. 11.19, to test  $H_0: \beta_1 = 0$  vs.  $H_a: \beta_1 \neq 0$ ,  $s^2 = 5.025$  and  $S_{xx} = 320$ . Then,  
 $|t| = \frac{|.475|}{\sqrt{5.025/320}} = 3.791$  with 8 degrees of freedom.
- From Table 5,  $P(|T| > 3.355) = 2(.005) = .01 > p\text{-value}$ .
  - Using the Applet,  $2P(T > 3.791) = 2(.00265) = .0053$ .
  - Reject  $H_0$ .
  - We cannot assume the linear trend continues – the number of errors could level off at some point.
  - A 95% CI for  $\beta_1$ :  $.475 \pm 2.306\sqrt{5.025/320} = .475 \pm .289$  or  $(.186, .764)$ . We are 95% confident that the expected change in number of errors for an hour increase of lost sleep is between  $(.186, .764)$ .
- 11.26** The summary statistics are:  $\bar{x} = 53.9$ ,  $\bar{y} = 7.1$ ,  $S_{xy} = 198.94$ ,  $S_{xx} = 1680.69$ ,  $S_{yy} = 23.6$ .
- The least squares line is:  $\hat{y} = 0.72 + 0.118x$ .
  - $SSE = 23.6 - .118(198.94) = .125$  so  $s^2 = .013$ . A 95% CI for  $\beta_1$  is  $0.118 \pm 2.776\sqrt{.013}\sqrt{.00059} = 0.118 \pm .008$ .
  - When  $x = 0$ ,  $E(Y) = \beta_0 + \beta_1(0) = \beta_0$ . So, to test  $H_0: \beta_0 = 0$  vs.  $H_a: \beta_0 \neq 0$ , the test statistic is  $|t| = \frac{|.72|}{\sqrt{.013}\sqrt{1.895}} = 4.587$  with 4 degrees of freedom. Since  $t_{.005} = 4.604$  and  $t_{.01} = 3.747$ , we know that  $.01 < p\text{-value} < .02$ .
  - Using the Applet,  $2P(T > 4.587) = 2(.00506) = .01012$ .
  - Reject  $H_0$ .
- 11.27** Assuming that the error terms are independent and normally distributed with 0 mean and constant variance  $\sigma^2$ :
- We know that  $Z = \frac{\hat{\beta}_i - \beta_{i,0}}{\sigma\sqrt{c_{ii}}}$  has a standard normal distribution under  $H_0$ .  
Furthermore,  $V = (n-2)S^2 / \sigma^2$  has a chi-square distribution with  $n-2$  degrees of freedom. Therefore, by Definition 7.2,

$$\frac{Z}{\sqrt{V/(n-2)}} = \frac{\hat{\beta}_i - \beta_{i,0}}{S\sqrt{c_{ii}}}$$

has a  $t$ -distribution with  $n - 2$  degrees of freedom under  $H_0$  for  $i = 1, 2$ .

- b. Using the pivotal quantity expressed above, the result follows from the material in Section 8.8.

**11.28** Restricting to  $\Omega_0$ , the likelihood function is

$$L(\Omega_0) = \frac{1}{(2\pi)^{n/2} \sigma^n} \exp\left[-\frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \beta_0)^2\right].$$

It is not difficult to verify that the MLEs for  $\beta_0$  and  $\sigma^2$  under the restricted space are  $\bar{Y}$  and  $\frac{1}{n} \sum_{i=1}^n (Y_i - \bar{Y})^2$  (respectively). The MLEs have already been found for the unrestricted space so that the LRT simplifies to

$$\lambda = \frac{L(\hat{\Omega}_0)}{L(\hat{\Omega})} = \left( \frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{\sum_{i=1}^n (y_i - \bar{y})^2} \right)^{n/2} = \left( \frac{\text{SSE}}{S_{yy}} \right)^{n/2}.$$

So, we reject if  $\lambda \leq k$ , or equivalently if

$$\frac{S_{yy}}{\text{SSE}} \geq k^{-2/n} = k'.$$

Using the result from 11.15,

$$\frac{\text{SSE} + \hat{\beta}_1 S_{xy}}{\text{SSE}} = 1 + \frac{\hat{\beta}_1 S_{xy}}{\text{SSE}} = 1 + \frac{\hat{\beta}_1^2 S_{xx}}{(n-2)S} = 1 + \frac{T^2}{(n-2)}.$$

So, we see that  $\lambda$  is small whenever  $T = \frac{\hat{\beta}_1}{S\sqrt{c_{11}}}$  is large in magnitude, where  $c_{11} = \frac{1}{S_{xx}}$ .

This is the usual  $t$ -test statistic, so the result has been proven.

**11.29** Let  $\hat{\beta}_1$  and  $\hat{\gamma}_1$  be the least-squares estimators for the linear models  $Y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$  and  $W_i = \gamma_0 + \gamma_1 c_i + \varepsilon_i$  as defined in the problem. Then, we have that:

- $E(\hat{\beta}_1 - \hat{\gamma}_1) = \beta_1 - \gamma_1$
- $V(\hat{\beta}_1 - \hat{\gamma}_1) = \sigma^2 \left( \frac{1}{S_{xx}} + \frac{1}{S_{cc}} \right)$ , where  $S_{cc} = \sum_{i=1}^m (c_i - \bar{c})^2$
- $\hat{\beta}_1 - \hat{\gamma}_1$  follows a normal distribution, so that under  $H_0$ ,  $\beta_1 - \gamma_1 = 0$  so that

$$Z = \frac{\hat{\beta}_1 - \hat{\gamma}_1}{\sigma \sqrt{\left( \frac{1}{S_{xx}} + \frac{1}{S_{cc}} \right)}} \text{ is standard normal}$$

- Let  $V = \text{SSE}_Y + \text{SSE}_W = \sum_{i=1}^n (Y_i - \hat{Y}_i)^2 + \sum_{i=1}^m (W_i - \hat{W}_i)^2$ . Then,  $V / \sigma^2$  has a chi-square distribution with  $n + m - 4$  degrees of freedom
- By Definition 7.2 we can build a random variable with a  $t$ -distribution (under  $H_0$ ):



$$T = \frac{Z}{\sqrt{V/(n+m-4)}} = \frac{\hat{\beta}_1 - \hat{\gamma}_1}{S \sqrt{\left(\frac{1}{S_{xx}} + \frac{1}{S_{cc}}\right)}}, \text{ where } S = (SSE_Y + SSE_W)/(n+m-4).$$

$H_0$  is rejected in favor of  $H_a$  for large values of  $|T|$ .

**11.30 a.** For the first experiment, the computed test statistic for  $H_0: \beta_1 = 0$  vs.  $H_a: \beta_1 \neq 0$  is  $t_1 = (.155)/(.0202) = 7.67$  with 29 degrees of freedom. For the second experiment, the computed test statistic is  $t_2 = (.190)/(.0193) = 9.84$  with 9 degrees of freedom. Both of these values reject the null hypothesis at  $\alpha = .05$ , so we can conclude that the slopes are significantly different from 0.

**b.** Using the result from Ex. 11.29,  $S = (2.04 + 1.86)/(31 + 11 - 4) = .1026$ . We can extract the values of  $S_{xx}$  and  $S_{cc}$  from the given values of  $V(\hat{\beta}_1)$ :

$$S_{xx} = \frac{SSE_Y/(n-2)}{V(\hat{\beta}_1)} = \frac{2.04/29}{(.0202)^2} = 172.397,$$

so similarly  $S_{cc} = 554.825$ . So, to test equality for the slope parameters, the computed test statistic is

$$|t| = \frac{|.155 - .190|}{\sqrt{.1024\left(\frac{1}{172.397} + \frac{1}{554.825}\right)}} = 1.25$$

with 38 degrees of freedom. Since  $t_{.025} \approx z_{.025} = 1.96$ , we fail to reject  $H_0$ : we cannot conclude that the slopes are different.

**11.31** Here, R is used to fit the regression model:

```
> x <- c(19.1, 38.2, 57.3, 76.2, 95, 114, 131, 150, 170)
> y <- c(.095, .174, .256, .348, .429, .500, .580, .651, .722)
> summary(lm(y~x))
```

```
Call:
lm(formula = y ~ x)
```

```
Residuals:
    Min       1Q   Median       3Q      Max
-1.333e-02 -4.278e-03 -2.314e-05  8.056e-03  9.811e-03
```

```
Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)  1.875e-02  6.129e-03   3.059   0.0183 *
x            4.215e-03  5.771e-05  73.040  2.37e-11 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Residual standard error: 0.008376 on 7 degrees of freedom
Multiple R-Squared:  0.9987,    Adjusted R-squared:  0.9985
F-statistic: 5335 on 1 and 7 DF,  p-value: 2.372e-11
```

From the output, the fitted model is  $\hat{y} = .01875 + .004215x$ . To test  $H_0: \beta_1 = 0$  against  $H_a: \beta_1 \neq 0$ , note that the  $p$ -value is quite small indicating a very significant test statistic.

Thus,  $H_0$  is rejected and we can conclude that peak current increases as nickel concentrations increase (note that this is a one-sided alternative, so the  $p$ -value is actually  $2.37\text{e-}11$  divided by 2).

- 11.32** a. From Ex. 11.5,  $\hat{\beta}_1 = 4.8417$  and  $S_{xx} = 42$ . From Ex. 11.15,  $s^2 = 3.0476$  so to test  $H_0: \beta_1 = 0$  vs.  $H_a: \beta_1 > 0$ , the required test statistic is  $t = 17.97$  with 6 degrees of freedom. Since  $t_{.01} = 3.143$ ,  $H_0$  is rejected: there is evidence of an increase.

b. The 99% CI for  $\beta_1$  is  $4.84 \pm 1.00$  or  $(3.84, 5.84)$ .

- 11.33** Using the coded  $x$ 's from 11.18,  $\hat{\beta}_1^* = .655$  and  $s^2 = 10.97$ . Since  $S_{xx} = \sum_{i=1}^{10} (x_i^*)^2 = 2360.2388$ , the computed test statistic is  $|t| = \frac{.655}{\sqrt{\frac{10.97}{2360.2388}}} = 9.62$  with 8 degrees of freedom. Since  $t_{.025} = 2.306$ , we can conclude that there is evidence of a linear relationship.

- 11.34** a. Since  $t_{.005} = 3.355$ , we have that  $p\text{-value} < 2(.005) = .01$ .

b. Using the Applet,  $2P(T < 9.61) = 2(.00001) = .00002$ .

- 11.35** With  $a_0 = 1$  and  $a_1 = x^*$ , the result follows since

$$\begin{aligned} V(\hat{\beta}_0 + \hat{\beta}_1 x^*) &= \frac{1 \cdot \frac{1}{n} \sum_{i=1}^n x_i^2 + (x^*)^2 - 2x^* \bar{x}}{S_{xx}} \sigma^2 = \frac{\frac{1}{n} (\sum_{i=1}^n x_i^2 - n\bar{x}^2) + (x^*)^2 - 2x^* \bar{x} + \bar{x}^2}{S_{xx}} \sigma^2 \\ &= \frac{\frac{1}{n} S_{xx} + (x^* - \bar{x})^2}{S_{xx}} \sigma^2 = \left[ \frac{1}{n} + \frac{(x^* - \bar{x})^2}{S_{xx}} \right] \sigma^2. \end{aligned}$$


This is minimized when  $(x^* - \bar{x})^2 = 0$ , so  $x^* = \bar{x}$ .

- 11.36** From Ex. 11.13 and 11.24, when  $x^* = 5$ ,  $\hat{y} = 452.119 - 29.402(5) = 305.11$  so that  $V(\hat{Y})$  is estimated to be 402.98. Thus, a 90% CI for  $E(Y)$  is  $305.11 \pm 1.782 \sqrt{402.98} = 305.11 \pm 35.773$ .

- 11.37** From Ex. 11.8 and 11.18, when  $x^* = 12$ ,  $\hat{y} = 7.15$  so that  $V(\hat{Y})$  is estimated to be  $10.97 \left[ .1 + \frac{(12 - 15.504)^2}{2359.929} \right] = 1.154$ . Thus, a 95% CI for  $E(Y)$  is  $7.15 \pm 2.306 \sqrt{1.154} = 7.15 \pm 2.477$  or  $(4.67, 9.63)$ .

- 11.38** Refer to Ex. 11.3 and 11.23, where  $s^2 = .1333$ ,  $\hat{y} = 1.5 - .6x$ ,  $S_{xx} = 10$  and  $\bar{x} = 0$ .

- When  $x^* = 0$ , the 90% CI for  $E(Y)$  is  $1.5 \pm 2.353 \sqrt{.1333(\frac{1}{5})}$  or (1.12, 1.88).
- When  $x^* = -2$ , the 90% CI for  $E(Y)$  is  $2.7 \pm 2.353 \sqrt{.1333(\frac{1}{5} + \frac{4}{10})}$  or (2.03, 3.37).
- When  $x^* = 2$ , the 90% CI for  $E(Y)$  is  $.3 \pm 2.353 \sqrt{.1333(\frac{1}{5} + \frac{4}{10})}$  or (-.37, .97).



On the graph, note the interval lengths.

- 11.39** Refer to Ex. 11.16. When  $x^* = 65$ ,  $\hat{y} = 25.395$  and a 95% CI for  $E(Y)$  is

$$25.395 \pm 2.228 \sqrt{19.033 \left[ \frac{1}{12} + \frac{(65-60)^2}{6000} \right]} \text{ or } 25.395 \pm 2.875.$$

- 11.40** Refer to Ex. 11.14. When  $x^* = .3$ ,  $\hat{y} = .7878$  and with  $SSE = .0155$ ,  $S_{xx} = .20625$ , and  $\bar{x} = .325$ , the 90% CI for  $E(Y)$  is  $.7878 \pm 1.86 \sqrt{\frac{.0155}{8} \left[ \frac{1}{10} + \frac{(.3-.325)^2}{.20625} \right]}$  or (.76, .81).

- 11.41** a. Using  $\hat{\beta}_0 = \bar{Y} - \hat{\beta}_1 \bar{x}$  and  $\hat{\beta}_1$  as estimators, we have  $\hat{\mu}_y = \bar{Y} - \hat{\beta}_1 \bar{x} + \hat{\beta}_1 \mu_x$  so that

$$\hat{\mu}_y = \bar{Y} - \hat{\beta}_1 (\bar{x} - \mu_x).$$

- b. Calculate  $V(\hat{\mu}_y) = V(\bar{Y}) + (\bar{x} - \mu_x)^2 V(\hat{\beta}_1) = \frac{\sigma^2}{n} + (\bar{x} - \mu_x)^2 \frac{\sigma^2}{S_{xx}} = \sigma^2 \left( \frac{1}{n} + \frac{(\bar{x} - \mu_x)^2}{S_{xx}} \right)$ .

From Ex. 11.4,  $s^2 = 7.1057$  and  $S_{xx} = 54,714$  so that  $\hat{\mu}_y = 72.1 + .99(74 - 72) = 74.08$

and the variance of this estimate is calculated to be  $7.1057 \left[ \frac{1}{10} + \frac{(74-72)^2}{54,714} \right] = .711$ . The two-standard deviation error bound is  $2\sqrt{.711} = 1.69$ .

- 11.42** Similar to Ex. 11.35, the variance is minimized when  $x^* = \bar{x}$ .

- 11.43** Refer to Ex. 11.5 and 11.17. When  $x = 9$  (year 1980),  $\hat{y} = 65.15$  and the 95% PI is

$$65.15 \pm 2.447 \sqrt{3.05 \left( 1 + \frac{1}{8} + \frac{(9-4.5)^2}{42} \right)} = 65.15 \pm 5.42 \text{ or } (59.73, 70.57).$$

- 11.44** For the year 1981,  $x = 10$ . So,  $\hat{y} = 69.99$  and the 95% PI is

$$69.99 \pm 2.447 \sqrt{3.05 \left( 1 + \frac{1}{8} + \frac{(10-4.5)^2}{42} \right)} = 69.99 \pm 5.80.$$

For the year 1982,  $x = 11$ . So,  $\hat{y} = 74.83$  and the 95% PI is

$$74.83 \pm 2.447 \sqrt{3.05 \left( 1 + \frac{1}{8} + \frac{(11-4.5)^2}{42} \right)} = 74.83 \pm 6.24.$$

Notice how the intervals get wider the further the prediction is from the mean. For the year 1988, this is far beyond the limits of experimentation. So, the linear relationship may not hold (note that the intervals for 1980, 1981 and 1982 are also outside of the limits, so caveat emptor).

- 11.45** From Ex. 11.8 and 11.18 (also see 11.37), when  $x^* = 12$ ,  $\hat{y} = 7.15$  so that the 95% PI is

$$7.15 \pm 2.306 \sqrt{10.97 \left[ 1 + \frac{1}{10} + \frac{(12-15.504)^2}{2359.929} \right]} = 7.15 \pm 8.03 \text{ or } (-.86, 15.18).$$

- 11.46** From 11.16 and 11.39, when  $x^* = 65$ ,  $\hat{y} = 25.395$  so that the 95% PI is given by

$$25.395 \pm 2.228 \sqrt{19.033 \left[ 1 + \frac{1}{12} + \frac{(65-60)^2}{6000} \right]} = 25.395 \pm 10.136.$$

- 11.47** From Ex. 11.14, when  $x^* = .6$ ,  $\hat{y} = .3933$  so that the 95% PI is given by

$$.3933 \pm 2.306 \sqrt{.00194 \left[ 1 + \frac{1}{10} + \frac{(.6-.325)^2}{.20625} \right]} = .3933 \pm .12 \text{ or } (.27, .51).$$

- 11.48** The summary statistics are  $S_{xx} = 380.5$ ,  $S_{xy} = 2556.0$ , and  $S_{yy} = 19,263.6$ . Thus,  $r = .944$ . To test  $H_0: \rho = 0$  vs.  $H_a: \rho > 0$ ,  $t = 8.0923$  with 8 degrees of freedom. From Table 7, we find that  $p\text{-value} < .005$ .

- 11.49** a.  $r^2$  behaves inversely to SSE, since  $r^2 = 1 - \text{SSE}/S_{yy}$ .  
b. The best model has  $r^2 = .817$ , so  $r = .90388$  (since the slope is positive,  $r$  is as well).

- 11.50** a.  $r^2$  increases as the fit improves.  
b. For the best model,  $r^2 = .982$  and so  $r = .99096$ .  
c. The scatterplot in this example exhibits a smaller error variance about the line.

- 11.51** The summary statistics are  $S_{xx} = 2359.929$ ,  $S_{xy} = 1546.459$ , and  $S_{yy} = 1101.1686$ . Thus,  $r = .9593$ . To test  $H_0: \rho = 0$  vs.  $H_a: \rho \neq 0$ ,  $|t| = 9.608$  with 8 degrees of freedom. From Table 7, we see that  $p\text{-value} < 2(.005) = .01$  so we can reject the null hypothesis that the correlation is 0.

- 11.52** a. Since the slope of the line is negative,  $r = -\sqrt{r^2} = -\sqrt{.61} = -.781$ .

b. This is given by  $r^2$ , so 61%.

c. To test  $H_0: \rho = 0$  vs.  $H_a: \rho < 0$ ,  $t = \frac{-.781\sqrt{12}}{\sqrt{1-(-.781)^2}} = -4.33$  with 12 degrees of freedom.

Since  $-t_{.05} = -1.782$ , we can reject  $H_0$  and conclude that plant density decreases with increasing altitude.

**11.53** a. This is given by  $r^2 = (.8261)^2 = .68244$ , or 68.244%.

b. Same answer as part a.

c. To test  $H_0: \rho = 0$  vs.  $H_a: \rho > 0$ ,  $t = \frac{.8261\sqrt{8}}{\sqrt{1-(.8261)^2}} = 4.146$  with 8 degrees of freedom. Since  $t_{.01} = 2.896$ , we can reject  $H_0$  and conclude that heights and weights are positively correlated for the football players.

d.  $p\text{-value} = P(T > 4.146) = .00161$ .

**11.54** a. The MOM estimators for  $\sigma_x^2$  and  $\sigma_y^2$  were given in Ex. 9.72.

b. By substituting the MOM estimators, the MOM estimator for  $\rho$  is identical to  $r$ , the MLE.

**11.55** Since  $\hat{\beta}_1 = S_{xy} / S_{xx}$  and  $r = \hat{\beta}_1 \sqrt{S_{xx} / S_{yy}}$ , we have that the usual  $t$ -test statistic is:

$$T = \frac{\hat{\beta}_1}{\sqrt{S / S_{xx}}} = \frac{\sqrt{S_{xx}} \hat{\beta}_1 \sqrt{n-2}}{\sqrt{S_{yy} - \hat{\beta}_1 S_{xy}}} = \frac{\sqrt{S_{xx} / S_{yy}} \hat{\beta}_1 \sqrt{n-2}}{\sqrt{1 - \hat{\beta}_1 S_{xy} / S_{yy}}} = \frac{r \sqrt{n-2}}{\sqrt{1-r^2}}.$$

**11.56** Here,  $r = .8$ .

a. For  $n = 5$ ,  $t = 2.309$  with 3 degrees of freedom. Since  $t_{.05} = 2.353$ , fail to reject  $H_0$ .

b. For  $n = 12$ ,  $t = 4.2164$  with 10 degrees of freedom. Here,  $t_{.05} = 1.812$ , reject  $H_0$ .

c. For part a,  $p\text{-value} = P(T > 2.309) = .05209$ . For part (b),  $p\text{-value} = .00089$ .

d. Different conclusions: note the  $\sqrt{n-2}$  term in the numerator of the test statistic.

e. The larger sample size in part b caused the computed test statistic to be more extreme. Also, the degrees of freedom were larger.

**11.57** a. The sample correlation  $r$  determines the sign.

b. Both  $r$  and  $n$  determine the magnitude of  $|t|$ .

**11.58** For the test  $H_0: \rho = 0$  vs.  $H_a: \rho > 0$ , we reject if  $t = \frac{r\sqrt{2}}{\sqrt{1-r^2}} \geq t_{.05} = 2.92$ . The smallest value of  $r$  that would lead to a rejection of  $H_0$  is the solution to the equation

$$r = \frac{2.92}{\sqrt{2}} \sqrt{1-r^2}.$$

Numerically, this is found to be  $r = .9000$ .

**11.59** For the test  $H_0: \rho = 0$  vs.  $H_a: \rho < 0$ , we reject if  $t = \frac{r\sqrt{18}}{\sqrt{1-r^2}} \leq -t_{.05} = -1.734$ . The largest value of  $r$  that would lead to a rejection of  $H_0$  is the solution to the equation

$$r = \frac{-1.734}{\sqrt{18}} \sqrt{1-r^2}.$$

Numerically, this is found to be  $r = -.3783$ .

- 11.60** Recall the approximate normal distribution of  $\frac{1}{2} \ln\left(\frac{1+r}{1-r}\right)$  given on page 606. Therefore, for sample correlations  $r_1$  and  $r_2$ , each being calculated from independent samples of size  $n_1$  and  $n_2$  (respectively) and drawn from bivariate normal populations with correlations coefficients  $\rho_1$  and  $\rho_2$  (respectively), we have that

$$Z = \frac{\frac{1}{2} \ln\left(\frac{1+r_1}{1-r_1}\right) - \frac{1}{2} \ln\left(\frac{1+r_2}{1-r_2}\right) - \left[\frac{1}{2} \ln\left(\frac{1+\rho_1}{1-\rho_1}\right) - \frac{1}{2} \ln\left(\frac{1+\rho_2}{1-\rho_2}\right)\right]}{\frac{1}{\sqrt{n_1-3}} + \frac{1}{\sqrt{n_2-3}}}$$

is approximately standard normal for large  $n_1$  and  $n_2$ .

Thus, to test  $H_0: \rho_1 = \rho_2$  vs.  $H_a: \rho_1 \neq \rho_2$  with  $r_1 = .9593$ ,  $n_1 = 10$ ,  $r_2 = .85$ ,  $n_2 = 20$ , the computed test statistic is

$$z = \frac{\frac{1}{2} \ln\left(\frac{1.9593}{.0407}\right) - \frac{1}{2} \ln\left(\frac{1.85}{.15}\right)}{\frac{1}{\sqrt{7}} + \frac{1}{\sqrt{17}}} = 1.52.$$

Since the rejection region is all values  $|z| > 1.96$  for  $\alpha = .05$ , we fail to reject  $H_0$ .

- 11.61** Refer to Example 11.10 and the results given there. The 90% PI is

$$.979 \pm 2.132(.045) \sqrt{1 + \frac{1}{6} + \frac{(1.5 - 1.457)^2}{.234}} = .979 \pm .104 \text{ or } (.875, 1.083).$$

- 11.62** Using the calculations from Example 11.11, we have  $r = \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}} = .9904$ . The proportion of variation described is  $r^2 = (.9904)^2 = .9809$ .

- 11.63** a. Observe that  $\ln E(Y) = \ln \alpha_0 - \alpha_1 x$ . Thus, the logarithm of the expected value of  $Y$  is linearly related to  $x$ . So, we can use the linear model

$$w_i = \beta_0 + \beta_1 x_i + \varepsilon_i,$$

where  $w_i = \ln y_i$ ,  $\beta_0 = \ln \alpha_0$  and  $\beta_1 = -\alpha_1$ . In the above, note that we are assuming an additive error term that is in effect after the transformation. Using the method of least squares, the summary statistics are:

$$\bar{x} = 5.5, \Sigma x^2 = 385, \bar{w} = 3.5505, S_{xw} = -.7825, S_{xx} = 82.5, \text{ and } S_{ww} = .008448.$$

Thus,  $\hat{\beta}_1 = -.0095$ ,  $\hat{\beta}_0 = 3.603$  and  $\hat{\alpha}_1 = -(-.0095) = .0095$ ,  $\hat{\alpha}_0 = \exp(3.603) = 36.70$ .

Therefore, the prediction equation is  $\hat{y} = 36.70e^{-.0095x}$ .

- b. To find a CI for  $\alpha_0$ , we first must find a CI for  $\beta_0$  and then merely transform the

endpoints of the interval. First, we calculate the SSE using  $SSE = S_{ww} - \hat{\beta}_1 S_{xw} =$

$.008448 - (-.0095)(-.782481) = .0010265$  and so  $s^2 = (.0010265)/8 = .0001283$ . Using the methods given in Section 11.5, the 90% CI for  $\beta_0$  is

$3.6027 \pm 1.86 \sqrt{.0001283 \left( \frac{385}{10(82.5)} \right)}$  or  $(3.5883, 3.6171)$ . So the 90% CI for  $\alpha_0$  is given by  $(e^{3.5883}, e^{3.6171}) = (36.17, 37.23)$ .

- 11.64** This is similar to Ex. 11.63. Note that  $\ln E(Y) = -\alpha_0 x^{\alpha_1}$  and  $\ln[-\ln E(Y)] = \ln \alpha_0 + \alpha_1 \ln x$ . So, we would expect that  $\ln(-\ln y)$  to be linear in  $\ln x$ . Define  $w_i = \ln(-\ln y_i)$ ,  $t_i = \ln x_i$ ,  $\beta_0 = \ln \alpha_0$ ,  $\beta_1 = \alpha_1$ . So, we now have the familiar linear model

$$w_i = \beta_0 + \beta_1 t_i + \varepsilon_i$$

(again, we are assuming an additive error term that is in effect after the transformation). The methods of least squares can be used to estimate the parameters. The summary statistics are

$$\bar{t} = -1.12805, \bar{w} = -1.4616, S_{tw} = 3.6828, \text{ and } S_{tt} = 1.51548$$

So,  $\hat{\beta}_1 = 2.4142$ ,  $\hat{\beta}_0 = 1.2617$  and thus  $\hat{\alpha}_1 = 2.4142$  and  $\hat{\alpha}_0 = \exp(1.2617) = 3.5315$ .

This fitted model is  $\hat{y} = \exp(-3.5315x^{2.4142})$ .

- 11.65** If  $y$  is related to  $t$  according to  $y = 1 - e^{-\beta t}$ , then  $-\ln(1 - y) = \beta t$ . Thus, let  $w_i = -\ln(1 - y_i)$  and we have the linear model

$$w_i = \beta t_i + \varepsilon_i$$

(again assuming an additive error term). This is the “no-intercept” model described in

Ex. 11.10 and the least squares estimator for  $\beta$  is given to be  $\hat{\beta} = \frac{\sum_{i=1}^n t_i w_i}{\sum_{i=1}^n t_i^2}$ . Now, using

similar methods from Section 11.4, note that  $V(\hat{\beta}) = \frac{\sigma^2}{\sum_{i=1}^n t_i^2}$  and  $\frac{SSE}{\sigma^2} = \frac{\sum_{i=1}^n (w_i - \hat{w})^2}{\sigma^2}$

is chi-square with  $n - 1$  degrees of freedom. So, by Definition 7.2, the quantity

$$T = \frac{\hat{\beta} - \beta}{S / \sum_{i=1}^n t_i^2},$$

where  $S = SSE/(n - 1)$ , has a  $t$ -distribution with  $n - 1$  degrees of freedom.

A  $100(1 - \alpha)\%$  CI for  $\beta$  is

$$\hat{\beta} \pm t_{\alpha/2} S \sqrt{\frac{1}{\sum_{i=1}^n t_i^2}},$$

and  $t_{\alpha/2}$  is the upper- $\alpha/2$  critical value from the  $t$ -distribution with  $n - 1$  degrees of freedom.

- 11.66** Using the matrix notation from this section,

$$\mathbf{X} = \begin{bmatrix} 1 & -2 \\ 1 & -1 \\ 1 & 0 \\ 1 & 1 \\ 1 & 2 \end{bmatrix} \quad \mathbf{Y} = \begin{bmatrix} 3 \\ 2 \\ 1 \\ 1 \\ .5 \end{bmatrix} \quad \mathbf{X}'\mathbf{Y} = \begin{bmatrix} 7.5 \\ -6 \end{bmatrix} \quad \mathbf{X}'\mathbf{X} = \begin{bmatrix} 5 & 0 \\ 0 & 10 \end{bmatrix}.$$

Thus,  $\hat{\beta} = \begin{bmatrix} 5 & 0 \\ 0 & 10 \end{bmatrix}^{-1} \begin{bmatrix} 7.5 \\ -6 \end{bmatrix} = \begin{bmatrix} .2 & 0 \\ 0 & .1 \end{bmatrix} \begin{bmatrix} 7.5 \\ -6 \end{bmatrix} = \begin{bmatrix} 1.5 \\ -.6 \end{bmatrix}$  so that  $\hat{y} = 1.5 - .6x$ .

$$11.67 \quad X = \begin{bmatrix} 1 & -1 \\ 1 & 0 \\ 1 & 1 \\ 1 & 2 \\ 1 & 3 \end{bmatrix} \quad Y = \begin{bmatrix} 3 \\ 2 \\ 1 \\ 1 \\ .5 \end{bmatrix} \quad X'Y = \begin{bmatrix} 7.5 \\ 1.5 \end{bmatrix} \quad X'X = \begin{bmatrix} 5 & 5 \\ 5 & 15 \end{bmatrix}$$

The student should verify that  $(X'X)^{-1} = \begin{bmatrix} .3 & -.1 \\ -.1 & .1 \end{bmatrix}$  so that  $\hat{\beta} = \begin{bmatrix} 2.1 \\ -.6 \end{bmatrix}$ . Note that the slope is the same as in Ex. 11.66, but the y-intercept is different. Since  $X'X$  is not a diagonal matrix (as in Ex. 11.66), computing the inverse is a bit more tedious.


$$11.68 \quad X = \begin{bmatrix} 1 & -3 & 9 \\ 1 & -2 & 4 \\ 1 & -1 & 1 \\ 1 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \end{bmatrix} \quad Y = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -1 \\ -1 \\ 0 \\ 0 \end{bmatrix} \quad X'Y = \begin{bmatrix} -1 \\ 4 \\ 8 \end{bmatrix} \quad X'X = \begin{bmatrix} 7 & 0 & 28 \\ 0 & 28 & 0 \\ 28 & 0 & 196 \end{bmatrix}.$$

The student should verify (either using Appendix I or a computer),

$$(X'X)^{-1} = \begin{bmatrix} .3333 & 0 & -.04762 \\ 0 & .035714 & 0 \\ -.04762 & 0 & .011905 \end{bmatrix} \text{ so that } \hat{\beta} = \begin{bmatrix} -.714285 \\ -.142857 \\ .142859 \end{bmatrix} \text{ and the fitted}$$

model is  $\hat{y} = -.714285 - .142857x + .142859x^2$ .





The graphed curve is above.

**11.69** For this problem, R will be used.

```
> x <- c(-7, -5, -3, -1, 1, 3, 5, 7)
> y <- c(18.5, 22.6, 27.2, 31.2, 33.0, 44.9, 49.4, 35.0)
```

**a. Linear model:**

```
> lm(y~x)
```

Call:

```
lm(formula = y ~ x)
```

Coefficients:

```
(Intercept)          x
    32.725         1.812
```

$$\hat{y} = 32.725 + 1.812x$$

**b. Quadratic model**

```
> lm(y~x+I(x^2))
```

Call:

```
lm(formula = y ~ x + I(x^2))
```

Coefficients:

```
(Intercept)          x      I(x^2)
    35.5625         1.8119     -0.1351
```

$$\hat{y} = 35.5625 + 1.8119x - 0.1351x^2$$

**11.70 a.** The student should verify that  $\mathbf{Y}'\mathbf{Y} = 105,817$ ,  $\mathbf{X}'\mathbf{Y} = \begin{bmatrix} 721 \\ 106155 \end{bmatrix}$ , and  $\hat{\boldsymbol{\beta}} = \begin{bmatrix} .719805 \\ .991392 \end{bmatrix}$ .

So,  $\text{SSE} = 105,817 - 105,760.155 = 56.845$  and  $s^2 = 56.845/8 = 7.105625$ .

**b.** Using the coding as specified, the data are:

$x_i^*$	-62	-60	-63	-45	-25	40	-36	169	-13	95
$y_i$	9	14	7	29	45	109	40	238	60	70

The student should verify that  $\mathbf{X}^{*'}\mathbf{Y} = \begin{bmatrix} 721 \\ 54243 \end{bmatrix}$ ,  $\mathbf{X}^{*'}\mathbf{X}^* = \begin{bmatrix} 10 & 0 \\ 0 & 54,714 \end{bmatrix}$  and

$\hat{\boldsymbol{\beta}} = \begin{bmatrix} .721 \\ .991392 \end{bmatrix}$ . So,  $\text{SSE} = 105,817 - 105,760.155 = 56.845$  (same answer as part a).

**11.71** Note that the vector  $\mathbf{a}$  is composed of  $k$  0's and one 1. Thus,

$$E(\hat{\beta}_i) = E(\mathbf{a}'\hat{\boldsymbol{\beta}}) = \mathbf{a}'E(\hat{\boldsymbol{\beta}}) = \mathbf{a}'\boldsymbol{\beta} = \beta_i$$

$$V(\hat{\beta}_i) = V(\mathbf{a}'\hat{\boldsymbol{\beta}}) = \mathbf{a}'E(\hat{\boldsymbol{\beta}})\mathbf{a} = \mathbf{a}'\sigma^2(\mathbf{X}'\mathbf{X})^{-1}\mathbf{a} = \sigma^2\mathbf{a}'(\mathbf{X}'\mathbf{X})^{-1}\mathbf{a} = c_{ii}\sigma^2$$

**11.72** Following Ex. 11.69, more detail with the R output is given by:

```
> summary(lm(y~x+I(x^2)))
```

Call:

```
lm(formula = y ~ x + I(x^2))
```

Residuals:

```
      1      2      3      4      5      6      7      8
2.242 -0.525 -1.711 -2.415 -4.239  5.118  8.156 -6.625
```

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )	
(Intercept)	35.5625	3.1224	11.390	9.13e-05	***
x	1.8119	0.4481	4.044	0.00988	**
I(x^2)	-0.1351	0.1120	-1.206	0.28167	

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 5.808 on 5 degrees of freedom

Multiple R-Squared: 0.7808, Adjusted R-squared: 0.6931

F-statistic: 8.904 on 2 and 5 DF, p-value: 0.0225

a. To test  $H_0: \beta_2 = 0$  vs.  $H_a: \beta_2 \neq 0$ , the computed test statistic is  $t = -1.206$  and  $p$ -value = .28167. Thus,  $H_0$  would not be rejected (no quadratic effect).

b. From the output, it is presented that  $\sqrt{V(\hat{\beta}_2)} = .1120$ . So, with 5 degrees of freedom,  $t_{.05} = 3.365$  so a 90% for  $\beta_2$  is  $-.1351 \pm (3.365)(.1120) = -.1351 \pm .3769$  or  $(-.512, .2418)$ . Note that this interval contains 0, agreeing with part a.

**11.73** If the minimum value is to occur at  $x_0 = 1$ , then this implies  $\beta_1 + 2\beta_2 = 0$ . To test this claim, let  $\mathbf{a}' = [0 \ 1 \ 2]$  for the hypothesis  $H_0: \beta_1 + 2\beta_2 = 0$  vs.  $H_a: \beta_1 + 2\beta_2 \neq 0$ . From Ex. 11.68, we have that  $\hat{\beta}_1 + 2\hat{\beta}_2 = .142861$ ,  $s^2 = .14285$  and we calculate  $\mathbf{a}'(\mathbf{X}'\mathbf{X})^{-1}\mathbf{a} = .083334$ . So, the computed value of the test statistic is  $|t| = \frac{.142861}{\sqrt{.14285(.083334)}} = 1.31$  with 4 degrees of freedom. Since  $t_{.025} = 2.776$ ,  $H_0$  is not rejected.

**11.74** a. Each transformation is defined, for each factor, by subtracting the midpoint (the mean) and dividing by one-half the range.

b. Using the matrix definitions of  $\mathbf{X}$  and  $\mathbf{Y}$ , we have that

$$\mathbf{X}'\mathbf{Y} = \begin{bmatrix} 338 \\ -50.2 \\ -19.4 \\ -2.6 \\ -20.4 \end{bmatrix} \quad \mathbf{X}'\mathbf{X} = \begin{bmatrix} 16 & 0 & 0 & 0 & 0 \\ 0 & 16 & 0 & 0 & 0 \\ 0 & 0 & 16 & 0 & 0 \\ 0 & 0 & 0 & 16 & 0 \\ 0 & 0 & 0 & 0 & 16 \end{bmatrix} \quad \text{so that } \hat{\boldsymbol{\beta}} = \begin{bmatrix} 21.125 \\ -3.1375 \\ -1.2125 \\ -.1625 \\ -1.275 \end{bmatrix}.$$

The fitted model is  $\hat{y} = 21.125 - 3.1375x_1 - 1.2125x_2 - .1625x_3 - 1.275x_4$ .

c. First, note that  $\text{SSE} = \mathbf{Y}'\mathbf{Y} - \hat{\boldsymbol{\beta}}'\mathbf{X}'\mathbf{Y} = 7446.52 - 7347.7075 = 98.8125$  so that  $s^2 = 98.8125/(16 - 5) = 8.98$ . Further, tests of  $H_0: \beta_i = 0$  vs.  $H_0: \beta_i \neq 0$  for  $i = 1, 2, 3, 4$ , are based on the statistic  $t_i = \frac{\hat{\beta}_i}{s\sqrt{c_{ii}}} = \frac{4\hat{\beta}_i}{\sqrt{8.98}}$  and  $H_0$  is rejected if  $|t_i| > t_{.005} = 3.106$ . The

four computed test statistics are  $t_1 = -4.19$ ,  $t_2 = -1.62$ ,  $t_3 = -.22$  and  $t_4 = -1.70$ . Thus, only the first hypothesis involving the first temperature factor is significant.

- 11.75** With the four given factor levels, we have  $\mathbf{a}' = [1 \ -1 \ 1 \ -1 \ 1]$  and so  $\mathbf{a}'(\mathbf{X}'\mathbf{X})^{-1}\mathbf{a} = 5/16$ . The estimate of the mean of  $Y$  at this setting is
- $$\hat{y} = 21.125 + 3.1375 - 1.2125 + .1625 - 1.275 = 21.9375$$
- and the 90% confidence interval (based on 11 degrees of freedom) is
- $$21.9375 \pm 1.796\sqrt{8.96}\sqrt{5/16} = 21.9375 \pm 3.01 \text{ or } (18.93, 24.95).$$

- 11.76** First, we calculate  $s^2 = \text{SSE}/(n - k - 1) = 1107.01/11 = 100.637$ .

a. To test  $H_0: \beta_2 = 0$  vs.  $H_0: \beta_2 < 0$ , we use the  $t$ -test with  $c_{22} = 8.1 \cdot 10^{-4}$ :

$$t = \frac{-.92}{\sqrt{100.637(.00081)}} = -3.222.$$

With 11 degrees of freedom,  $-t_{.05} = -1.796$  so we reject  $H_0$ : there is sufficient evidence that  $\beta_2 < 0$ .

- b. (Similar to Ex. 11.75) With the three given levels, we have  $\mathbf{a}' = [1 \ 914 \ 65 \ 6]$  and so  $\mathbf{a}'(\mathbf{X}'\mathbf{X})^{-1}\mathbf{a} = 92.76617$ . The estimate of the mean of  $Y$  at this setting is
- $$\hat{y} = 38.83 - .0092(914) - .92(65) + 11.56(6) = 39.9812$$
- and the 95% CI based on 11 degrees of freedom is
- $$39.9812 \pm 2.201\sqrt{100.637}\sqrt{92.76617} = 39.9812 \pm 212.664.$$

- 11.77** Following Ex. 11.76, the 95% PI is  $39.9812 \pm 2.201\sqrt{100.637}\sqrt{93.76617} = 39.9812 \pm 213.807$ .

- 11.78** From Ex. 11.69, the fitted model is  $\hat{y} = 35.5625 + 1.8119x - .1351x^2$ . For the year 2004,  $x = 9$  and the predicted sales is  $\hat{y} = 35.5625 + 1.8119(9) - .135(9^2) = 40.9346$ . With we have  $\mathbf{a}' = [1 \ 9 \ 81]$  and so  $\mathbf{a}'(\mathbf{X}'\mathbf{X})^{-1}\mathbf{a} = 1.94643$ . The 98% PI for *Lexus* sales in 2004 is then

$$40.9346 \pm 3.365(5.808)\sqrt{1 + 1.94643} = 40.9346 \pm 33.5475.$$

- 11.79** For the given levels,  $\hat{y} = 21.9375$ ,  $\mathbf{a}'(\mathbf{X}'\mathbf{X})^{-1}\mathbf{a} = .3135$ , and  $s^2 = 8.98$ . The 90% PI based on 11 degrees of freedom is  $21.9375 \pm 1.796\sqrt{8.98(1 + .3135)} = 21.9375 \pm 6.17$  or (15.77, 28.11).

- 11.80** Following Ex. 11.31,  $S_{yy} = .3748$  and  $SSE = S_{yy} - \hat{\beta}_1 S_{xy} = .3748 - (.004215)(88.8) = .000508$ . Therefore, the  $F$ -test is given by  $F = \frac{(.3748 - .000508)/1}{.000508/7} = 5157.57$  with 1 numerator and 7 denominator degrees of freedom. Clearly,  $p$ -value  $< .005$  so reject  $H_0$ .

- 11.81** From Definition 7.2, let  $Z \sim \text{Nor}(0, 1)$  and  $W \sim \chi_v^2$ , and let  $Z$  and  $W$  be independent.

Then,  $T = \frac{Z}{\sqrt{W/v}}$  has the  $t$ -distribution with  $v$  degrees of freedom. But, since  $Z^2 \sim \chi_1^2$ ,

by Definition 7.3,  $F = T^2$  has a  $F$ -distribution with 1 numerator and  $v$  denominator degrees of freedom. Now, specific to this problem, note that if  $k = 1$ ,  $SSE_R = S_{yy}$ . So, the reduced model  $F$ -test simplifies to

$$F = \frac{S_{yy} - (S_{yy} - \hat{\beta}_1 S_{xy})}{SSE_C / (n - 2)} = \frac{\hat{\beta}_1^2}{s^2 / S_{xx}} = T^2.$$

- 11.82** a. To test  $H_0: \beta_1 = \beta_2 = \beta_3 = 0$  vs.  $H_a$ : at least one  $\beta_i \neq 0$ , the  $F$ -statistic is

$$F = \frac{(10965.46 - 1107.01)/3}{1107.01/11} = 32.653,$$

with 3 numerator and 11 denominator degrees of freedom. From Table 7, we see that  $p$ -value  $< .005$ , so there is evidence that at least one predictor variable contributes.

- b. The coefficient of determination is  $R^2 = 1 - \frac{1107.01}{10965.46} = .899$ , so 89.9% of the variation in percent yield ( $Y$ ) is explained by the model.

- 11.83** a. To test  $H_0: \beta_2 = \beta_3 = 0$  vs.  $H_a$ : at least one  $\beta_i \neq 0$ , the reduced model  $F$ -test is

$$F = \frac{(5470.07 - 1107.01)/2}{1107.01/11} = 21.677,$$

with 2 numerator and 11 denominator degrees of freedom. Since  $F_{.05} = 3.98$ , we can reject  $H_0$ .

**b.** We must find the value of  $SSE_R$  such that  $\frac{(SSE_R - 1107.01)/2}{1107.01/11} = 3.98$ . The solution is  $SSE_R = 1908.08$

**11.84 a.** The result follows from

$$\frac{n - (k + 1)}{k} \left( \frac{R^2}{1 - R^2} \right) = \frac{n - (k + 1)}{k} \left( \frac{1 - SSE/S_{yy}}{SSE/S_{yy}} \right) = \frac{(S_{yy} - SSE)/k}{SSE/[n - (k + 1)]} = F.$$

**b.** The form is  $F = T^2$ .

**11.85** Here,  $n = 15$ ,  $k = 4$ .

**a.** Using the result from Ex. 11.84,  $F = \frac{10}{4} \left( \frac{.942}{1 - .942} \right) = 40.603$  with 4 numerator and 10 denominator degrees of freedom. From Table 7, it is clear that  $p\text{-value} < .005$ , so we can safely conclude that at least one of the variables contributes to predicting the selling price.

**b.** Since  $R^2 = 1 - SSE/S_{yy}$ ,  $SSE = 16382.2(1 - .942) = 950.1676$ .

**11.86** To test  $H_0: \beta_2 = \beta_3 = \beta_4 = 0$  vs.  $H_a$ : at least one  $\beta_i \neq 0$ , the reduced-model  $F$ -test is

$$F = \frac{(1553 - 950.16)/3}{950.1676/10} = 2.115,$$

with 3 numerator and 10 denominator degrees of freedom. Since  $F_{.05} = 3.71$ , we fail to reject  $H_0$  and conclude that these variables should be dropped from the model.

**11.87 a.** The  $F$ -statistic, using the result in Ex. 11.84, is  $F = \frac{2}{4} \left( \frac{.9}{.1} \right) = 4.5$  with 4 numerator and 2 denominator degrees of freedom. Since  $F_{.1} = 9.24$ , we fail to reject  $H_0$ .

**b.** Since  $k$  is large with respect to  $n$ , this makes the computed  $F$ -statistic small.

**c.** The  $F$ -statistic, using the result in Ex. 11.84, is  $F = \frac{40}{3} \left( \frac{.15}{.85} \right) = 2.353$  with 3 numerator and 40 denominator degrees of freedom. Since  $F_{.1} = 2.23$ , we can reject  $H_0$ .

**d.** Since  $k$  is small with respect to  $n$ , this makes the computed  $F$ -statistic large.

**11.88 a.** False; there are 15 degrees of freedom for SSE.

**b.** False; the fit ( $R^2$ ) cannot improve when independent variables are removed.

**c.** True

**d.** False; not necessarily, since the degrees of freedom associated with each SSE is different.

**e.** True.

**f.** False; Model III is not a reduction of Model I (note the  $x_1x_2$  term).

- 11.89** a. True.  
 b. False; not necessarily, since Model III is not a reduction of Model I (note the  $x_1x_2$  term).  
 c. False; for the same reason in part (b).
- 11.90** Refer to Ex. 11.69 and 11.72.  
 a. We have that  $SSE_R = 217.7112$  and  $SSE_C = 168.636$ . For  $H_0: \beta_2 = 0$  vs.  $H_a: \beta_2 \neq 0$ , the reduced model F-test is  $F = \frac{217.7112 - 168.636}{168.636/5} = 1.455$  with 1 numerator and 5 denominator degrees of freedom. With  $F_{.05} = 6.61$ , we fail to reject  $H_0$ .  
 b. Referring to the R output given in Ex. 11.72, the  $F$ -statistic is  $F = 8.904$  and the  $p$ -value for the test is .0225. This leads to a rejection at the  $\alpha = .05$  level.

- 11.91** The hypothesis of interest is  $H_0: \beta_1 = \beta_4 = 0$  vs.  $H_a$ : at least one  $\beta_i \neq 0, i = 1, 4$ . From Ex. 11.74, we have  $SSE_C = 98.8125$ . To find  $SSE_R$ , we fit the linear regression model with just  $x_2$  and  $x_3$  so that

$$\mathbf{X}'\mathbf{Y} = \begin{bmatrix} 338 \\ -19.4 \\ -2.6 \end{bmatrix} \quad (\mathbf{X}'\mathbf{X})^{-1} = \begin{bmatrix} 1/16 & 0 & 0 \\ 0 & 1/16 & 0 \\ 0 & 0 & 1/16 \end{bmatrix}$$

and so  $SSE_R = 7446.52 - 7164.195 = 282.325$ . The reduced-model  $F$ -test is

$$F = \frac{(282.325 - 98.8125)/2}{98.8125/11} = 10.21,$$

with 2 numerator and 11 denominator degrees of freedom. Thus, since  $F_{.05} = 3.98$ , we can reject  $H_0$  and conclude that either  $T_1$  or  $T_2$  (or both) affect the yield.

- 11.92** To test  $H_0: \beta_3 = \beta_4 = \beta_5 = 0$  vs.  $H_a$ : at least one  $\beta_i \neq 0$ , the reduced-model  $F$ -test is

$$F = \frac{(465.134 - 152.177)/3}{152.177/18} = 12.34,$$

with 3 numerator and 18 denominator degrees of freedom. Since  $F_{.005} = 5.92$ , we have that  $p$ -value  $< .005$ .

- 11.93** Refer to Example. 11.19. For the reduced model,  $s^2 = 326.623/8 = 40.83$ . Then,

$$(\mathbf{X}'\mathbf{X})^{-1} = \begin{bmatrix} 1/11 & 0 & 0 \\ 0 & 2/17 & 0 \\ 0 & 0 & 2/17 \end{bmatrix}, \mathbf{a}' = [1 \quad 1 \quad -1].$$

So,  $\hat{y} = \mathbf{a}'\hat{\boldsymbol{\beta}} = 93.73 + 4 - 7.35 = 90.38$  and  $\mathbf{a}'(\mathbf{X}'\mathbf{X})^{-1}\mathbf{a} = .3262$ . The 95% CI for  $E(Y)$  is  $90.38 \pm 2.306\sqrt{40.83(.3262)} = 90.38 \pm 8.42$  or (81.96, 98.80).

**11.94** From Example 11.19, tests of  $H_0: \beta_i = 0$  vs.  $H_0: \beta_i \neq 0$  for  $i = 3, 4, 5$ , are based on the

statistic  $t_i = \frac{\hat{\beta}_i}{s\sqrt{c_{ii}}}$  with 5 degrees of freedom and  $H_0$  is rejected if  $|t_i| > t_{.01} = 4.032$ .

The three computed test statistics are  $|t_3| = .58$ ,  $|t_4| = 3.05$ ,  $|t_5| = 2.53$ . Therefore, none of the three parameters are significantly different from 0.

**11.95** a. The summary statistics are:  $\bar{x} = -268.28$ ,  $\bar{y} = .6826$ ,  $S_{xy} = -15.728$ ,  $S_{xx} = 297.716$ , and  $S_{yy} = .9732$ . Thus,  $\hat{y} = -13.54 - 0.053x$ .

b. First,  $SSE = .9732 - (-.053)(-15.728) = .14225$ , so  $s^2 = .14225/8 = .01778$ . The test statistic is  $t = \frac{-.053}{\sqrt{\frac{.01778}{297.716}}} = -6.86$  and  $H_0$  is rejected at the  $\alpha = .01$  level.

c. With  $x = -273$ ,  $\hat{y} = -13.54 - .053(-273) = .929$ . The 95% PI is  $.929 \pm 2.306\sqrt{.01778}\sqrt{1 + \frac{1}{10} + \frac{(273-268.28)^2}{297.716}} = .929 \pm .33$ .

**11.96** Here, R will be used to fit the model:

```
> x <- c(.499, .558, .604, .441, .550, .528, .418, .480, .406, .467)
> y <- c(11.14, 12.74, 13.13, 11.51, 12.38, 12.60, 11.13, 11.70, 11.02, 11.41)
> summary(lm(y~x))
```

Call:

```
lm(formula = y ~ x)
```

Residuals:

Min	1Q	Median	3Q	Max
-0.77823	-0.07102	0.08181	0.16435	0.36771

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	6.5143	0.8528	7.639	6.08e-05 ***
x	10.8294	1.7093	6.336	0.000224 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.3321 on 8 degrees of freedom

Multiple R-Squared: 0.8338, Adjusted R-squared: 0.813

F-statistic: 40.14 on 1 and 8 DF, p-value: 0.0002241

a. The fitted model is  $\hat{y} = 6.5143 + 10.8294x$ .

b. The test  $H_0: \beta_1 = 0$  vs.  $H_a: \beta_1 \neq 0$  has a  $p$ -value of .000224, so  $H_0$  is rejected.

c. It is found that  $s = .3321$  and  $S_{xx} = .0378$ . So, with  $x = .59$ ,  $\hat{y} = 6.5143 + 10.8294(.59) = 12.902$ . The 90% CI for  $E(Y)$  is

$$12.902 \pm 1.860(.3321)\sqrt{\frac{1}{10} + \frac{(.59-.4951)^2}{.0378}} = 12.902 \pm .36.$$

**11.97** a. Using the matrix notation,

$$X = \begin{bmatrix} 1 & -3 & 5 & -1 \\ 1 & -2 & 0 & 1 \\ 1 & -1 & -3 & 1 \\ 1 & 0 & -4 & 0 \\ 1 & 1 & -3 & -1 \\ 1 & 2 & 0 & -1 \\ 1 & 3 & 5 & 1 \end{bmatrix}, Y = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}, X'Y = \begin{bmatrix} 10 \\ 14 \\ 10 \\ -3 \end{bmatrix}, (X'X)^{-1} = \begin{bmatrix} 1/7 & 0 & 0 & 0 \\ 0 & 1/28 & 0 & 0 \\ 0 & 0 & 1/84 & 0 \\ 0 & 0 & 0 & 1/6 \end{bmatrix}.$$

So, the fitted model is found to be  $\hat{y} = 1.4825 + .5x_1 + .1190x_2 - .5x_3$ .

**b.** The predicted value is  $\hat{y} = 1.4825 + .5 - .357 + .5 = 2.0715$ . The observed value at these levels was  $y = 2$ . The predicted value was based on a model fit (using all of the data) and the latter is an observed response.

**c.** First, note that  $SSE = 24 - 23.9757 = .0243$  so  $s^2 = .0243/3 = .008$ . The test statistic is  $t = \frac{\hat{\beta}_3}{s\sqrt{c_{ii}}} = \frac{-5}{\sqrt{.008(1/6)}} = -13.7$  which leads to a rejection of the null hypothesis.

**d.** Here,  $\mathbf{a}' = [1 \ 1 \ -3 \ -1]$  and so  $\mathbf{a}'(X'X)^{-1}\mathbf{a} = .45238$ . So, the 95% CI for  $E(Y)$  is  $2.0715 \pm 3.182\sqrt{.008}\sqrt{.45238} = 2.0715 \pm .19$  or (1.88, 2.26).

**e.** The prediction interval is  $2.0715 \pm 3.182\sqrt{.008}\sqrt{1 + .45238} = 2.0715 \pm .34$  or (1.73, 2.41).

**11.98** Symmetric spacing about the origin creates a diagonal  $X'X$  matrix which is very easy to invert.

**11.99** Since  $V(\hat{\beta}_1) = \frac{\sigma^2}{S_{xx}}$ , this will be minimized when  $S_{xx} = \sum_{i=1}^n (x_i - \bar{x})^2$  is as large as

possible. This occurs when the  $x_i$  are as far away from  $\bar{x}$  as possible. If  $-9 \leq x \leq 9$ , chose  $n/2$  at  $x = -9$  and  $n/2$  at  $x = 9$ .

**11.100** Based on the minimization strategy in Ex. 11.99, the values of  $x$  are:  $-9, -9, -9, -9, -9, 9, 9, 9, 9, 9$ . Thus  $S_{xx} = \sum_{i=1}^{10} (x_i - \bar{x})^2 = \sum_{i=1}^{10} x_i^2 = 810$ . If equal spacing is employed, the values of  $x$  are:  $-9, -7, -5, -3, -1, 1, 3, 5, 7, 9$ . Thus,  $S_{xx} = \sum_{i=1}^{10} (x_i - \bar{x})^2 = \sum_{i=1}^{10} x_i^2 = 330$ . The relative efficiency is the ratio of the variances, or  $330/810 = 11/27$ .

**11.101** Here, R will be used to fit the model:

```
> x1 <- c(0,0,0,0,0,1,1,1,1,1)
> x2 <- c(-2,-1,0,1,2,-2,-1,0,1,2)
> y <- c(8,9,9.1,10.2,10.4,10,10.3,12.2,12.6,13.9)
> summary(lm(y~x1+x2+I(x1*x2)))
```




```
Call:
lm(formula = y ~ x1 + x2 + I(x1 * x2))

Residuals:
    Min       1Q   Median       3Q      Max
-0.4900 -0.1925 -0.0300  0.2500  0.4000

Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)   9.3400     0.1561   59.834 1.46e-09 ***
x1             2.4600     0.2208   11.144 3.11e-05 ***
x2             0.6000     0.1104    5.436 0.00161 **
I(x1 * x2)     0.4100     0.1561    2.627 0.03924 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
Residual standard error: 0.349 on 6 degrees of freedom
Multiple R-Squared:  0.9754,    Adjusted R-squared:  0.963
F-statistic: 79.15 on 3 and 6 DF,  p-value: 3.244e-05
```

- a. The fitted model is  $\hat{y} = 9.34 + 2.46x_1 + .6x_2 + .41x_1x_2$ .
- b. For bacteria type A,  $x_1 = 0$  so  $\hat{y} = 9.34 + .6x_2$  (dotted line)  
For bacteria type B,  $x_1 = 1$  so  $\hat{y} = 11.80 + 1.01x_2$  (solid line)



- c. For bacteria A,  $x_1 = 0, x_2 = 0$ , so  $\hat{y} = 9.34$ . For bacteria B,  $x_1 = 1, x_2 = 0$ , so  $\hat{y} = 11.80$ . The observed growths were 9.1 and 12.2, respectively.
- d. The rates are different if the parameter  $\beta_3$  is nonzero. So,  $H_0: \beta_3 = 0$  vs.  $H_a: \beta_3 \neq 0$  has a  $p$ -value = .03924 (R output above) and  $H_0$  is rejected.
- e. With  $x_1 = 1, x_2 = 1$ , so  $\hat{y} = 12.81$ . With  $s = .349$  and  $\mathbf{a}'(X'X)^{-1}\mathbf{a} = .3$ , the 90% CI is  $12.81 \pm .37$ .
- f. The 90% PI is  $12.81 \pm .78$ .

**11.102** The reduced model  $F$  statistic is  $F = \frac{(795.23 - 783.9)/2}{783.9/195} = 1.41$  with 2 numerator and 195 denominator degrees of freedom. Since  $F_{.05} \approx 3.00$ , we fail to reject  $H_0$ : salary is not dependent on gender.

**11.103** Define  $\mathbf{I}$  as a column vector of  $n$  1's. Then  $\bar{y} = \frac{1}{n} \mathbf{I}' \mathbf{Y}$ . We must solve for the vector  $\mathbf{x}$  such that  $\bar{y} = \mathbf{x}' \hat{\boldsymbol{\beta}}$ . Using the matrix definition of  $\hat{\boldsymbol{\beta}}$ , we have

$$\bar{y} = \mathbf{x}' (\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}' \mathbf{Y} = \frac{1}{n} \mathbf{I}' \mathbf{Y}$$

$$\mathbf{x}' (\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}' \mathbf{Y} \mathbf{Y}' = \frac{1}{n} \mathbf{I}' \mathbf{Y} \mathbf{Y}'$$

which implies

$$\mathbf{x}' (\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}' = \frac{1}{n} \mathbf{I}'$$

$$\mathbf{x}' (\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}' \mathbf{X} = \frac{1}{n} \mathbf{I}' \mathbf{X}$$

so that

$$\mathbf{x}' = \frac{1}{n} \mathbf{I}' \mathbf{X}.$$

That is,  $\mathbf{x}' = [1 \ \bar{x}_1 \ \bar{x}_2 \ \dots \ \bar{x}_k]$ .

**11.104** Here, we will use the coding  $x_1 = \frac{P-65}{15}$  and  $x_2 = \frac{T-200}{100}$ . Then, the levels are  $x_1 = -1, 1$  and  $x_2 = -1, 0, 1$ .

$$\mathbf{a. Y} = \begin{bmatrix} 21 \\ 23 \\ 26 \\ 22 \\ 23 \\ 28 \end{bmatrix} \quad \mathbf{X} = \begin{bmatrix} 1 & -1 & -1 & 1 \\ 1 & -1 & 0 & 0 \\ 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & -1 & 1 \end{bmatrix} \quad \mathbf{X}' \mathbf{Y} = \begin{bmatrix} 143 \\ 3 \\ 11 \\ 97 \end{bmatrix} \quad (\mathbf{X}' \mathbf{X})^{-1} = \begin{bmatrix} .5 & 0 & 0 & -.5 \\ 0 & .1667 & 0 & 0 \\ 0 & 0 & .25 & 0 \\ -.5 & 0 & 0 & .75 \end{bmatrix}$$

So, the fitted model is  $\hat{y} = 23 + .5x_1 + 2.75x_2 + 1.25x_2^2$ .

**b.** The hypothesis of interest is  $H_0: \beta_3 = 0$  vs.  $H_a: \beta_3 \neq 0$  and the test statistic is (verify that  $\text{SSE} = 1$  so that  $s^2 = .5$ )  $|t| = \frac{|1.25|}{\sqrt{.5(.75)}} = 2.040$  with 2 degrees of freedom. Since  $t_{.025} = 4.303$ , we fail to reject  $H_0$ .

**c.** To test  $H_0: \beta_2 = \beta_3 = 0$  vs.  $H_a$ : at least one  $\beta_i \neq 0$ ,  $i = 2, 3$ , the reduced model must be fitted. It can be verified that  $\text{SSE}_R = 33.33$  so that the reduced model  $F$ -test is  $F = 32.33$  with 2 numerator and 2 denominator degrees of freedom. It is easily seen that  $H_0$  should be rejected; temperature does affect yield.

**11.105 a.**  $\hat{\beta}_1 = \frac{s_{xy}}{s_{xx}} = \frac{s_{xy}}{\sqrt{s_{xx}s_{yy}}} \sqrt{\frac{s_{yy}}{s_{xx}}} = r \sqrt{\frac{s_{yy}}{s_{xx}}}.$

b. The conditional distribution of  $Y_i$ , given  $X_i = x_i$ , is (see Chapter 5) normal with mean  $\mu_y + \rho \frac{\sigma_y}{\sigma_x}(x_i - \mu_x)$  and variance  $\sigma_y^2(1 - \rho^2)$ . Redefine  $\beta_1 = \rho \frac{\sigma_y}{\sigma_x}$ ,  $\beta_0 = \mu_y - \beta_1 \mu_x$ . So, if  $\rho = 0$ ,  $\beta_1 = 0$ . So, using the usual  $t$ -statistic to test  $\beta_1 = 0$ , we have

$$T = \frac{\hat{\beta}_1}{S / \sqrt{S_{xx}}} = \frac{\hat{\beta}_1}{\sqrt{\frac{SSE}{n-2}} \sqrt{\frac{1}{S_{xx}}}} = \frac{\hat{\beta}_1 \sqrt{(n-2)S_{xx}}}{\sqrt{(1-r^2)S_{yy}}}.$$

c. By part a,  $\hat{\beta}_1 = r \sqrt{\frac{S_{yy}}{S_{xx}}}$  and the statistic has the form as shown. Note that the distribution only depends on  $n - 2$  and not the particular value  $x_i$ . So, the distribution is the same unconditionally.

**11.106** The summary statistics are  $S_{xx} = 66.54$ ,  $S_{xy} = 71.12$ , and  $S_{yy} = 93.979$ . Thus,  $r = .8994$ . To test  $H_0: \rho = 0$  vs.  $H_a: \rho \neq 0$ ,  $|t| = 5.04$  with 6 degrees of freedom. From Table 7, we see that  $p\text{-value} < 2(.005) = .01$  so we can reject the null hypothesis that the correlation is 0.

**11.107** The summary statistics are  $S_{xx} = 153.875$ ,  $S_{xy} = 12.8$ , and  $S_{yy} = 1.34$ .

a. Thus,  $r = .89$ .

b. To test  $H_0: \rho = 0$  vs.  $H_a: \rho \neq 0$ ,  $|t| = 4.78$  with 6 degrees of freedom. From Table 7, we see that  $p\text{-value} < 2(.005) = .01$  so we can reject the null hypothesis that the correlation is 0.

**11.108** a.-c. Answers vary.