

Jack Melcher  
67574625  
EECS 101  
HW 6

1) a) What is the solid angle subtended by the moon as viewed from the earth if we assume the moon to be a sphere of radius  $R$  at a distance  $d$ ?

$$\Omega = \frac{A \cos \theta}{R^2}$$

$$\Omega = \frac{(\pi R^2) \cos(0)}{d^2}$$

$$\Omega = \frac{\pi R^2}{d^2}$$

b) What is the range of possible solid angles subtended by a flat circular plate of radius  $R$  at a distance  $d$ ?

$$\Omega = \frac{A \cos \theta}{R^2}$$

$$\frac{(\pi R^2) \cos(\pi)}{d^2} \leq \Omega \leq \frac{(\pi R^2) \cos(0)}{d^2}$$

$$0 \leq \Omega \leq \frac{(\pi R^2)}{d^2}$$

2) Consider a room in the shape of a cube of dimension 100 feet  $\times$  100 feet  $\times$  100 feet. Consider a square patch of size 1 foot by 1 foot on the ceiling. Suppose that the patch is exactly in the center of the ceiling.

a) What is the solid angle subtended by the square patch as viewed from a corner of the room on the floor?

$$\Omega = \frac{A \cos \theta}{R^2}$$

$$\Omega = \frac{(1^2) \frac{100}{\sqrt{15000}}}{(\sqrt{15000})^2}$$

$$\Omega = \frac{100}{15000^{3/2}}$$

b) What is the solid angle subtended by the square patch as viewed from a corner of the room on the ceiling?

$$\Omega = \frac{A \cos \theta}{R^2}$$

$$\Omega = \frac{(1^2) \cos(\pi)}{(\sqrt{15000})^2}$$

$$\Omega = 0$$

3) Consider a Lambertian plane in three dimensions defined by the equation

$$7x + \sqrt{50}y + z + 2 = 0$$

$$z = -7x - \sqrt{50}y - 2$$

a) What is the surface gradient (p, q) for the plane?

$$p = \frac{dz}{dx}, \quad q = \frac{dz}{dy}$$

$$p = -7, \quad q = -\sqrt{50}$$

b) Suppose that the plane is in a dark room with a single point light source. Consider the point P = (0, 0, -2) on the plane. Determine the location (x, y, z) where we should put the point light source so that the light source is a distance 20 from the point P and the reflected radiance from P in the direction of (0, 0, 0) is as large as possible.

Normal of given plane

$$n = (7, \sqrt{50}, 1)$$

Plane equation for light source

$$7(x - 20) + \sqrt{50}(y - 20) + z + 2 = 0$$

Point of light source

$$(20, 20, 2)$$

Radiance map equation

$$R(p, q) = \frac{1 + p_s p + q_s q}{\sqrt{1 + p_s^2 + q_s^2} \sqrt{1 + p^2 + q^2}}$$

Maximum when radiance map equals 1

This is satisfied when both p = p<sub>s</sub> and q = q<sub>s</sub>

$$1 = \frac{1 + 49 + 50}{\sqrt{1 + 49 + 50} \sqrt{1 + 49 + 50}}$$

$$1 = 1$$