

Generalised Pareto Distribution with Metropolis-Hastings MCMC updates

John O’Sullivan

November 4, 2019

Introduction

This document includes details on the model fitted by this code; namely, it describes in detail how we modelled a dataset of extreme observations using a generalised Pareto distribution and Metropolis-Hastings Markov chain Monte Carlo updates. Details for the Binomial component of the model are similar, and are omitted here for brevity.

The following pages include:

- the pseudocode in order to programme the algorithm;
- the model directed acyclic graph (DAG);
- details of the model notation;
- and the details of all equations needed for the MCMC updates of the parameters and the hyperparameters.

Algorithm details

Data: $z_k(x_i)$, the declustered threshold excesses at locations $x_i, i = 1 \dots n$, with $k = 1 \dots n_i$ excesses at location i ;

X_ϕ, X_ξ , $n \times (p + 1)$ and $n \times (q + 1)$ matrices of p and q covariates at locations $i = 1 \dots n$.

Result: Samples from the posterior distributions of $\phi = \log(\sigma)$ and ξ (the unknown parameters of interest), which can then be used to calculate return level estimates and other desired quantities

Initialisation;

Random starting values of ϕ and ξ ;

Hyperparameter values of $\alpha_\phi, \beta_\phi, \varsigma_\phi^2, \tau_\phi^2, \alpha_\xi, \beta_\xi, \varsigma_\xi^2$, and τ_ξ^2 ;

Number of iterations N ;

for iterations j from 1 to N **do**

 Generate $u \sim U(0, 1)$;

for grid locations i from 1 to n **do**

 Simulate $\phi_{new,i}$;

 Set $l_{new} = \log$ full conditional of new vector ϕ_{new} ;

 Set $l_{old} = \log$ full conditional of old vector ϕ ;

 Set $a = \exp(l_{new} - l_{old})$, that is, evaluate equation (??);

if $a > u$ **then**

 Set $\phi = \phi_{new}$;

end

 Simulate $\xi_{new,i}$;

 Set $l_{new} = \log$ full conditional of new vector ξ_{new} ;

 Set $l_{old} = \log$ full conditional of old vector ξ ;

 Set $a = \exp(l_{new} - l_{old})$, that is, evaluate equation (??);

if $a > u$ **then**

 Set $\xi = \xi_{new}$;

end

end

end

```

for iterations  $j$  from 1 to  $N$  (continued) do
  for each element  $c$  of the vector  $\alpha_\phi$  do
    Simulate  $\alpha_{\phi_{new},c}$ ;
    Set  $a$  = result of equation (??);
    If  $a > u$ , set  $\alpha_\phi = \alpha_{\phi_{new}}$ ;
  end

  for each element  $d$  of the vector  $\alpha_\xi$  do
    As above, but set  $a$  = result of equation (??);
  end

  for elements  $e$  in the lower-triangle of the matrix  $\beta_\phi$  do
    Simulate  $\beta_{\phi_{new},e}$ , repeat until symmetric positive definite  $\beta$  results;
    Set  $a$  = result of equation (??);
    If  $a > u$ , set  $\beta_\phi = \beta_{\phi_{new}}$ ;
  end

  for elements  $f$  in the lower-triangle of the matrix  $\beta_\xi$  do
    As above, but set  $a$  = result of equation (??);
  end

  Simulate  $\varsigma_{\phi_{new}}^2$ ;
  Set  $a$  = result of equation (??);
  If  $a > u$ , set  $\varsigma_\phi^2 = \varsigma_{\phi_{new}}^2$ ;
  Repeat for  $\varsigma_\xi^2$  with equation (??) ;

  Simulate  $\tau_{\phi_{new}}^2$ ;
  Set  $a$  = result of equation (??);
  If  $a > u$ , set  $\tau_\phi^2 = \tau_{\phi_{new}}^2$ ;
  Repeat for  $\tau_\xi^2$  with equation (??) ;

end

```

DAG




Figure 1: A directed acyclic graph (DAG) of the Bayesian spatial hierarchical models fitted to the wave data. On the left is the model for the excesses (using the GPD to model the data) and on the right is the model for the probability of an observation exceeding the threshold (using the Binomial distribution). The parameters of the distributions, modelled using Gaussian processes, are represented as circles in the middle layer, with the hyperparameters controlling these represented in the top layer. The data, modelled using the GPD for $z_k(x_i)$ and using a Binomial distribution for N_i , is represented in the bottom layer (in rectangles). Arrows run into nodes from their direct predecessors (often called parents). Given its parents, each node is independent of all other nodes in the graph except its descendants (often called children). Posterior estimates of the parameters' distributions can be used to form quantities of interest - typically return levels, as illustrated. Further details of each layer and the parameters involved may be found in the text.

Notation

- x_i Multivariate (bivariate or tri-variate) location values for location i , $i = 1, \dots, n$. Write the matrix of all locations as just x
- $z_k(x_i)$ k th observation at location i , $k = 1, \dots, n_i$ where n_i is the number of excesses at location i
- $\sigma(x_i)$ Scale parameter for location x_i
- $\phi(x_i) = \log(\sigma(x_i))$ The re-parameterised scale parameter
- $\xi(x_i)$ Shape parameter for location x_i
- $\mu_\phi(x), \mu_\xi(x)$ Means for the Gaussian processes
- Σ, Ψ Covariance matrices for the Gaussian processes
- τ_ϕ^2, τ_ξ^2 Nugget parameters for the Gaussian processes
- α_ϕ, α_ξ vectors of coefficients (including intercept) for Gaussian process means
- X_ϕ, X_ξ Matrices of covariates (including column for intercept term) on the x grid
- β_ϕ, β_ξ Length scale scalars or matrices
- $\varsigma_\phi^2, \varsigma_\xi^2$ Variance (partial sill) parameters for the Gaussian processes

Model outline

In hierarchical notation:

$$\begin{aligned}
z_k(x_i) &\sim GPD(\sigma(x_i), \xi(x_i)) \\
\log(\sigma(x)) = \phi(x) &\sim MVN(\mu_\phi(x), \Sigma(x, x)) \\
\xi(x) &\sim MVN(\mu_\xi(x), \Psi(x, x)) \\
\mu_\phi(x) &= X_\phi \alpha_\phi \\
\mu_\xi(x) &= X_\xi \alpha_\xi \\
\Sigma(x_k, x_l) &= \varsigma_\phi^2 \exp\left(-\left(\frac{d}{\beta_\phi}\right)^2\right) + \tau_\phi^2 \delta_{kl} \\
\Psi(x_k, x_l) &= \varsigma_\xi^2 \exp\left(-\left(\frac{d}{\beta_\xi}\right)^2\right) + \tau_\xi^2 \delta_{kl}
\end{aligned}$$

δ_{kl} is the Kronecker delta function. In the above formulae for Σ and Ψ , β_ϕ and β_ξ are treated as scalars. When considered as matrices, where d is the distance between two gridpoints, $(d/\beta)^2$ is replaced by $d^T \beta^{-1} d$.

Hyperparameter prior distributions (justifications for these values can be found in Section ??):

$$\begin{aligned}
\log(\varsigma_\phi^2), \log(\varsigma_\xi^2) &\sim N(0, 10) \\
\log(\tau_\phi^2), \log(\tau_\xi^2) &\sim N(0, 10) \\
\beta_\phi, \beta_\xi &\sim DU(0.001, 0.005, 0.01, 0.05, \dots, 100, 500, 1000) \\
\alpha_{\phi,j} &\sim N(0, 50) \\
\alpha_{\xi,j} &\sim N(0, 50)
\end{aligned}$$

Posterior distribution

The full posterior distribution is:

$$\begin{aligned}
p(\alpha_\phi, \alpha_\xi, \beta_\phi, \beta_\xi, \varsigma_\phi^2, \varsigma_\xi^2, \tau_\phi^2, \tau_\xi^2, \phi, \xi | x, z, X_\phi, X_\xi) \propto \\
\left[\prod_{i=1}^n \prod_{k=1}^{n_i} p(z_k(x_i) | \sigma(x_i) = \exp(\phi(x_i)), \xi(x_i)) \right] \times \\
p(\phi(x) | \mu_\phi, \Sigma) p(\xi(x) | \mu_\xi, \Psi) \times \\
p(\alpha_\phi) p(\alpha_\xi) p(\beta_\phi) p(\beta_\xi) \times \\
p(\tau_\phi^2) p(\tau_\xi^2) p(\varsigma_\phi^2) p(\varsigma_\xi^2)
\end{aligned}$$

Conditional posterior distributions: Layer 2

Updating the second layer of the DAG (parameters $\phi = \log(\sigma)$ and ξ): The conditional posterior distribution of ϕ is given by:

$$\begin{aligned}
\pi(\phi | z, x, X_\phi, X_\xi, \alpha_\phi, \alpha_\xi, \beta_\phi, \beta_\xi, \varsigma_\phi^2, \varsigma_\xi^2, \tau_\phi^2, \tau_\xi^2, \xi) \propto \\
p(z | \phi, x, X_\phi, X_\xi, \alpha_\phi, \alpha_\xi, \beta_\phi, \beta_\xi, \varsigma_\phi^2, \varsigma_\xi^2, \tau_\phi^2, \tau_\xi^2, \xi) \times \\
p(\phi | x, X_\phi, X_\xi, \alpha_\phi, \alpha_\xi, \beta_\phi, \beta_\xi, \varsigma_\phi^2, \varsigma_\xi^2, \tau_\phi^2, \tau_\xi^2, \xi) \propto \\
p(z | \sigma, \xi) p(\phi | \mu_\phi, \Sigma)
\end{aligned}$$

That is, the conditional posterior of ϕ is proportional to the product of the likelihood of the data z given ϕ and all other parameters, and the prior probability density of ϕ given all of the other parameters. In the final line, most parameters have dropped out from the right-hand side, as the densities are independent of these, given the remaining terms (see the DAG in Figure ?? for this). Remember that $\phi = \log(\sigma)$ throughout this, where σ is the scale parameter of the GPD, and note that μ_ϕ and Σ will have been calculated using the formulae in the model outline (above).

In briefer notation (to be used from now on):

$$\begin{aligned}
\pi(\phi|\dots) &\propto p(z|\sigma, \xi)p(\phi|\mu_\phi, \Sigma) \\
&= \left[\prod_{i=1}^n \prod_{k=1}^{n_i} p(z_k(x_i)|\sigma(x_i), \xi(x_i)) \right] p(\phi|\mu_\phi, \Sigma) \\
&= \left[\prod_{i=1}^n \prod_{k=1}^{n_i} \frac{1}{\sigma(x_i)} \left(1 + \xi(x_i) \frac{z_k(x_i)}{\sigma(x_i)} \right)^{-(1/\xi(x_i)+1)} \right] \times \\
&\quad \frac{1}{\sqrt{\det(2\pi\Sigma)}} e^{-\frac{1}{2}(\phi-\mu_\phi)'\Sigma^{-1}(\phi-\mu_\phi)}
\end{aligned}$$

Similarly:

$$\begin{aligned}
\pi(\xi|\dots) &\propto p(z|\phi, \xi)p(\xi|\mu_\xi, \Psi) \\
&= \left[\prod_{i=1}^n \prod_{k=1}^{n_i} p(z_k(x_i)|\sigma(x_i), \xi(x_i)) \right] p(\xi|\mu_\xi, \Psi) \\
&= \left[\prod_{i=1}^n \prod_{k=1}^{n_i} \frac{1}{\sigma(x_i)} \left(1 + \xi(x_i) \frac{z_k(x_i)}{\sigma(x_i)} \right)^{-(1/\xi(x_i)+1)} \right] \times \\
&\quad \frac{1}{\sqrt{\det(2\pi\Psi)}} e^{-\frac{1}{2}(\xi-\mu_\xi)'\Psi^{-1}(\xi-\mu_\xi)}
\end{aligned}$$

Metropolis-Hastings Markov chain Monte Carlo sampling

In order to sample from these conditional posterior distributions, we use the Metropolis-Hastings Markov chain Monte Carlo (MCMC) algorithm. New samples are accepted or rejected at random according to the algorithm outlined below.

A new value of ϕ_i is suggested: ϕ'_i , where i is a location on the grid, resulting in a new vector ϕ' .

Suggested updates are drawn from a Normal distribution centred on the old value and with a variance of a manually set tuning parameter used to control the size of the proposed steps.

We calculate:

$$\rho(\phi_i, \phi'_i) = \min \left(1, \frac{\pi(\phi' | \dots) q_t(\phi'_i \rightarrow \phi_i)}{\pi(\phi | \dots) q_t(\phi_i \rightarrow \phi'_i)} \right)$$

where $\pi(\phi | \dots)$ is as defined above, and $q_t(a \rightarrow b)$ is the transition probability of proposing value b given value a . Since updates are proposed using a Normal distribution, these transition probabilities above and below the line will always cancel, so the above simplifies to:

$$\rho(\phi_i, \phi'_i) = \min \left(1, \frac{\pi(\phi' | \dots)}{\pi(\phi | \dots)} \right)$$

Following this calculation, we always accept proposed value ϕ'_i when $\rho(\phi_i, \phi'_i)$ equals 1 and we reject accordingly when the ratio is smaller than 1 by simulating a random variable $u \sim U[0, 1]$ and accepting proposed value ϕ'_i when $u \leq \rho(\phi_i, \phi'_i)$.

Evaluating $\rho(\phi_i, \phi'_i)$ typically involves products and quotients of many terms which may be close to 0. In order to work with something far more computationally stable, we use the property that $x = \exp(\log(x))$. Note also that since the vectors ϕ and ϕ' differ in only one entry, the GPD calculations above and below will cancel for all other gridpoints apart from the one being updated, i .

Following these observations we need to evaluate:

$$\begin{aligned}
& \exp \left(\log \left(\frac{\pi(\phi' | \dots)}{\pi(\phi | \dots)} \right) \right) \\
&= \exp \left[\log(\pi(\phi' | \dots)) - \log(\pi(\phi | \dots)) \right] \\
&= \exp \left[\log \left(\left[\prod_{i=1}^n \prod_{k=1}^{n_i} p(z_k(x_i) | \sigma'(x_i), \xi(x_i)) \right] p(\phi' | \mu_\phi, \Sigma) \right) - \right. \\
&\quad \left. \log \left(\left[\prod_{i=1}^n \prod_{k=1}^{n_i} p(z_k(x_i) | \sigma(x_i), \xi(x_i)) \right] p(\phi | \mu_\phi, \Sigma) \right) \right] \\
&= \exp \left[\log \left(\left[\prod_{k=1}^{n_i} p(z_k(x_i) | \sigma'(x_i), \xi(x_i)) \right] p(\phi' | \mu_\phi, \Sigma) \right) - \right. \\
&\quad \left. \log \left(\left[\prod_{k=1}^{n_i} p(z_k(x_i) | \sigma(x_i), \xi(x_i)) \right] p(\phi | \mu_\phi, \Sigma) \right) \right]
\end{aligned}$$

Filling in the distribution details, this becomes:

$$\begin{aligned}
& \exp \left[\log \left(\left[\prod_{k=1}^{n_i} \frac{1}{\sigma'(x_i)} \left(1 + \xi(x_i) \frac{z_k(x_i)}{\sigma'(x_i)} \right)^{-(1/\xi(x_i)+1)} \right] \times \right. \right. \\
& \quad \left. \frac{1}{\sqrt{\det(2\pi\Sigma)}} e^{-\frac{1}{2}(\phi' - \mu_\phi)' \Sigma^{-1} (\phi' - \mu_\phi)} \right) \\
& \quad - \log \left(\left[\prod_{k=1}^{n_i} \frac{1}{\sigma(x_i)} \left(1 + \xi(x_i) \frac{z_k(x_i)}{\sigma(x_i)} \right)^{-(1/\xi(x_i)+1)} \right] \times \right. \\
& \quad \left. \left. \frac{1}{\sqrt{\det(2\pi\Sigma)}} e^{-\frac{1}{2}(\phi - \mu_\phi)' \Sigma^{-1} (\phi - \mu_\phi)} \right) \right] \\
& = \exp \left[\sum_{k=1}^{n_i} \log \left(\left[\frac{1}{\sigma'(x_i)} \left(1 + \xi(x_i) \frac{z_k(x_i)}{\sigma'(x_i)} \right)^{-(1/\xi(x_i)+1)} \right] \right) + \right. \\
& \quad \left. \log \left(\frac{1}{\sqrt{\det(2\pi\Sigma)}} \right) - \frac{1}{2}(\phi' - \mu_\phi)' \Sigma^{-1} (\phi' - \mu_\phi) \right. \\
& \quad - \sum_{k=1}^{n_i} \log \left(\left[\frac{1}{\sigma(x_i)} \left(1 + \xi(x_i) \frac{z_k(x_i)}{\sigma(x_i)} \right)^{-(1/\xi(x_i)+1)} \right] \right) - \\
& \quad \left. \log \left(\frac{1}{\sqrt{\det(2\pi\Sigma)}} \right) + \frac{1}{2}(\phi - \mu_\phi)' \Sigma^{-1} (\phi - \mu_\phi) \right] \\
& = \exp \left[\sum_{k=1}^{n_i} \log \left(\left[\frac{1}{\sigma'(x_i)} \left(1 + \xi(x_i) \frac{z_k(x_i)}{\sigma'(x_i)} \right)^{-(1/\xi(x_i)+1)} \right] \right) - \right. \\
& \quad \frac{1}{2}(\phi' - \mu_\phi)' \Sigma^{-1} (\phi' - \mu_\phi) - \\
& \quad \sum_{k=1}^{n_i} \log \left(\left[\frac{1}{\sigma(x_i)} \left(1 + \xi(x_i) \frac{z_k(x_i)}{\sigma(x_i)} \right)^{-(1/\xi(x_i)+1)} \right] \right) + \\
& \quad \left. \frac{1}{2}(\phi - \mu_\phi)' \Sigma^{-1} (\phi - \mu_\phi) \right]
\end{aligned}$$

The common term $\log(1/\sqrt{\det(2\pi\Sigma)})$ in both numerator and denominator above dropped out as it will be equal in both (it has no dependence on ϕ).

The GPD can clearly be simplified further using the properties of logs. Just taking one of the functions on its own for clarity:

$$\begin{aligned}
& \sum_{k=1}^{n_i} \log \left(\left[\frac{1}{\sigma'(x_i)} \left(1 + \xi(x_i) \frac{z_k(x_i)}{\sigma'(x_i)} \right)^{-(1/\xi(x_i)+1)} \right] \right) = \\
& \sum_{k=1}^{n_i} \left[\log \left(\frac{1}{\sigma'(x_i)} \right) + \log \left(1 + \xi(x_i) \frac{z_k(x_i)}{\sigma'(x_i)} \right)^{-(1/\xi(x_i)+1)} \right] = \\
& \sum_{k=1}^{n_i} \left[-\log(\sigma'(x_i)) - \left(\frac{1}{\xi(x_i)} + 1 \right) \log \left(1 + \xi(x_i) \frac{z_k(x_i)}{\sigma'(x_i)} \right) \right]
\end{aligned}$$

Using this form, the full term we need to evaluate in order to update ϕ is:

$$\begin{aligned}
= \exp & \left[\sum_{k=1}^{n_i} \left[-\log(\sigma'(x_i)) - \left(\frac{1}{\xi(x_i)} + 1 \right) \log \left(1 + \xi(x_i) \frac{z_k(x_i)}{\sigma'(x_i)} \right) \right] - \right. \\
& \frac{1}{2}(\phi' - \mu_\phi)' \Sigma^{-1}(\phi' - \mu_\phi) - \\
& \sum_{k=1}^{n_i} \left[-\log(\sigma(x_i)) - \left(\frac{1}{\xi(x_i)} + 1 \right) \log \left(1 + \xi(x_i) \frac{z_k(x_i)}{\sigma(x_i)} \right) \right] + \\
& \left. \frac{1}{2}(\phi - \mu_\phi)' \Sigma^{-1}(\phi - \mu_\phi) \right] \tag{1}
\end{aligned}$$

Similar reasoning leads to the update for ξ . We need to evaluate:

$$\rho(\xi_i, \xi'_i) = \min \left(1, \frac{\pi(\xi' | \dots) q_t(\xi'_i \rightarrow \xi_i)}{\pi(\xi | \dots) q_t(\xi_i \rightarrow \xi'_i)} \right)$$

The full term we need to evaluate is:

$$\begin{aligned}
= \exp & \left[\sum_{k=1}^{n_i} \left[-\log(\sigma(x_i)) - \left(\frac{1}{\xi'(x_i)} + 1 \right) \log \left(1 + \xi'(x_i) \frac{z_k(x_i)}{\sigma(x_i)} \right) \right] - \right. \\
& \frac{1}{2}(\xi' - \mu_\xi)' \Psi^{-1}(\xi' - \mu_\xi) - \\
& \sum_{k=1}^{n_i} \left[-\log(\sigma(x_i)) - \left(\frac{1}{\xi(x_i)} + 1 \right) \log \left(1 + \xi(x_i) \frac{z_k(x_i)}{\sigma(x_i)} \right) \right] + \\
& \left. \frac{1}{2}(\xi - \mu_\xi)' \Psi^{-1}(\xi - \mu_\xi) \right] \tag{2}
\end{aligned}$$

Conditional posterior distributions: Layer 3

α

Updating the third layer of the DAG (that is, all hyperparameters) starting with α_ϕ :

$$\begin{aligned}\pi(\alpha_\phi|z, \dots) &\propto p(z|\alpha_\phi, \dots)p(\alpha_\phi|\dots) \\ &\propto p(z|\sigma, \xi)p(\phi|\mu_\phi, \Sigma)p(\alpha_\phi) \\ &\propto p(\phi|\mu_\phi, \Sigma)p(\alpha_\phi)\end{aligned}$$

Essentially the GPD component is independent of α_ϕ once the other parameters are known, and so can be absorbed into the constant of proportionality. What we're left with is the *MVN* piece (since α_ϕ features in the calculation of μ_ϕ) and the prior on α_ϕ .

Then we have:

$$\begin{aligned}\pi(\alpha_\phi|\dots) &\propto p(\phi|\mu_\phi, \Sigma)p(\alpha_\phi) \\ &\propto \frac{1}{\sqrt{\det(2\pi\Sigma)}} e^{-\frac{1}{2}(\phi-\mu_\phi)'\Sigma^{-1}(\phi-\mu_\phi)} \times \\ &\quad \frac{1}{\sqrt{2\pi s^2}} e^{-\frac{(\alpha_\phi-m)^2}{2s^2}}\end{aligned}$$

where m is the prior mean and s^2 is the prior variance. In a slight abuse of notation in the final line above, α_ϕ is a scalar, the single element which is being updated from the vector α_ϕ .

Similarly:

$$\begin{aligned}\pi(\alpha_\xi|\dots) &\propto p(\xi|\mu_\xi, \Psi)p(\alpha_\xi) \\ &\propto \frac{1}{\sqrt{\det(2\pi\Psi)}} e^{-\frac{1}{2}(\xi-\mu_\xi)'\Psi^{-1}(\xi-\mu_\xi)} \times \\ &\quad \frac{1}{\sqrt{2\pi s^2}} e^{-\frac{(\alpha_\xi-m)^2}{2s^2}}\end{aligned}$$

where m is the prior mean and s^2 is the prior variance, and a similar slight abuse of notation applies in the final line for α_ξ .

As with the parameters ϕ and ξ , updates for α_ϕ are suggested element-wise: $\alpha_{\phi,\kappa} \rightarrow \alpha'_{\phi,\kappa}$, where the index κ runs over the vector of coefficients. The ratio $\rho(\alpha_{\phi,\kappa}, \alpha'_{\phi,\kappa})$ is then calculated. Similar manipulations to those used in the previous section lead to the following calculation needed to update α_ϕ :

$$\begin{aligned} & \exp \left[-\frac{1}{2} \log(\det(2\pi\Sigma)) - \frac{1}{2}(\phi - \mu'_\phi)' \Sigma^{-1}(\phi - \mu'_\phi) \right. \\ & \quad - \frac{1}{2} \log(2\pi s^2) - \frac{(\alpha'_{\phi,\kappa} - m)^2}{2s^2} \\ & \quad + \frac{1}{2} \log(\det(2\pi\Sigma)) + \frac{1}{2}(\phi - \mu_\phi)' \Sigma^{-1}(\phi - \mu_\phi) \\ & \quad \left. + \frac{1}{2} \log(2\pi s^2) + \frac{(\alpha_{\phi,\kappa} - m)^2}{2s^2} \right] = \\ & \exp \left[-\frac{1}{2}(\phi - \mu'_\phi)' \Sigma^{-1}(\phi - \mu'_\phi) - \frac{(\alpha'_{\phi,\kappa} - m)^2}{2s^2} \right. \\ & \quad \left. + \frac{1}{2}(\phi - \mu_\phi)' \Sigma^{-1}(\phi - \mu_\phi) + \frac{(\alpha_{\phi,\kappa} - m)^2}{2s^2} \right] \end{aligned} \quad (3)$$

And for α_ξ :

$$\begin{aligned} & \exp \left[-\frac{1}{2}(\xi - \mu'_\xi)' \Psi^{-1}(\xi - \mu'_\xi) - \frac{(\alpha'_{\xi,\kappa} - m)^2}{2s^2} \right. \\ & \quad \left. + \frac{1}{2}(\xi - \mu_\xi)' \Psi^{-1}(\xi - \mu_\xi) + \frac{(\alpha_{\xi,\kappa} - m)^2}{2s^2} \right] \end{aligned} \quad (4)$$

Updates for the other hyperparameters are similar. The matrices Σ and Ψ will change with any change in β, ζ^2 or τ^2 . The remaining hyperparameters either have a discrete update (in which case the prior probability will be $1/\#discrete.values$ and will cancel above and below the line, or have a univariate Normal prior on them (or their log).

β

For β_ϕ we have:

$$\pi(\beta_\phi | \dots) \propto p(\phi | \mu_\phi, \Sigma) p(\beta_\phi)$$

Since β_ϕ has a discrete prior, to update β_ϕ we need to evaluate:

$$\begin{aligned} \exp \left[-\frac{1}{2} \log(\det(2\pi\Sigma')) - \frac{1}{2}(\phi - \mu_\phi)' \Sigma'^{-1}(\phi - \mu_\phi) \right. \\ \left. + \frac{1}{2} \log(\det(2\pi\Sigma)) + \frac{1}{2}(\phi - \mu_\phi)' \Sigma^{-1}(\phi - \mu_\phi) \right] \end{aligned} \quad (5)$$

where Σ' has been formed using β'_ϕ , the new proposal value or matrix.

Then for β_ξ we have:

$$\pi(\beta_\xi | \dots) \propto p(\xi | \mu_\xi, \Psi) p(\beta_\xi)$$

To update β_ξ we need to evaluate:

$$\begin{aligned} \exp \left[-\frac{1}{2} \log(\det(2\pi\Psi')) - \frac{1}{2}(\xi - \mu_\xi)' \Psi'^{-1}(\xi - \mu_\xi) \right. \\ \left. + \frac{1}{2} \log(\det(2\pi\Psi)) + \frac{1}{2}(\xi - \mu_\xi)' \Psi^{-1}(\xi - \mu_\xi) \right] \end{aligned} \quad (6)$$

where Ψ' has been formed using β'_ξ , the new proposal value or matrix.

ς^2

ς_ϕ^2 has a conditional posterior distribution of:

$$\pi(\varsigma_\phi^2 | \dots) \propto p(\phi | \mu_\phi, \Sigma) p(\varsigma_\phi^2)$$

The prior distribution of the log of ς_ϕ^2 is a univariate Normal. So to update ς_ϕ^2 we need to evaluate:

$$\begin{aligned} \exp \left[-\frac{1}{2} \log(\det(2\pi\Sigma')) - \right. \\ \left. \frac{1}{2}(\phi - \mu_\phi)' \Sigma'^{-1}(\phi - \mu_\phi) - \frac{(\log(\varsigma_\phi^{2'}) - m)^2}{2s^2} + \right. \\ \left. \frac{1}{2} \log(\det(2\pi\Sigma)) + \right. \\ \left. \frac{1}{2}(\phi - \mu_\phi)' \Sigma^{-1}(\phi - \mu_\phi) + \frac{(\log(\varsigma_\phi^2) - m)^2}{2s^2} \right] \end{aligned} \quad (7)$$

where Σ' has been formed using $\varsigma_\phi^{2'}$, the new proposal value, and m and s are the prior mean and standard deviation repectively.

ς_ξ^2 has a conditional posterior distribution of:

$$\pi(\varsigma_\xi^2 | \dots) \propto p(\xi | \mu_\xi, \Psi) p(\varsigma_\xi^2)$$

To update ς_ξ^2 we need to evaluate:

$$\begin{aligned} \exp \left[-\frac{1}{2} \log(\det(2\pi\Psi')) - \right. \\ \left. \frac{1}{2}(\xi - \mu_\xi)' \Psi'^{-1}(\xi - \mu_\xi) - \frac{(\log(\varsigma_\xi^{2'}) - m)^2}{2s^2} + \right. \\ \left. \frac{1}{2} \log(\det(2\pi\Psi)) + \right. \\ \left. \frac{1}{2}(\xi - \mu_\xi)' \Psi^{-1}(\xi - \mu_\xi) + \frac{(\log(\varsigma_\xi^2) - m)^2}{2s^2} \right] \end{aligned} \quad (8)$$

where Ψ' has been formed using $\varsigma_\xi^{2'}$, the new proposal value, and m and s are the prior mean and standard deviation repectively.

τ^2

τ_ϕ^2 has a conditional posterior distribution of:

$$\pi(\tau_\phi^2 | \dots) \propto p(\phi | \mu_\phi, \Sigma) p(\tau_\phi^2)$$

The prior distribution of the log of τ_ϕ^2 is a univariate Normal. So to update τ_ϕ^2 we need to evaluate:

$$\begin{aligned} \exp \left[-\frac{1}{2} \log(\det(2\pi\Sigma')) - \right. \\ \left. \frac{1}{2} (\phi - \mu_\phi)' \Sigma'^{-1} (\phi - \mu_\phi) - \frac{(\log(\tau_\phi^{2'}) - m)^2}{2s^2} + \right. \\ \left. \frac{1}{2} \log(\det(2\pi\Sigma)) + \right. \\ \left. \frac{1}{2} (\phi - \mu_\phi)' \Sigma^{-1} (\phi - \mu_\phi) + \frac{(\log(\tau_\phi^2) - m)^2}{2s^2} \right] \end{aligned} \quad (9)$$

where Σ' has been formed using $\tau_\phi^{2'}$, the new proposal value, and m and s are the prior mean and standard deviation repectively.

τ_ξ^2 has a conditional posterior distribution of:

$$\pi(\tau_\xi^2 | \dots) \propto p(\xi | \mu_\xi, \Psi) p(\tau_\xi^2)$$

To update τ_ξ^2 we need to evaluate:

$$\exp \left[-\frac{1}{2} \log(\det(2\pi\Psi')) - \frac{1}{2}(\xi - \mu_\xi)' \Psi'^{-1}(\xi - \mu_\xi) - \frac{(\log(\tau_\xi^{2'}) - m)^2}{2s^2} + \frac{1}{2} \log(\det(2\pi\Psi)) + \frac{1}{2}(\xi - \mu_\xi)' \Psi^{-1}(\xi - \mu_\xi) + \frac{(\log(\tau_\xi^2) - m)^2}{2s^2} \right] \quad (10)$$

where Ψ' has been formed using $\tau_\xi^{2'}$, the new proposal value, , and m and s are the prior mean and standard deviation repectively.