

risks 1990		risks 2019	Mean percentage change in number of DALYs, 1990-2019	Mean percentage change in all-age of DALYs rate, 1990-2019
1 Child and maternal malnutrition		1 Child and maternal malnutrition	1.26	1.21
2 Air pollution		2 High systolic blood pressure	-0.35	-0.36
3 Unsafe water, sanitation, and handwashing		3 Tobacco	-0.13	-0.15
4 Tobacco		4 Air pollution	0.32	0.29
5 High systolic blood pressure		5 Dietary risks	-0.3	-0.31
6 Dietary risks		6 High fasting plasma glucose	-0.55	-0.56
7 High fasting plasma glucose		7 High body-mass index	-0.58	-0.59
8 High LDL cholesterol		8 High LDL cholesterol	-0.29	-0.31
9 Alcohol use		9 Alcohol use	-0.27	-0.29
10 High body-mass index		10 Unsafe water, sanitation, and handwashing	1.46	1.4
11 Occupational risks		11 Kidney dysfunction	-0.45	-0.46
12 Kidney dysfunction		12 Occupational risks	-0.05	-0.07
13 Non-optimal temperature		13 Unsafe sex	-0.57	-0.58
14 Unsafe sex		14 Non-optimal temperature	0.08	0.05
15 Other environmental risks		15 Drug use	-0.46	-0.47
16 Drug use		16 Other environmental risks	-0.27	-0.28
17 Low physical activity		17 Low bone mineral density	-0.48	-0.5
18 Low bone mineral density		18 Low physical activity	-0.45	-0.47
19 Intimate partner violence		19 Intimate partner violence	-0.39	-0.4
20 Childhood sexual abuse and bullying		20 Childhood sexual abuse and bullying	-0.36	-0.37