

# Markov Link Method for combining destructive measurements

Jackson Loper, Trygve Bakken, and Liam Paninski  
Columbia University

## Destructive measurements

- It is easy to calibrate thermometers
- RNAseq methods? Not so easy

## Setup

- $\ell$  – side information (region of brain, cre line, etc.)
- $X$  – result of experiment under one modality
- $Y$  – result of experiment under another modality

## Setup

- $\ell$  – side information (region of brain, cre line, etc.)
- $X$  – result of experiment under one modality
- $Y$  – result of experiment under another modality

## “Calibration”

$$\mathbb{Q}(Y = y|X = x) = ??$$

## Setup

- $\ell$  – side information (region of brain, cre line, etc.)
- $X$  – result of experiment under one modality
- $Y$  – result of experiment under another modality

## “Calibration”

$$\mathbb{Q}(Y = y | X = x) = ??$$

Problem: we can never observe  $X, Y$  together

## One solution: Markov Link Method Assumption

If  $X$  is *more fine-grained* than  $\ell$ , then in some cases we may assume that

$$\mathbb{P}(X, Y|\ell) = \mathbb{P}(X|\ell)\mathbb{P}(Y|X)$$

## One solution: Markov Link Method Assumption

If  $X$  is *more fine-grained* than  $\ell$ , then in some cases we may assume that

$$\mathbb{P}(X, Y|\ell) = \mathbb{P}(X|\ell)\mathbb{P}(Y|X)$$

And we can estimate the set of calibrations:

$$\Theta = \left\{ \mathbb{Q} : \mathbb{P}(Y|\ell) = \sum_X \mathbb{P}(X|\ell)\mathbb{Q}(Y|X) \right\}$$

## Analyze $\Theta$

- Rotationally Uniform eXtremal distribution
- Uniform distribution
- Diameter estimation
- Center of mass



## Empirical results

Tasic, Bosiljka, Zizhen Yao, Kimberly A. Smith, Lucas Graybuck, Thuc Nghi Nguyen, Darren Bertagnolli, Jeff Goldy et al. "Shared and distinct transcriptomic cell types across neocortical areas." bioRxiv (2017): 229542.

From this:



