

The Drinfeld center and topological symmetries

Jackson Van Dyke
UT Austin

April 24, 2023

3-dimensional TQFT

- Consider a 3-dimensional (framed) TQFT:

$$\mathbf{Bord}_3^{\text{fr}} \xrightarrow{F} \mathbf{Alg}_1(\mathbf{Cat}) .$$

- The point goes to some monoidal category:

$$F(\bullet) = (\mathcal{C}, *) .$$

- The interval goes to the identity bimodule:

$$F(\bullet \text{---} \bullet) = {}_c\mathcal{C}_c .$$

- The circle will be sent to some category:

$$F\left(\bigcirc\right) = ? \in \mathbf{Cat} \cong \mathbf{End}_{\mathbf{Alg}_1(\mathbf{Cat})}(1) .$$

Example of a 3-dimensional TQFT

- Before we identify $F(S^1)$, let's consider an example.
- Consider the category of vector spaces graded by a finite abelian group G :

$$\mathcal{C} = \mathbf{Vect}[G] . \quad (1)$$


The simple objects are given by 'skyscrapers' \mathbb{C}_g for $g \in G$.

- This has a tensor product given by convolution $*$. On simple objects \mathbb{C}_g it is simply:

$$\mathbb{C}_g * \mathbb{C}_h = \mathbb{C}_{gh} . \quad (2)$$

- The TQFT F associated to this particular fusion category is finite gauge theory with gauge group G . If τ is a cocycle for a class in $H^3(BG, \mathbb{C}^\times)$, we can define a nontrivial associator for $\mathbf{Vect}[G]$ using τ , resulting in *Dijkgraaf-Witten* theory for (G, τ) .

The assignment to the circle



- From this picture, we have an action:

$$F(\bigcirc) \xrightarrow{\sim} \text{End}_{c\text{-bimod}}(c\mathcal{C}) .$$

- This map turns out to be an equivalence.[DSS20, Section 3.2.2]
- This has more structure, e.g. a product map given by composition, which we will discuss in a couple of slides.
- But first, let's notice: this is the Drinfeld center!

The Drinfeld center

- The *Drinfeld center* of a tensor category $(\mathcal{C}, *)$ is:

$$\mathcal{Z}(\mathcal{C}) = \text{End}_{\mathcal{C}\text{-bimod}}({}_\mathcal{C}\mathcal{C}_\mathcal{C}) = \text{End}_{\mathcal{C} \otimes \mathcal{C}^{\text{op}}}(\mathcal{C}) .$$

- So the upshot of the previous slide is:

$$F(\bigcirc) \cong \mathcal{Z}(F(\bullet) = \mathcal{C}) .$$

- The Drinfeld center has a more concrete description: consider the category with objects given by pairs (X, σ_X) , where X is an object of \mathcal{C} , and σ_X is a natural transformation:

$$\sigma_X: X \otimes (-) \rightarrow (-) \otimes X .$$

The morphisms are (appropriately compatible) morphisms in \mathcal{C} . See [Eti+15, Prop. 7.13.8] for the equivalence between the two definitions.

Extra structure on $F(S^1)$

- We have seen that if $F(\bullet) = \mathcal{C}$, then:

$$F(\bigcirc) \cong \mathcal{Z}(\mathcal{C}) = \text{End}_{\mathcal{C} \otimes \mathcal{C}^{\text{op}}}(\mathcal{C}) .$$

- This is naturally a monoidal category: composition of endomorphisms is the same as the multiplication map induced by the pair of pants bordism:

$$F\left(\text{pair of pants bordism}\right) : F(\bigcirc) \otimes F(\bigcirc) \rightarrow F(\bigcirc) .$$

- In fact there is even a *braiding*, induced by moving one of the “legs” around the other.

Back to the example

- Recall our finite abelian gauge theory example: $F(\bullet) = \mathbf{Vect}[G]$.
- The Drinfeld center of this fusion category turns out to be:

$$\mathcal{Z}(\mathbf{Vect}[G]) \cong \mathbf{Vect}[G \oplus G^\vee] ,$$

where $G^\vee = \mathrm{Hom}(G, \mathbb{C}^\times)$ is the character dual.

- The monoidal structure is still convolution, and the braiding is given on simple objects by:

$$\mathbb{C}_{(g,\chi)} * \mathbb{C}_{(h,\omega)} \xrightarrow{\chi(h)\omega(g)\mathrm{id}} \mathbb{C}_{(h,\omega)} * \mathbb{C}_{(g,\chi)} . \quad (3)$$

What does any of this have to do with symmetry?

- A boundary theory $1 \rightarrow F$ should be thought of as a “2-dimensional theory with a \mathcal{C} -action”: usually a 2d theory sends the point to a category, and now it is sent to a \mathcal{C} -module category.
- In the finite gauge theory example, where $\mathcal{C} = \mathbf{Vect}[G]$, a \mathcal{C} -module structure on a category can be thought of as a categorical action of G itself.

Boundary theories for the 3-dimensional theory

- $F: \bullet \mapsto \mathcal{C}$ is the *Turaev-Viro (TV) theory associated to \mathcal{C}* .
- The theory sending the circle to a particular braided category \mathcal{B} is the *Reshetikhin-Turaev (RT) theory associated to \mathcal{B}* .
- So the RT theory for $\mathcal{Z}(\mathcal{C})$ agrees with the TV theory for \mathcal{C} , but not all RT theories may be of this form.

Theorem ([FT21])

An RT theory admits a nonzero boundary theory if and only if it is a TV theory.

4-dimensional theory associated to the Drinfeld center

- The Drinfeld center is a braided category, and turns out to be sufficiently dualizable [BJS21] to define a 4-dimensional TQFT:

$$\alpha: \mathbf{Bord}_4 \ni \bullet \mapsto \mathcal{Z}(\mathcal{C}) \in \mathbf{Alg}_2(\mathbf{Cat}) .$$

- This is the *Crane-Yetter (CY) theory* associated to the braided category $\mathcal{Z}(\mathcal{C})$.

Upgrading the 3-dimensional theory to a boundary theory

- The Drinfeld center of a tensor category manifestly acts on the original tensor category, since we have a forgetful functor

$$\mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C} .$$

- In terms of the theories, this means that F can be upgraded to a boundary condition:

$$\tilde{F}: 1 \rightarrow \alpha .$$

The value of \tilde{F} on the point is \mathcal{C} as a $\mathcal{Z}(\mathcal{C})$ -module.

- More specifically there is a (α, ρ) -module structure on F , in the sense of [FMT22].

Back to the example

- Consider our running example: if F is G -gauge theory, then the theory

$$\alpha: \text{pt} \mapsto \mathcal{Z}(\mathbf{Vect}[G]) \cong \mathbf{Vect}[G \oplus G^\vee] ,$$

can be described as the quantization (in the sense of [Fre+10]) of the groupoid $B^2(G \oplus G^\vee)$, twisted by a cocycle for the class

$$\text{ev} \in \text{Hom}(G \oplus G^\vee, \mathbb{C}^\times) \cong H^4(B^2(G \oplus G^\vee), \mathbb{C}^\times) .$$

- Lagrangian subgroups L of $(G \oplus G^\vee, \text{ev})$ now give rise to boundary theories $1 \rightarrow \alpha$, by quantizing the correspondence (as in [FMT22]):

$$\bullet \leftarrow B^2 L \rightarrow B^2(G \oplus G^\vee) .$$

- The boundary theories corresponding to $L = G$ and G^\vee are related by a “twice-categorified finite Fourier transform”, which is studied in detail in my upcoming work.

The running example as an anomalous theory

- We might wonder if the theory F can be upgraded to have an action of the automorphisms of the center or, more concretely, of the group $O(G \oplus G^\vee)$.
- At the level of the fusion category itself, this is answered by [ENO10]: this group acts if and only if specific obstructions are trivializable, and an action is determined by a trivialization.
- Passing the obstruction theory from [ENO10] through the “quantization of groupoids” formalism developed in [Fre+10] yields a collection of anomaly theories and symmetry theories for the theory associated to the fusion category we started with.
- This is spelled out in my upcoming work.

Summary:

TV for \mathcal{C} :

- ① governs \mathcal{C} -symmetry,
- ② is equivalent to RT for $\mathcal{Z}(\mathcal{C})$, and
- ③ is (can be upgraded to) a boundary theory for CY for $\mathcal{Z}(\mathcal{C})$.

References

- [BJS21] Adrien Brochier, David Jordan, and Noah Snyder. “On dualizability of braided tensor categories”. In: *Compos. Math.* 157.3 (2021), pp. 435–483. ISSN: 0010-437X. DOI: 10.1112/s0010437x20007630.
- [DSS20] Christopher L. Douglas, Christopher Schommer-Pries, and Noah Snyder. “Dualizable tensor categories”. In: *Mem. Amer. Math. Soc.* 268.1308 (2020), pp. vii+88.
- [ENO10] Pavel Etingof, Dmitri Nikshych, and Victor Ostrik. “Fusion categories and homotopy theory”. In: *Quantum Topol.* 1.3 (2010). With an appendix by Ehud Meir, pp. 209–273. ISSN: 1663-487X. DOI: 10.4171/QT/6.
- [Eti+15] Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik. *Tensor categories*. Vol. 205. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2015, pp. xvi+343.
- [FMT22] Daniel S. Freed, Gregory W. Moore, and Constantin Teleman. *Topological symmetry in quantum field theory*. 2022. DOI: 10.48550/ARXIV.2209.07471. URL: <https://arxiv.org/abs/2209.07471>.
- [Fre+10] Daniel S. Freed, Michael J. Hopkins, Jacob Lurie, and Constantin Teleman. “Topological quantum field theories from compact Lie groups”. In: *A celebration of the mathematical legacy of Raoul Bott*. Vol. 50. CRM Proc. Lecture Notes. Amer. Math. Soc., Providence, RI, 2010, pp. 367–403.
- [FT21] Daniel S. Freed and Constantin Teleman. “Gapped boundary theories in three dimensions”. In: *Comm. Math. Phys.* 388.2 (2021), pp. 845–892.