

Orderability and 3-manifold groups

Lectures by: Professor Cameron Gordon

Guest lectures given by Hannah Turner and Jonathan Johnson. Notes by: Jackson Van Dyke; All errors introduced are my own.

Contents

Chapter 1. Orders on groups; basic definitions and properties	4
1. Orderability of manifold groups	8
2. Three-manifold groups	9
3. Group rings	12
4. BO's on $\mathbb{Z} \times \mathbb{Z}$	14
5. BO's on \mathbb{R}	15
Chapter 2. The space of left-orders on a group	16
1. The cantor set	18
2. Surface groups	23
Chapter 3. Three-manifolds	27
1. Higher homotopy groups	28
2. Back to three-manifolds	29
Chapter 4. Seifert fiber(ed) spaces	34
Appendix A. Orderings of the braid group	40
1. Dehornoy's ordering	41
2. Nielsen-Thurston orderings on B_n	43
3. Isolated orderings	43
Appendix B. Orderability and knot groups	46
1. Generalized torsion	46
2. Knot groups as extensions	47
Appendix. Bibliography	50

CHAPTER 1

Orders on groups; basic definitions and properties

The book for the course is [2].

Lecture 1; January
21, 2020

Recall that a *strict total order* (STO) on a set X is a binary relation $<$ which satisfies:

- (1) $x < y$ and $y < z$ implies $x < z$;
- (2) $\forall x, y \in X$ exactly one of: $x < y$, $y < x$, $x = y$, holds.

A *left order* (LO) on a group G is an STO such that $g < h$ implies $fg < fh$ for all $f \in G$. G is *left-orderable* (LO) if there exists an LO on G . We similarly define a *right order* (RO) and *right orderability* (RO). A *bi-order* (BO) on G is an LO on G that is also an RO.

REMARK 1.1. (1) If G is abelian, $<$ is a LO iff $<$ is an RO iff $<$ is a BO.
(2) If $<$ is an LO on G , then \prec defined by:

$$(1.1) \quad g \prec h \iff h^{-1} < g^{-1}$$

is an RO on G . Therefore G is LO iff G is RO. We will stick to LO's.

- (3) For $H < G$, an LO (resp. BO) on G induces an LO (resp. BO) on H .

EXAMPLE 1.1. $(\mathbb{R}, +)$ with the usual $<$ is BO. The subgroups $\mathbb{Z} < \mathbb{Q} < \mathbb{R}$ are also BO.

Lemma 1.1. *Let $<$ be an LO on G . Then*

- (1) $g > 1, h > 1$ implies $gh > 1$;
- (2) $g > 1$ implies $g^{-1} < 1$;
- (3) $<$ is a BO iff $(g < h \implies f^{-1}gf < f^{-1}hf \forall f \in G)$ (i.e. $<$ is conjugation invariant).

PROOF. (1) $h > 1$ implies $gh > g \cdot 1g > 1$.

(2) $g > 1$ implies $g^{-1}g > g^{-1}$ implies $1 > g^{-1}$.

(3) (\implies) is immediate. (\impliedby) : We need to show $<$ is a RO. $g < h$ implies $fg < fh$ implies $f^{-1}(fg)f < f^{-1}(fh)f$ which implies $gf < hf$ as desired. □

Lemma 1.2. *If $<$ is a BO on G , then*

- (1) $g < h$ implies $g^{-1} > h^{-1}$;
- (2) $g_1 < h, g_2 < h_2$ implies $g_1g_2 < h_1h_2$.

PROOF. (1) If $g < h$, then $g^{-1}g < g^{-1}h$, which implies $1 < g^{-1}h$, which implies $1 \cdot h^{-1} < g^{-1}$, which implies $h^{-1} < g^{-1}$.

(2) $g_2 < h_2$ implies $g_1g_2 < g_1h_2 < h_1h_2$. □

WARNING 1.1. These don't necessarily true for LO's.

Lemma 1.3. *If G is LO then it is torsion free.*

PROOF. Consider $g \in G \setminus \{1\}$. If $g > 1$, then $g^2 > g > 1$, and similarly for all $n \geq 1$, $g^n > 1$. Similarly $g < 1$ implies $g^n < 1$ for all $n \geq 1$. \square

So LO is not preserved under taking quotients (e.g. $\mathbb{Z} \rightarrow \mathbb{Z}/n$).

Consider an indexed family of groups $\{G_\lambda \mid \lambda \in \Lambda\}$. Recall that the direct product

$$(1.2) \quad \prod_{\lambda \in \Lambda} G_\lambda = \{(g_\lambda)_{\lambda \in \Lambda}\}$$

with multiplication defined co-ordinatewise.

Recall a *well-order* (WO) on a set X is a STO \prec on X such that if $A \subset X$ and $A \neq \emptyset$ then there exists $a_0 \in A$ such that $a_0 \prec a$ for all $a \in A \setminus \{a_0\}$. Recall that the axiom of choice is equivalent to every set having a WO.

THEOREM 1.4. G_λ has a LO (resp. BO) for all $\lambda \in \Lambda$ iff $\prod_{\lambda \in \Lambda} G_\lambda$ has a LO (resp. BO).

PROOF. (\Leftarrow): $G_\lambda < \prod_{\lambda} G_\lambda$ so we are finished.

(\Rightarrow): Choose a WO \prec on Λ , and order $\prod_{\lambda} G_\lambda$ lexicographically. Let $g = (g_\lambda)$, $h = (h_\lambda)$, $g \neq h$. Then λ_0 be the \prec -least element of Λ such that $g_{\lambda_0} \neq h_{\lambda_0}$. Then define $g < h$ iff $g_{\lambda_0} < h_{\lambda_0}$ (in G_{λ_0}). Then $<$ is an LO (resp. BO) on $\prod_{\lambda} G_\lambda$. Left (resp. left and right) invariance is clear. Now we show transitivity. Suppose $f < g$, $g < h$. Let λ_0 be the \prec -least element of Λ such that $f_{\lambda_0} \neq g_{\lambda_0}$. Let μ_0 be the \prec -least element of Λ such that $g_{\mu_0} \neq h_{\mu_0}$.

- (1) ($\lambda_0 \preccurlyeq \mu_0$): Then $f_\lambda = g_\lambda = h_\lambda$ for all $\lambda \prec \lambda_0$. Then g_{λ_0} is $<$ (resp. $=$) h_{λ_0} if $\lambda_0 = \mu_0$ (resp. $\lambda_0 \prec \mu_0$). So $f_{\lambda_0} < g_{\lambda_0} \leq h_{\lambda_0}$, and therefore $f_{\lambda_0} < h_{\lambda_0}$.
- (2) ($\mu_0 < \lambda_0$): This follows similarly.

\square

Let $\sum_{\lambda \in \Lambda} G_\lambda$ be the *direct sum* of $\{G_\lambda\}$. Recall this is the subgroup of $\prod_{\lambda \in \Lambda} G_\lambda$ consisting of elements such that all but finitely many co-ordinates are 1.

Corollary 1.5. G_λ is LO (resp. BO) for all $\lambda \in \Lambda$ iff $\sum_{\lambda \in \Lambda} G_\lambda$ is LO (resp. BO).

Corollary 1.6. Free abelian groups are BO.

PROOF. Free abelian groups on Λ are $\sum_{\lambda \in \Lambda} \mathbb{Z}$. \square

Let $<$ be an LO on G . The *positive cone* $P = P_{<}$ of $<$ is $\{g \in G \mid g > 1\}$.

Lemma 1.7. Let P be as above.

- (1) $g, h \in P$, implies $gh \in P$ (i.e. $PP \subset P$).
- (2) $G = P \amalg P^{-1} \amalg \{1\}$.
- (3) $<$ is a BO on G iff $f^{-1}Pf \subset P$ for all $f \in G$.

PROOF. (1) This follows from Lemma 1.1 (1).

(2) This follows from Lemma 1.1 (2).

(3) This follows from Lemma 1.1 (3). \square

We say $P \subset G$ is a *positive cone* if P satisfies the conditions in Lemma 1.7.

Lemma 1.8. Let $P \subset G$ be a positive cone. Then $g < h$ implies $g^{-1}h \in P$ defines a LO $<$ on G (With $P_{<} = P$).

PROOF. $<$ is a STO, so:

- (i) $f < g, g < h$ implies $f^{-1}g \in P, g^{-1}h \in P$, which implies (by the first property) that $(f^{-1}g)(g^{-1}h) \in P$, which implies $f < h$.
- (ii) By the second property, for all $g, h \in G$ exactly one of the following holds: $g^{-1}h \in P, g^{-1}h \in P^{-1}$, and $g^{-1}h = 1$. Equivalently, $g < h, h < g$ (since $h^{-1}g \in P$), and $g = h$. Now we show left invariance. $g < h$ implies $g^{-1}h \in P$, but $g^{-1}h = (g^{-1}f^{-1})(fh)$ which implies $fg < fh$.

□

Lemmata 1.7 and 1.8 show that:

$$(1.3) \quad \{\text{LO's on } G\} \quad \leftrightarrow \quad \{\text{positive cones in } G\}$$

$$(1.4) \quad \{\text{BO's on } G\} \quad \leftrightarrow \quad \{\text{conjugacy-invariance positive cones in } G\} .$$

Consider the free group of rank n , F_n .

THEOREM 1.9. F_2 is LO.

PROOF BY SUSIC. Write $F_2 = F(a, b)$. $g \in F_2$ implies we can write it as a reduced word

$$(1.5) \quad (a^{m_1}) b^{n_1} \dots a^{m_k} (b^{n_k})$$

for $k \geq 0, m_i, n_i \in \mathbb{Z} \setminus \{0\}$. Recall 1 is the empty word, $k = 0$. Let $e(g)$ be the number of syllables in g with positive exponent, minus the number of syllables in g with negative exponent. Then define $j(g)$ so be the number of $a^m b^n$'s in f , minus the number of $b^n a^m$'s in G . So $j(g) = 0$, or ± 1 . For example:

$$(1.6) \quad j(a^* \dots a^*) = 0$$

$$(1.7) \quad j(b^* \dots b^*) = 0$$

$$(1.8) \quad j(a^* \dots b^*) = 1$$

$$(1.9) \quad j(b^* \dots a^*) = -1 .$$

Finally define

$$(1.10) \quad \tau(g) = e(g) + j(g) .$$

Note that

$$(1.11) \quad e(g^{-1}) = -e(g) \quad j(g^{-1}) = -j(g) .$$

Lemma 1.10. If $g \neq 1$, then $\tau(g) \equiv 1 \pmod{2}$.

PROOF. $e(f)$ is congruent to the number of syllables mod 2, and $j(g)$ is congruent to the number of syllables $+1 \pmod{2}$. □

Lemma 1.11. $|\tau(gh) - \tau(g) - \tau(h)| \leq 1$.

PROOF. If gh or g or $h = 1$ we are done. So suppose $gh, g, h \neq 1$. Clearly $e(gh) = e(g) + e(h) + \begin{Bmatrix} 0 \\ 1 \\ -1 \end{Bmatrix}$. Similarly:

$$(1.12) \quad j(gh) = j(g) + j(h) + \begin{Bmatrix} 0 \\ 1 \\ -1 \end{Bmatrix} .$$

Therefore:

$$(1.13) \quad |\tau(gh) - \tau(g) - \tau(h)| \leq 2$$

so by Lemma 1.10 we have

$$(1.14) \quad |\tau(gh) - \tau(g) - \tau(h)| \leq 1.$$

□

REMARK 1.2. Lemma 1.11 says that $\tau : F_2 \rightarrow \mathbb{Z}(< \mathbb{R})$ is what is called a *quasi-morphism*.

Define $P \subset F_2$ by

$$(1.15) \quad P = \{g \in F_2 \mid \tau(g) > 0\}.$$

Then $F_2 = P \amalg P^{-1} \amalg \{1\}$ by Lemma 1.10 and that $\tau(g^{-1}) = -\tau(g)$. Then $PP \subset P$ by Lemma 1.11 since

$$(1.16) \quad \tau(gh) \geq \tau(g) + \tau(h) - 1 \geq 1.$$

Therefore P is a positive cone for a LO on F_2 . ■

Corollary 1.12. *Any countable free group is LO.*

PROOF. A countable free group is a subgroup of F_2 . □

REMARK 1.3. (1) $\tau(a^{-1}b) = 1$, so $a^{-1}b > 1$, so $b > a$. On the other hand, $\tau(ab^{-1}) = 1$, so $ab^{-1} > 1$, so $b^{-1} > a^{-1}$. So τ does not define a BO on F_2 .

(2) We will see later that all free groups are LO.

(3) Even later we will see that all free groups are BO.

THEOREM 1.13. *Let $1 \rightarrow H \rightarrow G \rightarrow Q \rightarrow 1$ be a short-exact sequence of groups. Then*

(1) *H, Q LO implies G is LO;*

(2) *if Q is BO and H has a BO that is invariant under conjugation in G then G is BO.*

PROOF. Write $\varphi : G \rightarrow Q$ and regard H as $\ker \varphi < G$. Let P_H (resp. P_Q) be positive cones for LO's on H (resp. Q). Define $P = \varphi^{-1}(P_Q) \amalg P_H$.

CLAIM 1.1. P is a positive cone for an LO on G .

PROOF. We need to check (1) and (2) from Lemma 1.7. Let $g, h \in P$. Then we want to show $gh \in P$. We have three cases.

(a) $g, h \in \varphi^{-1}(P_Q)$: In this case $\varphi(g), \varphi(h) \in P_Q$, so $\varphi(gh) = \varphi(g)\varphi(h) \in P_Q$. Therefore $gh \in \varphi^{-1}(P_Q)$.

(b) $g, h \in P_H$: In this case $gh \in P_H$.

(c) $g \in \varphi^{-1}(P_Q), h \in P_H$: Then $\varphi(gh) = \varphi(g) \in P_Q$, so $gh \in \varphi^{-1}(P_Q)$. Similarly $hg \in \varphi^{-1}(P_Q)$.

Now we need to check $P \amalg P^{-1} \amalg \{1\}$. But this follows from the fact that:

$$(1.17) \quad G = (H \setminus \{1\}) \amalg \varphi^{-1}(Q \setminus \{1\}) \amalg \{1\} = \varphi^{-1}(P_Q) \amalg \varphi^{-1}(P_Q^{-1})$$

since $H \setminus \{1\} = P_H \amalg P_H^{-1}$. □

We leave (2) as an exercise. [Hint: Recall P is a positive cone for BO on G iff it is a conjugacy invariant cone for an LO.] ■

1. Orderability of manifold groups

EXAMPLE 1.2. Let X^2 be the Klein bottle. This has fundamental group

$$(1.18) \quad K = \pi_1(X^2) = \langle a, b \mid b^{-1}ab = a^{-1} \rangle .$$

This fits in the SES:

$$(1.19) \quad \begin{array}{ccccccc} 1 & \longrightarrow & \mathbb{Z} & \longrightarrow & K & \longrightarrow & \mathbb{Z} \longrightarrow 1 \\ & & \parallel & & & & \\ & & \langle a \rangle & & b \longmapsto gm & & \end{array}$$

which means K is LO by Theorem 1.13.

Note that K is *not* BO. We have that $a > 1$ iff $b^{-1}ab > 1$, but this is a^{-1} , so $a^{-1} > 1$ which is a contradiction.

Notice that \mathbb{Z} has exactly two LO's. The usual one, and the opposite. Therefore, if we choose an LO on $\langle a \rangle$ and $K/\langle a \rangle$, this gives 4 LO's on K determined by:

- (i) $a > 1, b > 1$;
- (ii) $a > 1, b < 1$;
- (iii) $a < 1, b > 1$;
- (iv) $a < 1, b < 1$.

THEOREM 1.14. *These are the only LO's on K .*

PROOF. It suffices to show that each of these conditions determines a unique positive cone.

- (i) $a > 1, b > 1$:

CLAIM 1.2. $a^k < b$ for all $k \in \mathbb{Z}$.

PROOF. $b < a^k$ implies $a^{-k}b < 1$. But $a^{-k}b = ba^k$ and $b > 1$, so $b < a^k$ implies $a^k > 1$, which implies $ba^k > 1$ which is a contradiction. \square

Note that every element in K has a unique representative of the form $a^m b^n$ for $m, n \in \mathbb{Z}$.

CLAIM 1.3. $a^m b^n > 1$ iff either $n > 0$ or $n = 0$ and $m > 0$.

PROOF. If $n = 0$, then this is clear. If $n > 0$, then $a^m b > 1$ for any m by claim 1 (for $k = -m$). But we also know $b > 1$ which implies $b^n > 1$, so we get $a^m b^n > 1$ for $n > 0$. On the other hand, if $m < 0$ then $a^m b^n = b^n a^{\pm m} = (a^{\mp m} b^{-n})^{-1}$. Then we know $a^{\mp m} b^{-n} > 1$ by the case above, so its inverse is < 1 . \square

If $<$ is an LO on G , and $\alpha : G \rightarrow G$ is an automorphism, then this induces an LO $<_\alpha$ on G given by: $g <_\alpha h$ iff $\alpha(g) < \alpha(h)$. Now notice that there are automorphisms α_1, α_2 of K such that

$$(1.20) \quad \alpha_1(a) = a, \quad \alpha_1(b) = b^{-1}$$

$$(1.21) \quad \alpha_1(a) = a^{-1}, \quad \alpha_1(b) = b.$$

In particular, α_1 is given by

$$(1.22) \quad \langle a, b \mid b^{-1}ab = a^{-1} \rangle \cong \langle a, b \mid bab^{-1} = a^{-1} \rangle$$

and similarly for α_2 .

Write $<_{(i)}$ for the unique LO on K determined by (i). Then $<_{(ii)}$ is induced by $<_{(i)}$ and α_1 , $<_{(iii)}$ is induced by $<_{(i)}$ and α_2 , and $<_{(iv)}$ is induced by $<_{(i)}$ and $\alpha_1 \alpha_2$.

■

FACT 1. *If G has only finitely many LO's, then the number of LO's is of the form 2^n .*

EXERCISE 1.1. Show that for all $n \geq 0$ there exists a group G with exactly 2^n LO's.

Corollary 1.15. *For any LO on K , if $h \in \langle a \rangle$, $g \in K \setminus \langle a \rangle$, and $g > 1$, then $g > h$.*

PROOF. It is sufficient to check this for the first LO, since the other three are determined by the above automorphisms. Let $a > 1$, $b > 1$. By claim 2 from above, we know $g = a^m b^n$ for $n > 0$. We now there is some k such that $h = a^k$, and therefore

$$(1.23) \quad h^{-1}g = a^{m-k}b^n > 1$$

by claim 2, so $g > h$. □

2. Three-manifold groups

Suppose M is a closed, orientable, connected three-manifold. Then we might ask if $\pi_1(M)$ is LO? BO?

Immediately we notice that not all such groups are. If M is a lens space, then $\pi_1(M) \cong \mathbb{Z}/n$ for $n > 1$, so this is not LO. More generally, for $\pi_1(M)$ nontrivial and finite is not LO. Recall that if $M = M_1 \# M_2$, then this implies $\pi_1(M) \cong \pi_1(M_1) * \pi_1(M_2)$. So, for example, if $M_1 \#$ lens space, then $\pi_1(M)$ has torsion, so not LO.

But at least some of them are. Consider $M \cong T^3 = S^1 \times S^1 \times S^1$. Then $\pi_1(M) = \mathbb{Z}^3$ is of course LO. Similarly $M = \#_n (S^1 \times S^2) \cong F_n$, so $\pi_1(M)$ is LO.

We will show that there exist (three-manifold) groups that are torsion-free, but not LO.

Let $p : T^2 \rightarrow X^2$ be a two-fold covering of the Klein bottle. Recall that

$$(1.24) \quad K > p_* (\pi_1(T^2)) = \langle a, b^2 \rangle \cong \mathbb{Z} \times \mathbb{Z}.$$

Let N be the mapping cylinder of p , namely:

$$(1.25) \quad N = (T^2 \times I) \amalg X^2 / ((x, 0) \sim p(x) \forall x \in T^2).$$

The orientation reversing curve representing b doesn't lift. So N is orientable. Note that $\partial N \cong T^2$. There is a strong deformation retraction $N \rightarrow X^2$, so $\pi_1(N) \cong K$. Let N_1, N_2 be two copies of N . Write

$$(1.26) \quad \pi_1(N_i) = \langle a_i, b_i \mid b_i^{-1} a_i b_i = a_i^{-1} \rangle.$$

Notice that $\pi_1(\partial N_i) \cong \mathbb{Z} \times \mathbb{Z} = \langle a_i, b_i^2 \rangle < \pi_1(N_i)$. Let $\varphi : \partial N_1 \rightarrow \partial N_2$ be a homeomorphism. Let $M_\varphi = N_1 \cup_\varphi N_2$. This is a closed, orientable three-manifold. Therefore

$$(1.27) \quad \pi_1(M_\varphi) = \pi_1(N_1) *_{\mathbb{Z} \times \mathbb{Z}} \pi_1(N_2) \cong K_1 *_{\mathbb{Z} \times \mathbb{Z}} K_2.$$

Since K is torsion-free, $\pi_1(M_\varphi)$ is torsion-free. But in fact we have the following theorem.

THEOREM 1.16. *If $H_1(M_\varphi)$ is finite, then $\pi_1(M_\varphi)$ is not LO.*

REMARK 1.4. We will see later that for M a prime three-manifold with $H_1(M)$ infinite has $\pi_1(M)$ LO.

PROOF. φ is determined up to isotopy, so the resulting manifold M_φ depends only on $\varphi_* : H_1(\partial N_1) \rightarrow H_1(\partial N_2)$. We know

$$(1.28) \quad \mathbb{Z} \oplus \mathbb{Z} = \mathbb{Z} \langle a_1, 2b_1 \rangle \quad \mathbb{Z} \oplus \mathbb{Z} = \mathbb{Z} \langle a_2, 2b_2 \rangle$$

so φ_* is given by some 2×2 matrix with \mathbb{Z} coefficients

$$(1.29) \quad \begin{bmatrix} p & r \\ q & s \end{bmatrix}$$

with determinant $ps - qr = \pm 1$. Specifically we have:

$$(1.30) \quad \varphi_*(a_1) = pa_2 + 2qb_2$$

$$(1.31) \quad \varphi_*(2b_1) = ra_2 + 2sb_2 .$$

Now we have $H_1(N_i) = \mathbb{Z} \oplus \mathbb{Z}_2$ with basis b_i and a_i respectively. Then $H_q(M_\varphi)$ is presented by

$$(1.32) \quad A = \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ -1 & 0 & p & 2q \\ 0 & -2 & r & 2s \end{bmatrix} .$$

where we order the basis as $\{a_1, b_1, a_2, b_2\}$. Interchanging columns 2 and 3 we get

$$(1.33) \quad \det A = 4 \left| \det \begin{bmatrix} 0 & 2q \\ -2 & 2s \end{bmatrix} \right| = 16 |q| .$$

Therefore $H_1(M_\varphi)$ is finite iff $q \neq 0$ iff $\varphi_*(a_1) \neq \pm a_2$.

Suppose $\pi_1(M_\varphi)$ is LO. Then we would get an induced LO on the common boundary $\partial N_1 = \partial N_2$. But there are only 4 LO's on $\pi_1(N_i)$ (for $i \in \{1, 2\}$). By Corollary 1.15, for any LO on $\pi_1(N)$, $\langle a \rangle$ is the unique \mathbb{Z} -summand of $\pi_1(\partial N) = \langle a, b^2 \rangle$ such that if $h \in \langle a \rangle$ and $g \in \pi_1(\partial N) \setminus \{1\}$, $g > 1$, then $g > h$. Therefore $\varphi_*(a_1) = \pm a_2$ which is a contradiction. \square

Let $<$ be an STO on a set X . Let $\mathcal{B}(X, <)$ be the group of $<$ -preserving bijections $X \rightarrow X$.

Lecture 3; January 28, 2020

THEOREM 1.17. $\mathcal{B}(X, <)$ is always LO.

PROOF. Let \prec be a WO on X . Let $f, g \in \mathcal{B}(X, <)$ such that $f \neq g$. Write

$$(1.34) \quad [f \neq g] = \{x \in X \mid f(x) \neq g(x)\} \neq \emptyset .$$

Let x_0 be the \prec -least element of $[f \neq g]$. Define

$$(1.35) \quad f < g \iff f(x_0) < g(x_0) .$$

Then we claim that this is an LO on $\mathcal{B}(X, <)$. Left-invariance is clear. To see this is a STO we need “trichotomy” and transitivity. Trichotomy is easy, and transitivity follows from the same argument as the proof of Theorem 1.4. \square

EXAMPLE 1.3. Let $<$ be the standard order on \mathbb{R} . Then $\mathcal{B}(\mathbb{R}, <)$ consists of the orientation-preserving homeomorphisms $\mathbb{R} \rightarrow \mathbb{R}$, written $\text{Homeo}_+(\mathbb{R})$.

Corollary 1.18. $\text{Homeo}_+(\mathbb{R})$ is LO.

REMARK 1.5. For $x \in \mathbb{R}$, let \prec_x be a WO on \mathbb{R} such that x is the \prec_x -least element of \mathbb{R} . Let $<_x$ be the LO on $\text{Homeo}_+(\mathbb{R})$ induced by \prec_x , as in the proof of Theorem 1.17. Given $x \neq y \in \mathbb{R}$, there exists $g \in \text{Homeo}_+(\mathbb{R})$ such that $g(x) > x$ and $g(y) < y$. But this means

$$g <_x 1 \qquad g <_y 1 .$$

which implies $<_x \neq <_y$. Therefore $\text{Homeo}_+(\mathbb{R})$ has uncountably many LO's.

REMARK 1.6. It is a fact that the number of LO's on a group G is either finite (and of the form 2^n) or uncountable.

Corollary 1.19. *A group G is LO iff G acts faithfully^{1.1} on a STO'd set $(X, <)$.*

PROOF. (\Leftarrow): This follows from Theorem 1.17.

(\Rightarrow): G acts faithfully on $(G, <)$ by left multiplication. \square

Corollary 1.18 implies that any subgroup of $\text{Homeo}_+(\mathbb{R})$ is LO. E.g. one can show that F_2 (the free group of rank 2) is a subgroup of $\text{Homeo}_+(\mathbb{R})$. (This is another way to show that countable free groups are LO.) In fact this characterizes countable LO groups.

THEOREM 1.20. *Let G be a countable group. Then G is LO iff there exists an injective homomorphism $G \rightarrow \text{Homeo}_+(\mathbb{R})$.*

PROOF. (\Leftarrow): This follows from Corollary 1.18.

(\Rightarrow): We actually prove something slightly stronger. This will follow from Theorem 1.21. \square

THEOREM 1.21. *Let $(G, <)$ be a countable group with an LO. Then there exists a LO on $\text{Homeo}_+(\mathbb{R})$ and an order-preserving injective homomorphism $(G, <) \rightarrow (\text{Homeo}_+(\mathbb{R}), <)$.*

SKETCH OF PROOF. Let $<$ be an LO on G . If $G = \{1\}$ this is immediate, so assume $G \neq \{1\}$. Therefore it is infinite, since LO groups are torsion free. Let g_1, g_2, \dots be some enumeration of the elements of G .

Define an embedding $e : G \rightarrow \mathbb{R}$ by $e(g_1) = 0$, and inductively by:

(i) If $g_{n+1} \begin{Bmatrix} > \\ < \end{Bmatrix} g_i$ for all $1 \leq i \leq n$, then set

$$(1.36) \quad e(g_{n+1}) = \begin{Bmatrix} \max \{e(g_i) \mid 1 \leq i \leq n\} + 1 \\ \min \{e(g_i) \mid 1 \leq i \leq n\} - 1 \end{Bmatrix}.$$

(ii) Otherwise let

$$g_l = \max \{g_i \mid 1 \leq i \leq n, g_i < g_{n+1}\}$$

$$g_r = \min \{g_i \mid 1 \leq i \leq n, g_i > g_{n+1}\}$$

and set

$$e(g_{n+1}) = \frac{e(g_l) + e(g_r)}{2}.$$

REMARK 1.7. (1) e is order-preserving, i.e. $a < b \implies e(a) < e(b)$.

(2) $e(g_{n+1}) \in \mathbb{Z}$ iff (i) holds.

(3) If $g > 1$ then $g^2 > g$ and $g^{-1} < g$. If $g < 1$ then $g^2 < g$ and $g^{-1} > g$, which implies $\mathbb{Z} \subset e(G) = \Gamma$.

(4) G acts on Γ by $g(e(a)) = e(ga)$. In fact, G acts on $(\Gamma, <)$ (where $<$ is the restriction of $<$ on \mathbb{R}) since $e(a) < e(b)$ iff $a < b$ iff $ga < gb$ iff $e(ga) < e(gb)$ iff $g(e(a)) < g(e(b))$.

To see that this action extends to an action of G on \mathbb{R} , we have a few steps.

Step 1: The action of G on Γ is continuous,

Step 2: The action of G on Γ extends to a continuous action of G on $\bar{\Gamma}$.

^{1.1}Recall this means $g(x) = x$ for all $x \in X$ iff $g = 1$.

Step 3: $\mathbb{R} \setminus \bar{\Gamma}$ is a countable Π of open intervals (a_i, b_i) ; the action of G is defined on $\{a_i, b_i\}$; and extends to $[a_i, b_i]$.

Note, to ensure Step 1:, it is not enough to take e to be an order-preserving of G in \mathbb{R} . It must be continuous.

To define an LO on $\text{Homeo}_+(\mathbb{R})$ that restricts to the LO on Γ from G , first pick any $\gamma \in \Gamma$. Then $g > 1$ (resp. < 1) iff $g(\gamma) > \gamma$ (resp. $< \gamma$). Let \prec be a WO on \mathbb{R} such that γ is the \prec -least element of \mathbb{R} . Then let \leq be the LO on $\text{Homeo}_+(\mathbb{R})$ induced by \prec . Then $g > 1$ (resp. < 1) in G iff $g \succ 1$ (resp. \prec) in $\text{Homeo}_+(\mathbb{R})$. \square

3. Group rings

Let R be a ring (with 1).

- $a \in R$ is a *unit* if there exists $b \in R$ such that $ab = ba = 1$.
- $a \in R$ is a *zero-divisor* if $a \neq 0$ and there exists $b \neq 0$ such that either $ab = 0$ or $ba = 0$.
- $a \in R$ is a *non-trivial idempotent* if $a^2 = a$ but $a \neq 0$ and $a \neq 1$.

Let G be a group and R a ring. Then the R -group ring of G consists of formal sums:

$$(1.37) \quad RG := \left\{ \sum r_g g \mid g \in G, r_g \in R, r_g \neq 0 \forall \text{ but f'tly many } g \in G \right\}.$$

RG is a ring with respect to the obvious operations. For $g \in G$ and $r \in R$ a unit, then rg is a unit in RG . A unit in RG is *non-trivial* if it is not of this form.

REMARK 1.8. If $\tilde{X} \rightarrow X$ is a universal covering, then $\pi = \pi_1(X)$ acts on \tilde{X} so $H_*(\tilde{X}, \mathbb{Z})$ is a $\mathbb{Z}\pi$ -module.

THEOREM 1.22. Suppose G has non-trivial torsion, and K is a field of characteristic 0.

- (1) KG has zero divisors,
- (2) KG has non-trivial units,
- (3) KG has non-trivial idempotents.

PROOF. Let $g \in G$ have order $n \geq 2$. Define

$$\sigma = 1 + g + g^2 + \dots + g^{n-1} \in KG.$$

First notice that

$$(1.38) \quad g\sigma = \sigma$$

which implies $(1 - g)\sigma = 0$ so we have zero divisors.

(1.38) also gives us that $\sigma^2 = n\sigma$. Therefore

$$(1 - \sigma) \left(1 - \frac{1}{n-1} \sigma \right) = 1$$

so we have a nontrivial unit for $n > 2$. If $n = 2$, $1 - \sigma = -g$, but we still have:

$$(1.39) \quad (1 - 2\sigma) \left(1 - \frac{2}{3} \sigma \right) = 1.$$

Finally, we have that

$$(1.40) \quad \left(\frac{1}{n} \sigma \right)^2 = \left(\frac{1}{n^2} \right) \sigma^2 = \frac{1}{n} \sigma$$

so we have nontrivial idempotents. \square

Note that the proof of (1) works even for $\mathbb{Z}G$.

REMARK 1.9. If $n \notin \{2, 3, 4, 6\}$ then $\mathbb{Z}G$ has nontrivial units. This is a theorem of Higman.

EXAMPLE 1.4. For $n = 5$,

$$(1.41) \quad (1 - g - g^4)(1 - g^2 - g^3) = 1 .$$

But what if G is torsion free? This brings us to the famous Kaplansky conjectures.

CONJECTURE 1 (Kaplansky). *If G is torsion free and K is a field, then:*

I (Units conjecture): KG has no non-trivial units,

II (Zero-divisors conjecture): KG has no zero divisors,

III (Idempotents conjecture): KG has no non-trivial idempotents.

REMARK 1.10. Clearly II implies III since $a^2 = a$ implies $a(a - 1) = 0$, which by II implies $a = 0$ or $a = 1$ which implies III. In fact they're all equivalent, but this is nontrivial to see.

Lecture 4; January
30, 2020

REMARK 1.11. Note that if R is an integral domain (e.g. \mathbb{Z}) then R is contained in its field of fractions. In this case items I and II and item III for its field of fractions imply the corresponding versions of items I and II and item III for R .

REMARK 1.12. We know this is true for LO groups. As we have seen, we should think of LO as being a stronger version of torsion free.

THEOREM 1.23. *If G is LO then KG satisfies items I and II and item III.*

PROOF. Since item I implies item III by the above remark we show item I and item II.
item I: Suppose

$$(1.42) \quad \left(\sum_{i=1}^m \alpha_i g_i \right) \left(\sum_{j=1}^n \beta_j h_j \right) = 1$$

with m, n not both 1, $\alpha_i, \beta_j \neq 0 \in K$, distinct $g_i \in G$, and distinct $h_i \in G$. Note this product can be rewritten as the following sum with mn terms:

$$(1.43) \quad \sum_{i,j} (\alpha_i \beta_j) (g_i h_j) .$$

Assume WLOG that $h_1 < h_2 < \dots < h_n$. Let $g_k h_l$ be a minimal element of

$$(1.44) \quad S = \{g_i h_j \mid 1 \leq i \leq m, 1 \leq j \leq n\} \subset G .$$

We know $h_1 < h_j$ for $j > 1$, so $g_k h_1 < g_k h_j$ for all $j > 1$. Therefore $l = 1$. Also $gh_1 = g'h_1$ which implies $g = g'$. Therefore $g_k h_1$ is the unique

$$(1.45) \quad (k, 1) \in \{(i, j) \mid 1 \leq i \leq m, 1 \leq j \leq n\}$$

such that $g_k h_1$ is a minimal element of S .

Similarly, there is a unique

$$(1.46) \quad (r, n) \in \{(i, j) \mid 1 \leq i \leq m, 1 \leq j \leq n\}$$

such that $g_r h_n$ is a maximal element of S .

CLAIM 1.4. $g_k h_1 \neq g_r h_n$.

If they were equal, then $r = k$, $n = 1$, so $m > 1$. So $g_k h_1 = g_r h_1$, and therefore $g_r = g_k$. But this cannot be the case since they are distinct by assumption.

This implies that (1.43) has ≥ 2 terms after cancellation, so it cannot be 1.

item II: Now suppose

$$(1.47) \quad \left(\sum_{i=1}^m \alpha_i g_i \right) \left(\sum_{j=1}^n \beta_j h_j \right) = 0$$

for $m, n \geq 1$. Then there is a unique minimal element and nonzero coefficient, which means it is nonzero. \square

CONJECTURE 2 (Isomorphism conjecture). *If G is torsion free, then $\mathbb{Z}G \cong \mathbb{Z}H$ implies $G \cong H$.*

REMARK 1.13. In [5] a finite counterexample to the conjecture for arbitrary groups was provided, i.e. it is shown that there exists finite G, H such that $\mathbb{Z}G \cong \mathbb{Z}H$, $G \not\cong H$.

Corollary 1.24 ([8]). *If G is LO, then G satisfies the isomorphism conjecture.*

PROOF. Theorem 1.23 implies that $\mathbb{Z}G$ has no nontrivial units. Call $\mathcal{U}_{\mathbb{Z}G}$ the group of units in $\mathbb{Z}G = \mathbb{Z}/2 \times G$. Suppose $\mathbb{Z}G \cong \mathbb{Z}H$. Theorem 1.23 says that $\mathbb{Z}G$ has no 0-divisors. This implies $\mathbb{Z}H$ has no 0-divisors, which means (by Theorem 1.22) that H is torsion-free. Now $H < \mathcal{U}_{\mathbb{Z}H} \cong \mathcal{U}_{\mathbb{Z}G} \cong \mathbb{Z}/2 \times G$ which implies $H < G$ (since H is torsion-free), which implies H is LO (since G is), which implies $\mathcal{U}_{\mathbb{Z}H} \cong \mathbb{Z}/2 \times H$, which implies $\mathbb{Z}/2 \times H \cong \mathbb{Z}/2 \times G$ which implies $H \cong G$ (since H, G are torsion free). \square

REMARK 1.14. We might wonder if it is ever the case that (for $G \neq 1$) $(G * \mathbb{Z}) / \langle \langle w \rangle \rangle = 1$? This is known for G torsion free [6].

COUNTEREXAMPLE 1. If we consider the question of whether we can ever have $(A * B) / \langle \langle w \rangle \rangle = 1$ for A, B nontrivial, a counterexample is given by:

$$\mathbb{Z}/2 * \mathbb{Z}/3 / (a = b) .$$

4. BO's on $\mathbb{Z} \times \mathbb{Z}$

Recall we have 2 orders on \mathbb{Z} . Consider a line of slope α in $\mathbb{Z} \times \mathbb{Z}$. Then we have two cases.

- (1) α irrational: The associated positive cone is everything above the line. Specifically, $P \subset \mathbb{Z} \times \mathbb{Z}$ is given by

$$(1.48) \quad P = \{(m, n) \mid n > m\alpha\} .$$

It is easy to check that this is a positive cone. This means there are uncountable many BO's on $\mathbb{Z} \times \mathbb{Z}$.

- (2) α rational: Notice that now

$$(1.49) \quad \{(m, n) \mid n = m\alpha\} \cong \mathbb{Z} < \mathbb{Z} \times \mathbb{Z} .$$

Now let P_0 be one of the two positive cones on \mathbb{Z} . Then we can check that

$$P = P_0 \amalg \{(m, n) \mid n > m\alpha\}$$

is a positive cone for $\mathbb{Z} \times \mathbb{Z}$.

REMARK 1.15. (1) (Up to reversal) these are all the BOs on $\mathbb{Z} \times \mathbb{Z}$. I.e. for α rational we get two, and for α irrational we get 4.

- (2) This generalizes in the obvious way to \mathbb{Z}^n .

5. BO's on \mathbb{R}

Regard \mathbb{R} as a vector space on \mathbb{Q} with uncountable bases Λ . Recall Λ exists by the axiom of choice. Therefore $\mathbb{R} \subset \mathbb{Q}^\Lambda$. In particular it is the elements of \mathbb{Q}^Λ with only finitely many non-zero coordinates. There are uncountable many WO's on Λ , and each gives rise to a lexicographic BO on \mathbb{Q}^Λ . This gives us uncountably many BOs on \mathbb{R} .

CHAPTER 2

The space of left-orders on a group

The basic idea is that since lefts orders are determined by positive cones, we can give this space a topology. Consider a family of sets $\{X_\lambda \mid \lambda \in \Lambda\}$. Then write

$$X = \prod_{\lambda \in \Lambda} X_\lambda$$

and $\pi_\lambda : X \rightarrow X_\lambda$ for the projection. If X_λ is a topological space, then X can be given the product topology. This is the largest topology on X such that π_λ is continuous for all $\lambda \in \Lambda$. So X has subbasis

$$(2.1) \quad \left\{ \pi_\lambda^{-1}(U_\lambda) = U_\lambda \times \prod_{\mu \neq \lambda} X_\mu \mid U_\lambda \subset X_\lambda \text{ open, } \lambda \in \Lambda \right\}.$$

THEOREM. *If X_λ is compact for all $\lambda \in \Lambda$ then $\prod_{\lambda \in \Lambda} X_\lambda$ is compact.*

REMARK 2.1 (Exercises). (1) X_λ Hausdorff (for all $\lambda \in \Lambda$) implies $\prod_{\lambda \in \Lambda} X_\lambda$ is Hausdorff.

(2) A space X is totally disconnected if the only nonempty connected subspaces are singletons $\{x\}$ for $x \in X$. This is equivalent to the connected components of X all being $\{x\}$. Show that X_λ totally disconnected (for all $\lambda \in \Lambda$) implies $\prod_{\lambda \in \Lambda} X_\lambda$ is totally disconnected.

Let X be a set, let $\mathcal{S}(X)$ be the set of subsets of X (i.e. the power set). Then we have a correspondence:

$$\mathcal{S}(X) \quad \leftrightarrow \quad \{f : X \rightarrow \{0, 1\}\}$$

which sends:

$$A \subset X \quad \leftrightarrow \quad f_A : X \rightarrow \{0, 1\}$$

where

$$f_A(x) = \begin{cases} 1 & x \in A \\ 0 & x \notin A \end{cases}.$$

Give $\{0, 1\}$ the discrete topology, and give

$$\mathcal{S}(X) = \{0, 1\}^X = 2^X = \prod_{x \in X} \{0, 1\}$$

the product topology. Note $\{0, 1\}$ is a compact, Hausdorff, totally-disconnected space, which means $\mathcal{S}(X)$ is too. For $x \in X$ let

$$\begin{aligned} U_x &= \pi_x^{-1}(1) = \{A \subset X \mid x \in A\} \\ V_x &= \pi_x^{-1}(0) = \{A \subset X \mid x \notin A\}. \end{aligned}$$

Note that $V_x = \mathcal{S}(X) \setminus U_x$ so U_x and V_x are open and closed. Then

$$(2.2) \quad \{U_x \mid x \in X\} \cup \{V_x \mid x \in X\}$$

is a subbasis for $\mathcal{S}(X)$.

Lecture 5; February
4, 2020

Lemma 2.1. *Suppose $B \subset X$. Then*

$$\{A \subset X \mid B \not\subset A\} \quad \{A \subset X \mid A \cap B \neq \emptyset\}$$

are open subsets of $\mathcal{S}(X)$.

PROOF.

$$\{A \subset X \mid B \not\subset A\} = \bigcup_{b \in B} \{A \subset X \mid b \notin A\} = \bigcup_{b \in B} V_b$$

so it is open. The argument for the other set is similar. \square

If G is a group, let

$$(2.3) \quad \text{LO}(G) = \{\text{positive cones } \subset G\} \subset \mathcal{S}(G)$$

and equip it with the subspace topology. We call this the *space of left-orders on G* .

EXAMPLE 2.1. $\text{LO}(\mathbb{Z}) = \text{pt} \amalg \text{pt}$. $\text{LO}(\mathbb{Z} \times \mathbb{Z})$ is the cantor set.

THEOREM 2.2. $\text{LO}(G)$ is closed in $\mathcal{S}(G)$ and hence compact.

PROOF. We show $\mathcal{S}(G) \setminus \text{LO}(G)$ is open. Suppose $A \in \mathcal{S}(G) \setminus \text{LO}(G)$, i.e. $A \subset G$ is not a positive cone. So either:

- (i) $\exists g, h \in A$ such that $gh \notin A$ or
- (ii) $\exists g \in G$ such that $g, g^{-1} \in A$ or
- (iii) $1 \in A$ or
- (iv) $\exists g, g \neq 1$ such that $g \notin A$ and $g^{-1} \notin A$.

Now the point is that these are open conditions since we can write them in terms of the U_x 's and V_x 's. In particular:

$$\begin{aligned} (i) &\iff A \in U_g \cap U_h \cap V_{gh} & (ii) &\iff A \in U_g \cap U_{g^{-1}} \\ (iii) &\iff A \in U_1 & (iv) &\iff A \in \bigcup_{g \neq 1} (V_g \cap V_{g^{-1}}) . \end{aligned}$$

Therefore $\text{LO}(G)$ is compact, Hausdorff, and totally disconnected. \square

Similarly one can define the space of biorders on G , $\text{BO}(G)$, to be the set of conjugation invariant positive cones in G .

EXERCISE 2.1. Show that $\text{BO}(G)$ is closed inside of $\text{LO}(G)$.

Therefore $\text{BO}(G)$ is compact, Hausdorff, and totally disconnected.

1. The cantor set

The cantor set $C \subset I \subset \mathbb{R}$ is defined as follows. First write

$$\begin{aligned} C_1 &= [0, 1/3] \cup [2/3, 1] \\ C_2 &= ([0, 1/9] \cup [2/9, 1/3]) \cup ([2/3, 7/9] \cup [8/9, 1]) \\ &\dots \end{aligned}$$

then define

$$(2.4) \quad C = \bigcap_{n=1}^{\infty} C_n .$$

The idea is that we keep removing the middle thirds.

C is uncountable, totally-disconnected, closed in I . Therefore it is also compact and Hausdorff. This is a very surprising example. We can easily write down something uncountable and totally-disconnected, such as the irrationals, but they do not form a compact set.

Any $x \in I$ has a ternary expansion:

$$x = 0.x_1x_2\dots = \sum_{n=1}^{\infty} \frac{x_n}{3^n}$$

which is unique up to:

$$\dots x_k 22 \dots = \dots (x_{k+1}) 00 \dots$$

Now notice

$$x_1 = 1 \quad \Longleftrightarrow \quad x \in (1/3, 2/3)$$

with the convention that

$$\frac{1}{3} = 0.022\dots$$

Similarly (with the same convention) we have

$$x_1 \neq 1, x_2 = 1 \quad \Longleftrightarrow \quad x \in (1/9, 2/9) \cup (7/9, 8/9)$$

and so on. Then

$$(2.5) \quad C = \{x \in I \mid x = 0.x_1x_2\dots \mid \forall n, x_n = 0 \text{ or } 2\} .$$

Now give $\{0, 2\}^{\mathbb{N}}$ the product topology.

EXERCISE 2.2. Show that the map sending

$$(2.6) \quad 0.x_1x_2\dots \mapsto (x_1, x_2, \dots)$$

defines a homeomorphism

$$(2.7) \quad C \xrightarrow{\cong} \{0, 2\}^{\mathbb{N}} .$$

Now recall that $\text{LO}(G)$ is compact in $\{0, 1\}^G$, so if G is countable, then $\text{LO}(G)$ is homeomorphic to a subspace of C .

We say $x \in X$ is *isolated* if $\{x\}$ is open. We say X is *perfect* if it has no isolated points. As it turns out, the Cantor set is perfect.

THEOREM. *If X is a compact, totally-disconnected, and perfect metric space, then $X \cong C$.*

Therefore, if G is countable, $\text{LO}(G) \neq \emptyset$, and has no isolated points, then $\text{LO}(G) \cong C$.

EXAMPLE 2.2. In 2004 [18] it was shown that if $n > 1$ then $\text{LO}(\mathbb{Z}^n) = \text{BO}(\mathbb{Z}^n) \cong C$.

EXAMPLE 2.3. In 1985 [10] it was shown that $\text{LO}(F_n) \cong C$. It is unknown if $\text{LO}(F_n)$ has isolated points.

REMARK 2.2. As it turns out, the braid group is LO. The first proof of this fact was not topological, so topologists started to think of a topological proof. When someone asked Thurston, he said “of course the braid group is left-orderable!”

If $X \subset G$, let $S(X)$ be the semigroup generated by X in G . This is the same as the non-empty product of elements in X . There is a characterization of left orderability in terms of finite subsets of G .

THEOREM 2.3. G is LO iff for all finite $F \subset G \setminus \{1\}$, there exists $\epsilon : F \rightarrow \{\pm 1\}$ such that

$$(2.8) \quad 1 \notin S\left(\left\{f^{\epsilon(f)} \mid f \in F\right\}\right) (= S(F, \epsilon)) .$$

REMARK 2.3. It follows from this that, given a solution to the word problem in G , there exists a machine such that if G is not LO, the machine will eventually tell you that. Nathan Dunfield has an explicit algorithm for three-manifold groups.

REMARK 2.4. If we take the n -fold cyclic branch cover of the knot 5_2 , then we can consider $\pi_1(\Sigma_n(5_2))$. For $n = 2$, this is a lens space so π_1 is finite. It is also not LO for $n = 3, 4$, and 5 . But it is unknown for $n = 6, 7$, and 8 . (If the L -space conjecture is true,^{2.1} then it should be LO for these values of n .) For $n \geq 9$ it is known to be LO.

PROOF. (\implies): Define

$$\epsilon(f) = \begin{cases} +1 & f > 1 \\ -1 & f < 1 \end{cases} .$$

(\impliedby): Let $F \subset G \setminus \{1\}$ be finite, $\epsilon : F \rightarrow \{\pm 1\}$. Define

$$Q(F, \epsilon) := \left\{ Q \subset G \setminus \{1\} \mid S(F, \epsilon) \subset Q, S(F, \epsilon)^{-1} \cap Q = \emptyset \right\} .$$

Note that $Q(F, \epsilon) \neq \emptyset$ iff (2.8) holds. Let

$$Q(F) = \cup_{\epsilon} Q(F, \epsilon) .$$

Note this is a finite union.

CLAIM 2.1. $Q(F)$ is closed in $S(G)$.

PROOF. It is sufficient to show that $Q(F, \epsilon)$ is closed, i.e. $S(G) \setminus Q(F, \epsilon)$ is open. Suppose $A \subset G$, $A \not\subset Q(F, \epsilon)$ i.e. either $1 \in A$, or $S(F, \epsilon) \not\subset A$, or $S(F, \epsilon)^{-1} \cap A \neq \emptyset$. These conditions are all open by Lemma 2.1. \square

Note that if $F \subset F'$, then

$$(2.9) \quad S(F, \epsilon'|_{F'}) \subset S(F', \epsilon')$$

and therefore

$$(2.10) \quad Q(F') \subset Q(F) .$$

^{2.1}Which is looking quite likely. It has been checked for something like three-hundred thousand manifolds.

Let F_1, F_2, \dots, F_n be finite subsets of $G \setminus \{1\}$. Then

$$\bigcap_{i=1}^n Q(F_i) \supset Q(F_1 \cup F_2 \cup \dots \cup F_n) \neq \emptyset$$

since (2.8) holds. This means $\{Q(F)\}$ has the *finite intersection property* (FIP) and each one is closed. Therefore, since $\mathcal{S}(G)$ is compact,

$$\bigcap_{F \subset G \setminus \{1\} \text{ finite}} Q(F) \neq \emptyset.$$

So let $P \in \bigcap Q(F)$.

CLAIM 2.2. P is a positive cone for G .

PROOF. First notice $1 \notin P$ since $1 \notin Q(F)$ for any finite $F \subset G \setminus \{1\}$.

Now we show $g, h \in P$ implies $gh \in P$. Let $F = \{g, h\}$. Then there are $\epsilon(g), \epsilon(h) \in \{\pm 1\}$ such that

$$S(g^{\epsilon(g)}, h^{\epsilon(h)}) \subset P \quad S(g^{\epsilon(g)}, h^{\epsilon(h)})^{-1} \cap P = \emptyset.$$

Therefore $\epsilon(g) = \epsilon(h) = +1$, which implies $gh \in S(g^{\epsilon(g)}, h^{\epsilon(h)}) \subset P$.

Now we show $P \cap P^{-1} = \emptyset$. Let $g \in P$, and $F = \{g\}$. Therefore $S(g) \subset P$, which means $S(g)^{-1} \cap P = \emptyset$, so $g^{-1} \notin P$.

Finally we show $P \amalg P^{-1}G \setminus \{1\}$. Take $g \in G$ such that $g \neq 1$. Let $F = \{g\}$. Then there exists $\epsilon = \pm 1$ such that $S(g^\epsilon) \subset P$ (and $S(g^{-1}) \cap P = \emptyset$) which implies $g^\epsilon \in P$. \square

REMARK 2.5. There exists an analogue of this for BO.

THEOREM 2.4. G is BO if and only if for all finite $F \subset G \setminus \{1\}$ there is some $\epsilon : F \rightarrow \{\pm 1\}$ such that $1 \notin T(F, \epsilon)$ where $T(F, \epsilon)$ is the smallest semigroup which

- (i) contains $S(F, \epsilon)$, and
- (ii) for all $g, h \in T(F, \epsilon)$, $g, h, g^{-1}, g^{-1}hg \in T(F, \epsilon)$.

EXERCISE 2.3. Prove Theorem 2.4.

Let P be a property of groups. A group G is *locally* P if and only if every finitely generated subgroup of G has property P . (So $\text{loc}(\text{loc}(P)) \equiv \text{loc}(P)$.) P is a *local property* if $\text{loc}(P) \implies P$.

THEOREM 2.5. G is locally LO (resp. BO) if and only if G is LO (resp. BO).

PROOF. (\Leftarrow): LO and BO are inherited by subgroups.

(\Rightarrow): Let G be a finite set contained in $G \setminus \{1\}$. Then $\langle F \rangle < G$ is finitely generated. G loc(LO) implies $\langle F \rangle$ is LO. Therefore there exists ϵ such that (2.8) holds (from Theorem 2.3). This is true for all F , so G is LO by Theorem 2.3. The argument for BO is similar, using Theorem 2.4 instead. \square

Corollary 2.6. An abelian group is BO iff it is torsion free.

PROOF. (\Rightarrow): This follows from Lemma 1.3.

(\Leftarrow): G is LO iff G is loc(LO). For H finitely generated inside of torsion free G , then $H \cong \mathbb{Z}^n$, so it is LO. \square

Corollary 2.7. *An arbitrary free group is LO.*

PROOF. Let F be a free group. For H a finitely generated subgroup of F , $H \cong F_n$ for some n . Then H is LO by Corollary 2.7, so F is LO by Theorem 2.5. \square

THEOREM 2.8. *Let $\{G_\lambda\}_{\lambda \in \Lambda}$ be a collection of groups. Then G_λ is LO for all $\lambda \in \Lambda$ if and only if $\ast_{\lambda \in \Lambda} G_\lambda$ is LO.*

PROOF. (\Leftarrow): $G_\lambda < \ast_{\lambda \in \Lambda} G_\lambda$.

(\Rightarrow): There exists a homomorphism

$$G = \ast_{\lambda \in \Lambda} G_\lambda \xrightarrow{\varphi} \prod_{\lambda \in \Lambda} G_\lambda$$

$$g_\lambda \longmapsto (1, \dots, 1, g_\lambda, 1, \dots)$$

So we get a SES

$$(2.11) \quad 1 \rightarrow H \rightarrow \ast_\Lambda G_\lambda \xrightarrow{\varphi} \prod_\Lambda G_\lambda \rightarrow 1$$

where $H = \ker \varphi$. By the Kurosh subgroup theorem

$$H = \left(\ast_\mu H_\mu \right) \ast F$$

where H_μ is a subgroup of a conjugate of G_{λ_μ} in G , and F is a free group. But $H = \ker \varphi$, and $\varphi|_{G_\lambda}$ is injective for all $\lambda \in \Lambda$. Therefore for all $\lambda \in \Lambda$ and $g \in G$ we have $H \cap g^{-1}G_\lambda g = \{1\}$. Therefore $H = F$.

But now G_λ LO for all $\lambda \in \Lambda$ implies $\prod_{\lambda \in \Lambda} G_\lambda$ is LO by Theorem 1.4, and $F = H$ is LL by Corollary 2.7, so G is LO by Theorem 1.13. \square

Let P be a property of groups. A group G is residually P , $\text{res}(P)$, if and only if for all $g \in G \setminus \{1\}$ there exists an epimorphism $\varphi : G \rightarrow H$ such that H has property P , and $\varphi(g) \neq 1$.

REMARK 2.6. Note that P implies $\text{res}(P)$, and $\text{res}(\text{res}(P))$ implies $\text{res}(P)$.

We say P is a *residual property* if and only if $\text{res}(P)$ implies P .

EXAMPLE 2.4. Finiteness is not a residual property. E.g. \mathbb{Z} is $\text{res}(\text{finite})$.

Lemma 2.9. *If P is closed under taking subgroups and direct products, then P is a residual property.*

Corollary 2.10. *LO and BO are residual properties.*

PROOF OF LEMMA 2.9. Suppose G is $\text{res}(P)$. Then for all $g \in G \setminus \{1\}$ there is an epimorphism $\varphi_g : G \rightarrow H_g$ such that H_g has P , and $\varphi_g(g) \neq 1$. The collection of these $\{\varphi_g \mid g \in G \setminus \{1\}\}$ induces a homomorphism

$$\varphi : G \rightarrow \prod_{g \in G \setminus \{1\}} H_g .$$

Then this is injective, and $\varphi_g(g) \neq 1$. H_g has P for all $g \in G \setminus \{1\}$. Therefore $\prod_{g \in G \setminus \{1\}} H_g$ has P . But

$$G \cong \varphi(G) < \prod_{g \in G \setminus \{1\}} H_g$$

so G has P . \square

REMARK 2.7. Residual properties are related to areas of active research. For example the geometrization conjecture is related to residual finiteness of 3-manifolds.

REMARK 2.8. Let G be a group. Let $\text{FQ}(G)$ consist of the finite quotients of G . Then the following is an open question. Let F_2 be a free group of rank 2. If G is a residually finite group such that $\text{FQ}(G) = \text{FQ}(F_2)$ is $G \cong F_2$? Note that $\text{FQ}(F_2)$ consists of the finite groups generated by two elements. So this is really quite concrete.

Another open question is if G_1 and G_2 are residually finitely presented, then does $\text{FQ}(G_1) = \text{FQ}(G_2)$ imply $G_1 \cong G_2$?

EXAMPLE 2.5. $\text{LO}(\mathbb{Z}^n)$ and $\text{LO}(F_n)$ are both the cantor set.

EXAMPLE 2.6. Let B_n denote the braid group. As it turns out $\text{LO}(B_n)$ has isolated points [4].

The following is a strengthening of the fact that LO is a local property.

WARNING 2.1. At this point it is convenient to make the convention that $\{1\}$ is *not* LO.

THEOREM 2.11 (Burns-Hale). *G is LO iff every non-trivial finitely generated subgroup $H < G$ has an LO quotient.*

PROOF. (\implies): G is LO implies H is LO.

(\impliedby): $F = \{g_1, \dots, g_n\} \subset G \setminus \{1\}$ for $n \geq 1$. We show by induction on n that the condition on F in Theorem 2.3 holds. Let $n = 1$. Then $\langle g_1 \rangle$ has an LO quotient by assumption. Therefore g_1 has infinite order, so $1 \notin S(g_1)$. Now suppose $n > 1$. By assumption, there exists a nontrivial homomorphism $\varphi : \langle g_1, \dots, g_n \rangle \rightarrow L$ where L is LO. For some m there exists

$$\varphi(g_i) = \begin{cases} +1 & 1 \leq i \leq m \\ -1 & m < i \leq n \end{cases}$$

By the induction hypothesis there exists $\epsilon_1, \dots, \epsilon_m \in \{\pm 1\}$ such that $1 \notin S(\{g_i^{\epsilon_i} \mid 1 \leq i \leq m\})$. Let $<$ be an LO on L . Define $\epsilon_i \in \{\pm 1\}$ ($m < i \leq n$) so that

$$(2.12) \quad \varphi(g_i^{\epsilon_i}) > 1$$

Then $1 \notin S(\{g_i^{\epsilon_i} \mid 1 \leq i \leq n\})$. □

A group G is *indicible* if either $G = \{1\}$ or there is an epimorphism $G \rightarrow \mathbb{Z}$.

Corollary 2.12. *G is locally indicible implies G is LO.*

REMARK 2.9. Free groups are loc (indicible) so this gives another proof that free groups are LO.

REMARK 2.10. Note that G having an LO quotient does not imply G is LO.

COUNTEREXAMPLE 2. $\mathbb{Z} * \mathbb{Z}/2$ has LO quotient, but is not LO.

We do however have:

THEOREM 2.13. *Let G be a group such that every finitely generated subgroup of infinite index is indicible. Then G is LO if and only if G has an LO quotient.*

PROOF. (\implies): This direction is immediate.

(\impliedby): Apply Theorem 2.11. Let $H < G$, $H \neq \{1\}$, finitely generated.

- Case 1: $[G : H] = \infty$. By hypothesis, H is indicable, so therefore (since H is nontrivial) G has quotient \mathbb{Z} .
- Case 2: $[G : H]$ finite. By hypothesis there exists an epimorphism $\varphi : G \rightarrow Q$ where Q is LO. Therefore Q is infinite, so $\varphi(H) \neq \{1\}$, (since $[Q : \varphi(H)]$ is finite) and therefore H has LO quotient $\varphi(H)$.

□

REMARK 2.11. It turns out that G BO implies G is locally indicable.

REMARK 2.12. We will eventually apply Theorem 2.13 to three-manifold groups. But first we look at surfaces.

2. Surface groups

An n -manifold is a second-countable Hausdorff space M such that for all $x \in M$ x has a neighborhood U such that either

$$(U, x) \cong (\mathbb{R}^n, 0) \quad \text{or} \quad (U, x) \cong (\mathbb{R}_+^n, 0) .$$

Define the interior and boundary as:

$$\begin{aligned} \text{int}(M) &= \{x \in M \mid x \text{ has a neighborhood of the first type}\} \\ \partial M &= \{x \in M \mid x \text{ has a neighborhood of the second type}\} . \end{aligned}$$

Note that $(\text{int}(M)) \cap \partial M = \emptyset$. Also note that $\text{int}(M)$ is an n -manifold with empty boundary, and ∂M is an $(n-1)$ -manifold with empty boundary. M is *closed* if M is compact and $\partial M = \emptyset$.

A *triangulation* of M is a homeomorphism $M \cong |K|$, where K is a locally finite simplicial complex. Whether or not a manifold has a triangulation is a subtle question which wasn't settled until recently [9].

FACT 2. *Every n -manifold has a triangulation for $n \leq 3$.*

This was shown for $n = 2$ in [17] and for $n = 3$ in [12].

For us, a *surface* is a 2-manifold. There is the well-known classification of closed surfaces. In particular, they all either look like S^2 , T^2 , a connect sum of copies of T^2 , the projective plane \mathbb{P}^2 , or connect sums of copies of \mathbb{P}^2 .

There is also a classification of non-compact surfaces.

EXAMPLE 2.7. Consider the plane. Now attach handles as in fig. 1. This is an infinite genus non-compact surface. Now consider the infinite genus surface in fig. 2. Are these homeomorphic? See remark 2.13 for the answer.

Now we consider the following question.

QUESTION 1. Which surface groups $\pi_1(S)$ are LO?

We want to use Theorem 2.11, so we will consider finitely generated subgroups of surface groups. First, recall the following.

Lemma 2.14. *If M is a closed n -manifold, N is a connected n -manifold, and $f : M \rightarrow N$ is an injective map, then f is a homeomorphism.*

This uses the Jordan-Brouwer theorem for S^{n-1} s in S^n . For M compact, N Hausdorff, it is enough to show f is onto.

Lemma 2.15. *Let S be a non-compact surface. Then $H_2(S) = 0$.*




FIGURE 1. The Loch-Ness monster surface obtained by attached infinitely many handles to the plane.



FIGURE 2. The Jacob's ladder surface.

PROOF. Triangulate S . Then we can get compact surfaces $S_1 \subset S_2 \dots \subset S$ such that

$$S = \bigcup_{i=1}^n S_i .$$

$\partial S_i \neq \emptyset$ by Lemma 2.14, so $S_i \simeq$ some 1-complex. Therefore $H_2(S_i) = 0$, for all i . And every 2-cycle in S is contained in some S_i . Therefore $H_2(S) = 0$. \square

Lemma 2.16. *Let S be a surface, δ a circle component of ∂S such that $\pi_1(\delta) \rightarrow \pi_1(S)$ is not injective. Then $S \cong D^2$.*

PROOF. For S compact, this is true by the classification. So let S be non-compact. Let $S^* = S \cup D^2$ glued along δ . Then we have that the following commutes

$$\begin{array}{ccc} \pi_1(\delta) & \longrightarrow & \pi_1(S) \\ \downarrow \cong & & \downarrow \\ H_1(\delta) & \longrightarrow & H_1(S) \end{array} .$$

But now since $\pi_1(\delta) \rightarrow \pi_1(S)$ is not injective, $H_1(\delta) \rightarrow H_1(S)$ cannot be injective either. So now applying Mayer-Vietoris, we get

$$(2.13) \quad H_2(S^*) \cong \ker(H_1(\delta) \rightarrow H_1(S)) ,$$

so by definition this is nonzero. But S^* is noncompact, so this contradicts Lemma 2.15. \square

REMARK 2.13. Have you answered the question from example 2.7 yet? The answer has to do with the number of *ends*, which is defined as follows. Remove compact subsets and count the remaining components. If we minimize the number of components, then this is the number of ends. This is clearly a topological invariant. The loch-ness monster has 1, and Jacob's ladder has 2.

We can also define the notion of the number of ends of a group. As it turns out, $e(G) = 0$ iff G is finite. Then, for example, we have

$$\begin{aligned} e(\mathbb{Z}) &= 2 \\ e(\mathbb{Z}^n) &= 1 \quad (n \geq 2) \\ e(F_n) &= \infty. \end{aligned}$$

Then it turns out that for all G , $e(G) = 0, 1, 2$, or ∞ .

THEOREM 2.17 (Compact core theorem for surfaces). *Let S be a connected surface with $\pi_1(S)$ finitely generated. Then there exists a compact connected $S_0 \xrightarrow{i} S$ such that $i_* : \pi_1(S_0) \rightarrow \pi_1(S)$ is an isomorphism. We call S_0 a compact core of S .*

PROOF. Triangulate S . Let $\gamma_1, \dots, \gamma_n$ be simplicial loops in S such that $\{[\gamma_1], \dots, [\gamma_n]\}$ are generators of $\pi_1(S)$. Let N be a regular neighborhood of $\bigcup_{i=1}^n \gamma_i$ in S . N is a compact surface with $\partial N \neq \emptyset$ (and we can in fact assume it is connected) and $\pi_1(N) \rightarrow \pi_1(S)$ is onto.

Let S_0 be N union with any disk components of S cut along ∂N . S_0 is a compact surface, and $\pi_1(S_0) \rightarrow \pi_1(S)$ is onto. If $\partial S_0 = \emptyset$ then we are done since $S_0 = S$.

So suppose $\partial S_0 \neq \emptyset$. Let δ be a component of ∂S_0 . Since $\pi_1(S_0) \rightarrow \pi_1(S)$ is onto, δ separates S . (If not, there exists a loop $\gamma \subset S$ such that $\gamma \cap \delta$ is a single point. Therefore γ cannot be in S_0 but $\pi_1(S_0) \rightarrow \pi_1(S)$ is onto.)

Let S_1 be the component of S cut along δ such that $S_0 \not\subset S_1$. By definition of S_0 S_1 is not a disk. Therefore by Lemma 2.16 $\pi_1(\delta) \rightarrow \pi_1(S_1)$ is one-to-one. If S_0 is a disk, then $\pi_1(S) = \{1\}$ and we are done. So assume S_0 is not a disk. Then $\pi_1(\delta) \rightarrow \pi_1(S_0)$ is injective. So do this for all the boundary components δ of S_0 . Then we see by Van-Kampen that this is just a big free product:

$$\pi_1(S) \cong \operatorname{colim} \left(\begin{array}{ccccccc} \pi_1(S_1) & & \pi_1(S_2) & & \pi_1(S_3) & \dots & \pi_1(S_k) \\ & \searrow & & \searrow & \downarrow & & \swarrow \\ & & & & \pi_1(S_0) & & \end{array} \right)$$

but by definition this means $\pi_1(S_0) \rightarrow \pi_1(S)$ is injective. □

Lecture 8; February
14, 2020

REMARK 2.14. There is an analogue of this theorem for three-manifolds as well. This is related to the the *Whitehead manifold*, which is a contractible three-manifold not homeomorphic to \mathbb{R}^3 . Whitehead invented this as a counterexample to his own theorem. Professor Cameron says this tells us it is okay to make mistakes as long as you're the one to find the counterexample.

REMARK 2.15. Now Theorem 2.11 implies that if G is locally indicable (and nontrivial) then G is LO.

THEOREM 2.18. *Let S be a surface not homeomorphic to \mathbb{RP}^2 . Then $\pi_1(S)$ is locally indicable.*

PROOF. Let $H < \pi_1(S)$, H finitely generated and nontrivial. Then we want to show it maps to \mathbb{Z} . The point is that there exists a connected covering space $\tilde{S} \rightarrow S$ such that $\pi_1(\tilde{S}) \cong H$. By Theorem 2.17, $H \cong \pi_1(S_0)$ for S_0 a compact surface. Of course $\pi_1(S_0) \neq \{1\}$ (since H was).

Now we claim $S_0 \not\cong \mathbb{RP}^2$. If it was, then $\tilde{S} \cong \mathbb{RP}^2$, so $S \cong \mathbb{RP}^2$, which is a contradiction. Not by the classification of compact surfaces, there exists an epimorphism $H_1(S_0) \rightarrow \mathbb{Z}$, so we can just pre-compose with the map $\pi_1(S_0) \rightarrow H_1(S_0)$, so we get an epimorphism $H \rightarrow \mathbb{Z}$. \square

Corollary 2.19. *Let S be a surface. Then $\pi_1(S)$ is LO if and only if $\pi_1(S) \neq \{1\}$ and $S \not\cong \mathbb{RP}^2$.*

REMARK 2.16. (1) If S is the Klein bottle then $\pi_1(S)$ is locally indicable. But $\pi_1(S)$ is not BO (there exists $a \in \pi_1(S)$ such that a is conjugate to a^{-1}). This shows:

- (a) locally indicable and nontrivial does not imply BO, and
- (b) there is no analog of Burns Hale for BO.
- (2) Locally indicable (and nontrivial) implies LO, but the converse is false. We will see that there are three manifolds M with $H_1(M)$ finite^{2.2} and $\pi_1(M)$ LO.
- (3) It can be shown that if S is a non-compact surface, then $\pi_1(S)$ is free. For example, $\mathbb{R} \setminus$ a cantor set has π_1 isomorphic to a free group on a countably infinite number of generators.
- (4) It can be shown that $\pi_1(S) = 1$ if and only if $S \cong S^2$ or $D^2 \setminus X$ for X a closed subgroup of S^1 .

REMARK 2.17. Colin Adams is a knot theorist who gives lectures in different personas. E.g. a sleazy real-estate agent selling property in hyperbolic space. Once he attended a class posing as a student. He started heckling the lecturer, and eventually the lecturer said “well if you know so much, you come teach the class!” so he did. Some of the students were responding to his heckling, saying “shut up man, he’s doing a great job!” so they were in for surprise when he revealed who he is.

^{2.2}So in particular $\pi_1(M)$ is not locally indicable.

CHAPTER 3

Three-manifolds

Our three-manifolds will always be connected, orientable. They may have boundary and may be non-compact. We will always be working in the PL or smooth category.

Let M_1 and M_2 be oriented 3-manifolds with balls $B_i \subset \text{int}(M_i)$, $B_i \cong B^3$ for $i = 1, 2$. The *connect sum* of M_1 and M_2 is the oriented manifold

$$M_1 \# M_2 = (M_1 \setminus \text{int}(B_1)) \cup_h (M_2 \setminus \text{int}(B_2))$$

for $h : \partial B_1 \rightarrow \partial B_2$ an orientation-reversing homeomorphism. It turns out that $M_1 \# M_2$ is well-defined (up to orientation-preserving homeomorphism). The operation $\#$ is associative, and commutative. Note that $M \# S^3 \cong M$ for all M . Also note that

$$\pi_1(M_1 \# M_2) \cong \pi_1(M_1) * \pi_1(M_2) .$$

We say M is *prime* if $M \cong M_1 \# M_2$ implies M_1 or $M_2 \cong S^3$.

THEOREM (Kneser[7], Milnor[11]). *Let M be a compact, oriented 3-manifold. Then*

$$M \cong \#_{i=1}^n M_i$$

(orientation preserving (op)) where M_i is prime (and not $\cong S^3$) for $1 \leq i \leq n$. Moreover the M_i are unique up to order and orientation-preserving homeomorphism.

Note S^3 corresponds to $n = 0$.

REMARK 3.1. In the same paper [7] Kneser proved some other things which relied on Dehn's lemma. So he was looking closer at Dehn's proof, and found some holes. He wrote to Dehn who was on vacation to find out that he agreed there was something fishy. Thus ensued a great correspondence between the two trying to fix it. It was eventually fixed in [13].

For M compact, and

$$M \cong \#_{i=1}^n M_i$$

where the M_i are prime, we have

$$\pi_1(M) \cong \bigast_{i=1}^n \pi_1(M_i) .$$

So $\pi_1(M)$ is LO iff $\pi_1(M_i)$ is LO (for $1 \leq i \leq n$). This is also true for BO.

EXERCISE 3.1. Show $\pi_1(M)$ is locally indicable iff $\pi_1(M_i)$ is locally indicable for all $1 \leq i \leq n$. [Hint: Use the Kurosh subgroup theorem.]

The upshot is that for M compact, to answer LO, BO, or locally indicable, we may assume M is prime.

REMARK 3.2. There are noncompact three manifolds that cannot be expressed as $\#$ of prime manifolds.

M is irreducible if every $S^2 \subset M$ bounds a $B^3 \subset M$.

FACT 3. M is irreducible iff M is prime and not homeomorphic (op) to $S^1 \times S^2$.

The point being that $S^1 \times S^2$ is prime.

THEOREM (Perelman[14–16]). Let M be a closed 3-manifold with universal cover \tilde{M} .

- (1) If $\pi_1(M)$ is finite, then $\tilde{M} \cong S^3$ and the action of $\pi_1(M)$ on S^3 is as a subgroup of $\mathrm{SO}(4)$.
- (2) If $\pi_1(M)$ is infinite and M is irreducible, then $\tilde{M} \cong \mathbb{R}^3$.

COROLLARY (Poincaré conjecture). If M is closed and $\pi_1(M) = 1$, $M \cong S^3$.

REMARK 3.3. We know $\pi_1(M)$ infinite implies \tilde{M} is noncompact. Then M irreducible implies $\pi_2(M) = 0$ (as we will see soon) so by standard stuff, \tilde{M} is contractible. But, there are contractible non-compact 3-manifolds without boundary which are not homeomorphic to \mathbb{R}^3 .

The 3-manifolds with π_1 finite can be completely described. They're all Seifert fiber spaces.

EXAMPLE 3.1. Let $p, q \in \mathbb{Z}$ such that $p \geq 2$ $(p, q) = 1$. Recall we have a \mathbb{Z}/p action on \mathbb{C}^2 by

$$(z, w) \mapsto (e^{2\pi i/p} z, e^{2\pi q i/p} w)$$

Now the restriction of this action to S^3 is free, so we can quotient by it to get the lens space $L(p, q)$. Then

$$\pi_1(L(p, q)) = \mathbb{Z}/p.$$

Nonetheless, Alexander showed that $L(5, 1) \not\cong L(5, 2)$.

THEOREM (Redemeister). $L(p, q)$ is homeomorphic to $L(p, q')$ iff either $q \cong q' \pmod{p}$ or $qq' \cong 1 \pmod{p}$.

The \Leftarrow direction is easy.

THEOREM (Perelman[14–16]). For M and M' closed three-manifolds, M prime and not a lens space, then $\pi_1(M) \cong \pi_1(M')$ implies $M' \cong M$.

So “prime three-manifolds are pretty much determined by their fundamental group”.

REMARK 3.4. The restriction to prime is necessary here. Let M be an oriented three-manifold such that M is not homeomorphic (op) to $-M$. For example, $M = L(3, 1)$ or the Poincaré homology sphere.

Then $\pi_1(M \# M) \cong \pi_1(M \# (-M))$, but by prime decomposition, $M \# M \not\cong M \# (-M)$.

Lecture 11;
February 25, 2020

1. Higher homotopy groups

We will give a basic overview of higher homotopy groups. Define

$$(3.1) \quad \pi_n(X, x_0) = \left\{ \text{homotopy classes of maps } (S^n, s_0) \xrightarrow{f} (X, x_0) \right\}.$$

As it turns out, we can define a composition on this set with respect to which this becomes a group. This is called the n th homotopy group of (X, x_0) . As soon as $n \geq 2$, this group is abelian. Note that for $f : (X, x_0) \rightarrow (Y, y_0)$, we get an induced homomorphism $f_* :$

$\pi_n(X, x_0) \rightarrow \pi_n(Y, y_0)$. I.e. the π_n are covariant functors. If X is path-connected, this implies that for all $x_0, x_1 \in X$,

$$(3.2) \quad \pi_n(X, x_0) \cong \pi_n(X, x_1) .$$

Let $f : S^n \rightarrow X$. This induces a homomorphism

$$(3.3) \quad f_* : \underbrace{H_n(S^n)}_{\cong \mathbb{Z}} \rightarrow H_n(X) .$$

Now $f \simeq g$ implies $f_* = g_*$, so we can define

$$(3.4) \quad h : \pi_n(X) \rightarrow \pi_n(X)$$

by sending $h([f]) = f_*(1)$. This is well-defined, and it can be shown that this is in fact a homomorphism. This is called the Hurewicz homomorphism. For $n = 1$, h is just the abelianization.

Any covering projection $p : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$ induces an isomorphism $p_* : \pi_n(\tilde{X}, \tilde{x}_0) \rightarrow \pi_n(X, x_0)$ for $n \geq 2$. This follows from the fact that $\pi_1(S^n) = 1$ if $n \geq 2$, and the lifting criterion.

2. Back to three-manifolds

Let M be a connected, orientable (sometimes oriented) 3-manifold, possibly non-compact and possibly with boundary. Let $S \subset M$ be an embedded 2-sphere. We say S is *essential*, if $S \not\simeq \text{pt}$. I.e. the homotopy class $[S] \neq 0 \in \pi_2(M)$. We say S is *incompressible* if S does not bound a 3-ball in M . Recall a manifold M is *irreducible* if and only if every 2-sphere bounds a ball, i.e. every 2-sphere is compressible.

THEOREM (Sphere theorem [13], [19]). *Let M be a 3-manifold with $\pi_2(M) \neq 0$. Then there exists an embedded essential 2-sphere in M .*

This implies the asphericity of knots: if K is a knot in S^3 , then $\pi_2(S^3 \setminus K) = 0$. The proof uses the “tower construction”.

THEOREM 3.1. *Let M be a three-manifold and S a 2-sphere in M . Then S is incompressible if and only if S is essential.*

PROOF. (\Leftarrow): If S bounds a $B^3 \subset M$ then $S \simeq \text{pt}$.

(\Rightarrow): Let S be incompressible. Suppose (for a contradiction) that $S \simeq \text{pt}$, i.e. $[S] = 0 \in \pi_2(M)$, so $[S] = 0 \in H_2(M)$ (after applying h from above). Therefore S separates M . [Recall for S non-separating, then there is a loop $\gamma \subset M$ such that γ meets S transversely in 1 point, so $[S] \neq 0 \in H_2(M)$.] So this separates M into M_1 and M_2 . So $M = M_1 \cup_S M_2$. Now $[S] = 0 \in H_2(M)$ implies S bounds a 3-chain in M . Therefore M_1 (say) is compact and $\partial M_1 = S$.

If $\pi_1(M_1) = 1$, then $M_1 \cong B^3$ (by the Poincaré conjecture). Therefore S is compressible, which is a contradiction. So $\pi_1(M) \cong \pi_1(M_1) * \pi_1(M_2)$.

Case (1) $\pi_1(M_2) \neq 1$: Let \tilde{M} be the universal covering of M . Then $\pi_1(M_1) \neq 1$, so \tilde{M}_1 has more than one boundary component. Let $\tilde{S} \subset \tilde{M}$ be a lift of S . Then every component of \tilde{M} cut along \tilde{S} is non-compact. Therefore \tilde{S} does not bound a (finite) 3-chain in \tilde{M} , so $[\tilde{S}] \neq 0 \in H_2(\tilde{M})$ so $[\tilde{S}] \neq 0 \in \pi_2(\tilde{M})$. Therefore

$$(3.5) \quad [S] = p_*([\tilde{S}]) \neq 0 \in \pi_2(M) .$$

Case (2) $\pi_1(M_2) = 1$: Then \tilde{M} is \tilde{M}_1 with some copies of noncompact M_2 attached. By hypothesis S is incompressible so $M_2 \not\cong B^3$. Therefore $\pi_1(M_2) = 1$, so either M_2 is noncompact or $\partial M_2 \setminus S \neq \emptyset$. Now pick some lift \tilde{S} . It cannot bound a finite chain in either direction since both either have nontrivial boundary, or are noncompact. So as before, $[\tilde{S}] \neq 0 \in H_2(\tilde{M})$, so nonzero in $\pi_1(\tilde{M})$, so $[S] \neq 0 \in \pi_1(M)$. \square

Corollary 3.2. *If M is a 3-manifold, then M is irreducible if and only if $\pi_2(M) = 0$.*

PROOF. Combine Theorem 3.1 with the sphere theorem. \square

Corollary 3.3. *For $\tilde{M} \rightarrow M$ a covering, M is irreducible iff \tilde{M} is irreducible.*

PROOF. This follows from Theorem 3.1 since $\pi_2(\tilde{M}) \cong \pi_2(M)$. \square

REMARK 3.5. This falls out for nothing by Theorem 3.1, but we did use the Poincaré conjecture in our proof. This was actually known before the Poincaré conjecture. The converse is easy, but the forward implication is hard.

THEOREM. *Let M be a three-manifold with $\pi_1(M)$ finitely generated. Then there exists a compact three-manifold $N \xrightarrow{i} M$ such that $i_* : \pi_1(N) \rightarrow \pi_1(M)$ is an isomorphism.*

Recall a group G is *coherent* if $H < G$ finitely generated implies it is finitely presentable.

COROLLARY. *For M a three-manifold, $\pi_1(M)$ is coherent.*

PROOF. For H a finitely generated subgroup of $\pi_1(M)$, $H \cong \pi_1(\tilde{M})$ for $\tilde{M} \rightarrow M$ a covering. Then apply the Scott core theorem to \tilde{M} . \square

REMARK 3.6. $\mathrm{SL}_2(\mathbb{Z})$ is *virtually* free. (It has a free subgroup of finite index.)

Free groups are coherent, so $\mathrm{SL}(2, \mathbb{Z})$ is coherent. We know $\mathrm{SL}_n(\mathbb{Z})$ is incoherent for $n \geq 4$.

QUESTION 2 (Serre). Is $\mathrm{SL}_3(\mathbb{Z})$ coherent?

REMARK 3.7. Once at a conference in the UK, Professor Gordon was playing table tennis. He asked a guy standing nearby if he wanted to play, and the guy beat him. He asked for his name and he replied “Jean-Pierre Serre”.

Lemma 3.4. *Let M be a closed orientable n -manifold, n odd. Then $\chi(M) = 0$.*

PROOF. Let \mathbb{F} be a field. By Poincaré duality,

$$(3.6) \quad H_i(M; \mathbb{F}) \cong H^{n-i}(M; \mathbb{F}) \cong H_{n-i}(M; \mathbb{F})$$

where the second equality follows from the universal coefficient theorem. Therefore

$$(3.7) \quad \chi(F) = \sum_{i=0}^n (-1)^i \dim H_i(M; \mathbb{F}) = 0$$

if n is odd. \square

Lemma 3.5. *Let M be a compact three-manifold and \mathbb{F} a field. Then*

$$\dim H_1(M; \mathbb{F}) \geq \frac{1}{2} \dim H_1(\partial M; \mathbb{F}) .$$

PROOF. Let $2M$ be the *double* of M . This is the union of M with M (i.e. M with the opposite orientation) glued along the boundary. Let n be the number of ∂ -components of M . Then by Lemma 3.4,

$$(3.8) \quad 0 = \chi(2M) = 2\chi(M) - \chi(\partial M)$$

which means $\chi(M) = \chi(\partial M)/2$.

We will use the Betti number notation:

$$(3.9) \quad \beta_2(M) = \dim H_2(M; \mathbb{F}) .$$

Then

$$(3.10) \quad \chi(M) = 1 - \beta_1(M) + \beta_2(M)$$

Then

$$(3.11) \quad \chi(\partial M) = n - \beta_1(\partial M) + n .$$

Now by the universal coefficient theorem and Poincaré-Lefschetz duality we get

$$(3.12) \quad H_2(M; \mathbb{F}) \cong H^2(M; \mathbb{F}) \cong H_1(M, \partial M; \mathbb{F}) .$$

So now we have the exact sequence of the pair $(M, \partial M)$:

$$(3.13) \quad \dots \longrightarrow H_1(M, \partial M; \mathbb{F}) \longrightarrow \underbrace{H_0(\partial M; \mathbb{F})}_{\mathbb{F}^n} \longrightarrow \underbrace{H_0(M; \mathbb{F})}_{\mathbb{F}} \longrightarrow 0$$

which means

$$(3.14) \quad \beta_2(M) = \beta_1(M, \partial M) \geq n - 1 .$$

Therefore

$$(3.15) \quad 1 - \beta_1(M) + n - 1 \leq \chi(M) = \frac{1}{2}\chi(\partial M) = n - \frac{1}{2}\beta_1(\partial M)$$

so

$$(3.16) \quad \beta_1(M) \geq \frac{1}{2}\beta_1(\partial M)$$

as desired. \square

Corollary 3.6. *Let M be a compact 3-manifold with a boundary component not homeomorphic to S^2 . Then $H_1(M)$ is infinite.*

PROOF. $\dim H_1(M; \mathbb{R}) \geq 1$ and by the universal coefficient theorem, this is $H_1(M) \otimes \mathbb{R} \cong \mathbb{R}^n$. Then this is equivalent to

$$(3.17) \quad H_1(M) \cong \mathbb{Z}^n \oplus A$$

for A some finitely abelian group. \square

Lecture 12;
February 27, 2020

Lemma 3.7. *Let M be a prime three-manifold, H a finitely generated subgroup of $\pi_1(M)$ such that either*

- (1) *H has infinite index in $\pi_1(M)$, or*
- (2) *M is not closed.*

Then H is indicable (i.e. either $H = \{1\}$ or there is some epimorphism $H \rightarrow \mathbb{Z}$).

PROOF. If $M = S^1 \times S^2$ this is clear, since $\pi_1(M) \cong \mathbb{Z}$. So we may assume M is irreducible. In case (2), replacing M by $M \setminus \partial M$, we may assume M is noncompact. There exists a covering $\tilde{M} \rightarrow M$ with $\pi_1(\tilde{M}) \cong H$. In both cases, \tilde{M} is noncompact. Now we can use the Scott Core theorem to show that there is some compact submanifold N of \tilde{M} such that the inclusion $\pi_1(N) \xrightarrow{\cong} \pi_1(\tilde{M})$ is an isomorphism. Let S be a 2-sphere component of ∂N . Corollary 3.3 implies \tilde{M} is irreducible. Therefore S bounds a 3-ball $B \subset \tilde{M}$. If $N \subset B$, then $\pi_1(N) \rightarrow \pi_1(\tilde{M})$ factors through $\pi_1(B) = 1$. So $\pi_1(\tilde{M}) \cong H = \{1\}$. So H is indicable. So assume $H \neq \{1\}$. Then we have $N \not\subset B$. So replace N with $N \cup B$, so $\pi_1(N \cup B) \cong \pi_1(N)$. So now we can assume N has no 2-sphere boundary components. If N is closed, then $N = \tilde{M}$, but this contradicts the fact that \tilde{M} is noncompact. Therefore $\partial N \neq \emptyset$, so by Corollary 3.6 $H_1(N)$ is infinite, so $\pi_1(N) \cong H$ maps onto \mathbb{Z} , so H is indicable. \square

Corollary 3.8. *For M a prime 3-manifold, $\pi_1(M)$ is infinite. Then $\pi_1(M)$ is torsion-free.*

PROOF. Suppose we have an element $g \in \pi_1(M)$ of finite order $n > 1$. Therefore $\langle g \rangle \cong \mathbb{Z}/n$. But this has infinite index in $\pi_1(M)$, but by Lemma 3.7 this means it is indicable. But of course it is nontrivial and does not map onto \mathbb{Z} , so this is a contradiction. \square

Let M be a closed three-manifold with $H_1(M; \mathbb{Z}) = 0$. Then

$$(3.18) \quad H_2(M; \mathbb{Z}) \cong H^1(M; \mathbb{Z}) \cong \text{Hom}(H_1(M; \mathbb{Z}), \mathbb{Z}) = 0$$

where the first isomorphism follows from Poincaré duality, and the second follows from the universal coefficient theorem. Now note that $H_3(M; \mathbb{Z}) \cong H^0(M; \mathbb{Z}) \cong \mathbb{Z}$, and $H_q(M; \mathbb{Z}) = 0$ for $q \geq 4$, so

$$(3.19) \quad H_*(M; \mathbb{Z}) \cong H_*(S^3; \mathbb{Z}) .$$

In this case we say M is an integral homology sphere, or \mathbb{Z} HS. Similarly, if $H_1(M; \mathbb{Q}) = 0$ (equivalently $H_1(M; \mathbb{Z})$ finite) then

$$(3.20) \quad H_*(M; \mathbb{Q}) \cong H_*(S^3; \mathbb{Q}) .$$

In this case we say M is a rational homology sphere, or \mathbb{Q} HS. As it turns out, there are infinitely many \mathbb{Z} HS's M with $\pi_1(M) \neq 1$. There are also infinitely many \mathbb{Q} HS's that are not \mathbb{Z} HS's. For example, lens spaces are \mathbb{Q} HS's.

Let $K \subset S^3$ be a knot in S^3 . Then the tubular neighborhood is $N(K) \cong S^2 \times D^2$. Now take

$$(3.21) \quad X = \text{Cl}(S^3 \setminus N(K)) .$$

Note that $\partial X \cong T^2$. Take μ to be some meridional curve in the boundary, and λ some longitudinal one. There is (up to isotopy) a unique such λ in X such that $\lambda \sim 0$. Now for any α an essential simply closed curve in ∂X , $\alpha \sim p\mu = q\lambda$ for $(p, q) = 1$. Now consider

$$(3.22) \quad X \cup_h (S^1 \times D^2)$$

where $h : S^1 \times D^2 \rightarrow \partial X$ such that

$$h(\text{pt} \times \partial D^2) = \alpha .$$

So this gives a closed three-manifold $K(\alpha) = K(p/q)$. This is called the α (or p/q) Dehn surgery on K . Recall $H_1(X) \cong \mathbb{Z}$ given by μ . So when we add in this solid torus, we're killing α . λ was dead anyway, so we're really just killing the p th power of the meridian. So

$$(3.23) \quad H_1(K(p/q)) \cong \mathbb{Z}/p.$$

For $p = 1$ we get an ZHS, and for $p > 1$ we get a QHS. The following is an open question:

QUESTION 3. Does a prime ZHS (which is not S^3 or the Poincaré homology sphere) have left-orderable fundamental group?

This is related to the so-called L -space conjecture.

THEOREM 3.9. *Let M be a prime 3-manifold. Then $\pi_1(M)$ is locally indicable. Equivalently, M is S^3 , or M is not a QHS.*

PROOF. (\implies): Suppose M is a QHS. Then $\pi_1(M)$ is finitely generated. Therefore $\pi_1(M)$ LI implies $\pi_1(M)$ is indicable, so $\pi_1(M) = 1$ or $H_1(M)$ is infinite.

(\impliedby): Suppose M is not a QHS. Then either

- (i) M is closed and $H_1(M)$ is infinite, or
- (ii) M is not closed.

Let H be a finitely generated subgroup of $\pi_1(M)$. In the first case, $[\pi_1(M) : H] = \infty$ so H is indicable by Lemma 3.7. If $[\pi_1(M) : H]$ is finite, then $[H_1(M) : h(H)]$ is finite. Therefore $h(H)$ is infinite, and so H maps onto \mathbb{Z} . Case (ii) follows from Lemma 3.7. \square

Since (for $G \neq \{1\}$) locally indicable implies LO (from Corollary 2.12).

Corollary 3.10. *If M is a prime 3-manifold which is not a QHS, then $\pi_1(M)$ is LO.*

In fact, we can do better, using Theorem 2.13.

THEOREM 3.11 ([1]). *Let M be a prime three-manifold. Then $\pi_1(M)$ is LO iff $\pi_1(M)$ has a LO quotient.*

PROOF. (\implies): this direction is trivial.

(\impliedby): this follows from Lemma 3.7 and Theorem 2.13. \square

CHAPTER 4

Seifert fiber(ed) spaces

DEFINITION 4.1. A *Seifert fiber space* (SFS) is a compact (orientable) 3-manifold M that is a disjoint union of circles (called the *fibers*) such that each circle has a neighborhood which is a union of fibers and is isomorphic to a *fibered solid torus*. This is

$$(4.1) \quad (D^2 \times I) / ((x, 1) \sim (h(x), 0))$$

where $h : D^2 \rightarrow D^2$ is rotation through $2\pi q/p$ for $p > 0$ and $(p, q) = 1$. The fibers are the images of $0 \times I$ (the central fiber) and

$$(4.2) \quad (x \times I) \cup (h(x) \times I) \cup \dots \cup (h^{p-1}(x) \times I)$$

for $x \neq 0$. If $p > 1$, the central fiber is *exceptional*: other fibers are *ordinary*.

M compact implies there are only finitely many exceptional fibers. Let $\pi : M \rightarrow F$ be the quotient map defined by identifying each fiber to a point. For a fibered solid torus, the quotient $\cong D^2$. So F is a surface (called the base surface) and $\pi : \partial M \rightarrow \partial F$ is an S^1 -bundle projection. Therefore ∂M is the disjoint union of finitely many tori (possibly empty). If M has no exceptional fibers, then $\pi : M \rightarrow F$ is an S^1 -bundle projection. Conversely, an orientable S^1 -bundle over F is a SFS.

Let $N_0 \subset \text{int}(M)$ be a fibered solid torus neighborhood of an ordinary fiber, and (disjoint) $N_i \subset \text{int}(M)$ solid tori neighborhood of the exceptional fibers with parameters p_i, q_i .

Let $D_i = \pi(N_i)$. These are disks in F . Then the restriction

$$(4.3) \quad \pi : \underbrace{(M \setminus \bigcup_{i=1}^n N_i)}_{M_0} \rightarrow \underbrace{F \setminus \bigcup_{i=1}^n D_i}_{F_0}$$

is an S^1 -bundle projection. Let $\alpha_1, \dots, \alpha_k \subset F_0$ be disjoint properly embedded arcs such that if we cut F_0 along $\bigcup_{i=1}^k \alpha_i$, to get $F_0|_{\bigcup_{i=1}^k \alpha_i}$, this is a disk B . Then $\pi^{-1}\alpha_i$ is an S^1 bundle over α_i , i.e. an annulus $A_i \subset M_0$. So we have an S^1 -bundle over B given by

$$(4.4) \quad M_0|_{\bigcup_{i=1}^k A_i} \cong B \times S^1$$

with copies A_i^\pm of A_i inside $\partial(B \times S^1)$. Therefore we can recover M_0 by taking the quotient:

$$M_0 \cong (B \times S^1) / (A_i^+ \sim A_i^-, 1 \leq i \leq k) .$$

We can isotope these identifications so that $\alpha_i^- \sim \alpha_i^+$. So we actually get a copy of $F_0 \subset M_0$ which is a section of the S^1 -bundle:

$$(4.5) \quad F_0 \subset M_0 \xrightarrow[\text{id}]{\pi} F_0 .$$

Note that $\pi_1(B \times S^1) \cong \mathbb{Z}$, generated by the class of the ordinary fiber, written h . Now there are two cases:

Lecture 13; March 3, 2020

- (O) F orientable: $g(F) = g \geq 0$, $|\partial F| = m \geq 0$. Now $\pi_1(F_0)$ has generators $a_1, b_1, \dots, a_g, b_g$ corresponding to the genus; d_1, \dots, d_m corresponding to the boundary components; and then c_1, \dots, c_n for the exceptional fibers. Then since $M_0 \cong F_0 \times S^1$, we have that

$$(4.6) \quad \pi_1(M_0) \cong \pi_1(F_0) \times \mathbb{Z}$$

where this copy of \mathbb{Z} is generated by h .

- (N) F non-orientable: Now $\pi_1(F_0)$ has generators a_1, \dots, a_g (where these now come from twisted strips); d_1, \dots, d_m ; and c_1, \dots, c_n . Now M_0 is a twisted S^1 -bundle over F_0 . The presentation is explicitly:

$$(4.7) \quad \pi_1(M_0) \cong \left\langle a_1, \dots, a_g, d_1, \dots, d_m, c_1, \dots, c_n, h \left| \begin{array}{l} a_i^{-1} h a_i = h^{-1}, 1 \leq i \leq g, \\ h \leftrightarrow d_i, h \leftrightarrow c_i \end{array} \right. \right\rangle.$$

In both cases, $\pi_1(\partial N_i) \cong \mathbb{Z} \times \mathbb{Z}$ generated by h and C_i for $0 \leq i \leq n$. Now let λ_i and μ_i be a longitude, meridian pair for N_i , $0 \leq i \leq n$. Then $\pi_1(\partial N_i) \cong \mathbb{Z}^2$ is generated by $\{\lambda_i, \mu_i\}$. In terms of this basis, an ordinary fiber is:

$$(4.8) \quad h = \lambda_i^{p_i} \mu_i^{q_i}$$

where the (p_i, q_i) are the Seifert invariants of the fibered solid torus N_i . Then we can write the c_i 's as:

$$(4.9) \quad c_i = \lambda_i^{r_i} \mu_i^{s_i}$$

where $p_i s_i - q_i r_i = 1$. Therefore we get the relation

$$(4.10) \quad \mu_i = c_i^{p_i} h^{-r_i}.$$

Therefore by van Kampen

$$(4.11) \quad \pi_1(M) \cong \pi_1(M_0) / (c_i^{p_i} = h^{r_i}, 0 \leq i \leq n).$$

For $p_0 = 1$ let $b = -r_0$. Note that (in $\pi_1(F)$)

$$(4.12) \quad c_0^{-1} = \begin{cases} \prod_{i=1}^g [a_i, b_i] \prod_{i=1}^m d_i \prod_{i=1}^n c_i & \text{Case (O)} \\ \prod_{i=1}^g a_i^2 \prod_{i=1}^m d_i \prod_{i=1}^n c_i & \text{Case (N)} \end{cases}$$

THEOREM 4.1. *Let M be a SFS. Then in case (O), $\pi_1(M)$ has presentation:*

$$(4.13) \quad \left\langle a_1, b_1, \dots, a_g, b_g, d_1, \dots, d_m, c_1, \dots, c_n, h \left| \begin{array}{l} h \leftrightarrow a_i, b_i, d_i, c_i; c_i^{p_i} = h^{r_i}; \\ \prod [a_i, b_i] \prod d_i \prod c_i = h^b \end{array} \right. \right\rangle.$$

In case (N), $\pi_1(M)$ has presentation:

$$(4.14) \quad \left\langle a_1, \dots, a_g, d_1, \dots, d_m, c_1, \dots, c_n, h \left| \begin{array}{l} a_i^{-1} h a_i = h^{-1}; h \leftrightarrow c_i, d_i; c_i^{p_i} = h^{r_i}; \\ \prod a_i^2 \prod d_i \prod c_i = h^b \end{array} \right. \right\rangle.$$

REMARK 4.1. Recall the a_i (and b_i) come from the base surface, (in the (N) case these are orientation reversing) the d_i come from the boundary components, the c_n come from the singular fibers.

- REMARK 4.2.** (1) $\langle h \rangle$ is central in $\pi_1(M)$ in case (O), and normal in case (N).
 (2) The fundamental group of F is

$$\pi_1(F) \cong \pi_1(M) / (h = 1, c_i = 1, 1 \leq i \leq n)$$

so

$$\pi_*: \pi_1(M) \rightarrow \pi_1(F)$$

is onto.

- (3) Suppose $n > 0$, i.e. we have at least one singular fiber. Let α be a properly embedded arc in F_0 joining ∂ -components C_i and C_j for $0 \leq i, j \leq n$, $i \neq j$. Then $\pi^{-1}(\alpha)$ is an annulus H in M_0 , with boundary components in ∂N_i and ∂N_j . Now by Dehn twisting M_0 along A , k times gives a homeomorphism $M_0 \rightarrow M_0$. This is the identity outside of a neighborhood of A , and

$$\begin{array}{lll} \text{on } \partial N_i & h \mapsto h & c_i \mapsto c_i h^k = c'_i \\ \text{on } \partial N_j & h \mapsto h & c_j \mapsto c_j h^{-k} = c'_j \end{array}$$

DIGRESSION 1 (Dehn twists). Let A be a properly embedded^{4.1} annulus inside of a 3-manifold M . A has a neighborhood $N(A) \cong A \times I \subset M$. Then Dehn twist along A is the homeomorphism $h : M \rightarrow M$ such that $h|_{\text{Cl}(M \setminus A)} = \text{id}$, and $h|_{A \times I} : A \times I \rightarrow A \times I$ sends

$$(4.15) \quad ((\theta, s), t) \mapsto ((\theta = 2\pi t, s)) .$$

So the meridians go to:

$$\mu_i = c_i^{p_i} h^{-r_i} \mapsto c_i^{p_i} h^{-r'_i} \quad \mu_j = c_j^{p_j} h^{-r_j} \mapsto c_j^{p_j} h^{-r'_j}$$

where $r'_i = r_i + kp_i$ and $r'_j = r_j - kp_j$. Therefore

$$(4.16) \quad \sum_{i=0}^n \frac{r_i}{p_i}$$

is unchanged. Now recall $p_0 = 1$, $r_0 = -b$ so

$$(4.17) \quad -b + \sum_{i=1}^n \frac{r_i}{p_i}$$

is also unchanged. This is called the Euler number of the Seifert structure.

When $n = 0$, we just have a circle bundle, and this gives us the Euler number of this circle bundle. Using this, we can either normalize the r_i so that $0 < r_i < p_i$, or if $n > 0$, we can take $b = 0$.

- (4) One can show that M is irreducible unless $M \cong S^1 \times S^2$ or $\mathbb{RP}^3 \# \mathbb{RP}^3$.
 (5) For most SFS's, the SF structure is unique.
 (6) It follows from Perelman that if M is a closed three-manifold with $\pi_1(M)$ finite, then M is a SFS. Those with finite π_1 are

$$\begin{array}{llll} F = S^2 & n \leq 2 & M \not\cong S^1 \times S^2 & (S^2, \text{lens spaces}) \\ F = S^2 & n = 3 & \sum_{i=1}^3 \frac{1}{p_i} > 1 & \left(\begin{array}{l} \text{platonic triples: } (2, 2, p), \\ (2, 3, 3), (2, 3, 4), (2, 3, 5) \end{array} \right) \\ F = \mathbb{RP}^2 & n = 0, b \neq 0 & & (n = b = 0 \leadsto \mathbb{RP}^3 \# \mathbb{RP}^3) \\ & n = 1 & & \end{array}$$

- (7) SFS's are one of the building blocks of the JSJ decomposition of three-manifolds. The other pieces are hyperbolic three-manifolds.

^{4.1}Recall this means $A \cap \partial M = \partial A$.

THEOREM 4.2. *Let M be a SFS. Then $\pi_1(M)$ is BO if and only if M is either $S^1 \times S^2$ or an S^1 -bundle over an orientable surface not homeomorphic to S^2 .*

REMARK 4.3. The S^1 bundles over S^2 are $S^1 \times S^2 = L(0, 1)$, $S^3 = L(1, 1)$, and $L(p, 1)$ for $p \geq 2$.

PROOF. (\Leftarrow): $S^1 \times S^2$ is immediate. Let M be an S^1 bundle over an orientable surface $F \not\cong S^2$. For any bundle we have a homotopy exact sequence:

$$(4.18) \quad \underbrace{1}_{\pi_2(F)} \rightarrow \underbrace{\pi_1(S^1)}_{=\mathbb{Z}} \rightarrow \pi_1(M) \rightarrow \pi_1(F) \rightarrow 1.$$

FACT 4. $\pi_1(F) \neq 1$ implies $\pi_1(F)$ is BO.

We will prove this later. M orientable implies the conjugation action of $\pi_1(F)$ on \mathbb{Z} is by the identity, so the BO on \mathbb{Z} is conjugation invariant, so $\pi_1(M)$ is BO by Theorem 1.13.

(\Rightarrow): We first need the following lemma.

Lemma 4.3. *In case (N), $h \neq 1 \in \pi_1(M)$.*

PROOF. First take $n = 0$. Then we can kill the d_i 's and a_i 's for $i > 1$ to get the quotient of $\pi_1(M)$:

$$(4.19) \quad \langle a, h \mid a^{-1}ha = h^{-1}, a^2 = h^b \rangle.$$

For b even, this has quotient

$$(4.20) \quad \langle a, h \mid a^2 = h^2 = (ah)^2 = 1 \rangle \cong \mathbb{Z}/2 \times \mathbb{Z}/2.$$

For b odd, this has quotient

$$(4.21) \quad \langle a, h \mid a^{-1}ha = h^{-1}, h^2 = 1, a^2 = h \rangle \cong \langle a \mid a^4 = 1 \rangle \cong \mathbb{Z}/4.$$

For $n > 0$, we also kill the c_i for $i \geq 2$, so we get a quotient

$$(4.22) \quad \langle a, c, h \mid h^2 = 1, a \leftrightarrow h, c^p = h^r, a^2c = 1 \rangle \cong \langle a, h \mid h^2 = 1, a \leftrightarrow h, a^{2p} = h^{-1} \rangle$$

$$(4.23) \quad \cong \begin{cases} \mathbb{Z}/2 \times \mathbb{Z}/2p & r \text{ even} \\ \mathbb{Z}_{4p} & r \text{ odd} \end{cases}.$$

□

Lemma 4.4. *In case (N), $\pi_1(M)$ is not BO.*

PROOF. $a^{-1}ha = h^{-1}$, and $h \neq 1$ by Lemma 4.3, so $\pi_1(M)$ is not BO. □

Lemma 4.5. *Suppose G is BO. If $h^m \leftrightarrow g$ for some $n \neq 0$, then $h \leftrightarrow g$.*

PROOF. Since $x \leftrightarrow g$ iff $x^{-1} \leftrightarrow g$, we may assume $n > 0$. Suppose $h \not\leftrightarrow g$, i.e. $ghg^{-1}h^{-1} \neq 1$. WLOG this means $ghg^{-1}h^{-1} > 1$, so $ghg^{-1} > h$. Therefore $(ghg^{-1})^n = gh^n g^{-1} > h^n$, so $gh^n g^{-1}h^{-n} > 1$ which is a contradiction. □

Lemma 4.6. *In case (O), if $g > 0$ and $n > 0$ then $\pi_1(M)$ is not BO.*

PROOF. Assume $\pi_1(M)$ is BO. We have the relation

$$(4.24) \quad c_1^{p_1} = h^{r_1}.$$

Now recall in case (O), h is central, so $c_1^{p_1}$ is central, so c_1 is central (by Lemma 4.5). Now kill the a_i 's, b_i 's, d_i 's and c_i 's for $i > 1$, and h . So we get a quotient

$$(4.25) \quad \langle a, b, c \mid c^p = 1, c = [a, b] \rangle \cong \langle a, b \mid [a, b]^p = 1 \rangle$$

for $p \geq 2$. Now we have the following lemma which will finish the proof:G

Lemma 4.7. $[a, b]$ is not central in $G_p = \langle a, b \mid [a, b]^p = 1 \rangle$ (for $p \geq 2$).

PROOF. Define a representation $G_p \rightarrow S_{2p}$ by

$$(4.26) \quad a \longmapsto (1\ 2)(3\ 4)\dots(2p-1\ 2p)$$

$$b \longmapsto (1\ 2\ 3\ \dots\ 2p)$$

so we have that

$$(4.27) \quad b^{-1}ab = (2\ 3)(4\ 5)\dots(2p\ 1) .$$

Therefore

$$(4.28) \quad [a, b] = a^{-1}b^{-1}ab = (2\ 4\ 6\dots 2p)(1\ 2p-1\ \dots\ 3)$$

is of order p . Furthermore, $a^2 = 1$ in S_{2p} , so we have

$$(4.29) \quad a^{-1}[a, b]a = a^{-1}a^{-1}b^{-1}aba = b^{-1}aba = (a^{-1}b^{-1}ab)^{-1} = [a, b]^{-1}$$

so $[a, b]$ is not central if $p \geq 3$. In the case $p = 2$ we can instead send $a \mapsto (1\ 2\ 3)$
 $b \mapsto (2\ 3\ 4)$. □

This lemma completes the proof. □

Lemma 4.8. If case (O), if $g = 0$, $n > 0$, and $m = n \geq 3$, then $\pi_1(M)$ is not BO.

PROOF. Recall $c_i^{p_i} = h^{r_i}$. This implies $c_i^{p_i}$ is central, which implies c_i is central (if $\pi_1(M)$ is BO). Then we have three-subcases:

- (a) $m \geq 2$: Kill h , and all the c_i 's except 1, use least to eliminate d_m , kill all remaining d_i 's except d_1 . So we get a quotient

$$(4.30) \quad \langle c, d \mid c^p = 1 \rangle \cong \mathbb{Z} * \mathbb{Z}/p$$

so c is not central.

- (b) $m = 1$, $n \geq 2$: Kill h and eliminate d_1 . Now kill all but 2 c_i 's to get a quotient

$$(4.31) \quad \langle c_1, c_2 \mid c_1^{p_1} = c_2^{p_2} = 1 \rangle \cong \mathbb{Z}/p_1 * \mathbb{Z}/p_2$$

which is non-abelian.

- (c) $m = 0$, $n \geq 3$: Kill h and all but 3 c_i 's. So we get a quotient

$$(4.32) \quad \langle c_1, c_2, c_3 \mid c_i^{p_i} = 1, 1 \leq i \leq 3, c_1c_2c_3 = 1 \rangle = T(p_1, p_2, p_3) .$$

Then c_1 , c_2 , and c_3 central implies $T(p_1, p_2, p_3)$ is abelian.

Lemma 4.9. $T(p_1, p_2, p_3)$ is nonabelian unless $p_1 = p_2 = p_3$, in which case it is $\mathbb{Z}/2 \times \mathbb{Z}/2$.

PROOF. Let $A_1A_2A_3$ be a geodesic triangle with angles $\pi/p_1, \pi/p_2, \pi/p_3$. If the sum

$$(4.33) \quad \frac{1}{p_1} + \frac{1}{p_2} + \frac{1}{p_3}$$

is > 1 this is in the sphere \mathbb{S}^2 , if it is $= 1$ this is in Euclidean space \mathbb{E}^2 , and if this is < 1 , this is the hyperbolic plane \mathbb{H}^2 . Let ρ_{ij} be reflection of the plane ($\mathbb{S}^2, \mathbb{E}^2$, or \mathbb{H}^2) over A_iA_j . Then

$$\gamma_1 = \rho_{13}\rho_{12} \quad \gamma_2 = \rho_{21}\rho_{23} \quad \gamma_3 = \rho_{23}\rho_{12} = (\gamma_1\gamma_2)^{-1} .$$

We can visualize γ_i as rotation about A_i through $2\pi/p_i$. Let $\Gamma(p_1, p_2, p_3)$ be the subgroup of Isom^+ (of whatever plane we are in) generated by γ_1, γ_2 . Since $\gamma_i^{p_i} = 1$, $1 \leq i \leq 3$ and $\gamma_1\gamma_2\gamma_3 = 1$. Therefore $\Gamma(p_1, p_2, p_3)$ is a quotient of $T(p_1, p_2, p_3)$. Clearly $\Gamma(p_1, p_2, p_3)$ is non-abelian unless $p_1 = p_2 = p_3$. Therefore $T(p_1, p_2, p_3)$ is nonabelian unless $p_1 = p_2 = p_3$. □

□

■

APPENDIX A

Orderings of the braid group

We will follow [3].

Let $z_1, \dots, z_n \in \mathbb{D}^2$. A *braid on n strands* is a subset $\beta \subset \mathbb{D}^2 \times I$ such that β is a union of smoothly embedded intervals (called *strands*) in $\mathbb{D}^2 \times I$ such that

Lecture 9 (Hannah Turner); February 18, 2020

- (1) $\beta \cap (D^2 \times \{1\}) = \{(z_1, 1), \dots, (z_n, 1)\}$,
- (2) $\beta \cap (D^2 \times \{0\}) = \{(z_1, 0), \dots, (z_n, 0)\}$,
- (3) $\beta \cap (\mathbb{D}^2 \times \{t\})$ in n points.

We should think of braids as these strands weaving around one another as in fig. 1.

We say two braids are equivalent if there is a deformation from one to the other through braids. There is an operation on braids called *stacking*. This takes two braids and stacks them to make a new braid.






FIGURE 1. A braid on 3 strands.

FIGURE 2. The generator σ_i of B_n .FIGURE 3. The half-Dehn twist about the straight arc connecting z_i and z_{i+1} .

THEOREM A.1 (Artin). *The set of n -strand braids form a group B_n with group operation given by stacking. In particular, it has the following presentation:*

$$(A.1) \quad B_n = \left\langle \sigma_1, \dots, \sigma_{n-1} \mid \begin{array}{l} |i-j| > 1 \implies \sigma_i \sigma_j = \sigma_j \sigma_i, \\ \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \end{array} \right\rangle.$$

Geometrically, the generators σ_i correspond to braids as in fig. 2. Now a braid β is an equivalence class of words in the σ_i .

There is a map $B_n \rightarrow \text{MCG}(D_n)$ from the braid group to the mapping class group of D_n , i.e. the group of orientation preserving homeomorphisms of \mathbb{D}^2 with n punctures such that the punctures are fixed setwise, and $\partial\mathbb{D}^2$ is fixed pointwise. The map sends the generators

$$\sigma_i \mapsto h_{\sigma_i} : D_n \curvearrowright$$

to half-Dehn twists about the straight arc connecting z_i and z_{i+1} . See fig. 3.

CLAIM A.1. This map is an isomorphism.

1. Dehornoy's ordering

DEFINITION A.1. A braid word w is said to be σ -positive (resp. σ -negative) if, among the letters $\sigma_i^{\pm 1}$ that occur in w , the one with lowest index occurs with only positive (resp. negative) exponent, i.e. σ_i occurs but not σ_i^{-1} . In this case we say w is σ_i positive.

REMARK A.1. Usually we don't care for which i the word is σ_i positive. In this scenario we just say ω is σ -positive.

EXAMPLE A.1. $\sigma_1\sigma_2$ and $\sigma_1\sigma_2^{-1}$ are both σ_1 positive. $\sigma_1^{-1}\sigma_2$ is σ_1 -negative.

WARNING A.1. Some braids are neither, e.g. $\sigma_2^{-1}\sigma_3\sigma_2$.

DEFINITION A.2. We say $1 <_{Deh} \beta$ if β is σ -positive.

Note $\beta_1 <_{Deh} \beta_2$ iff $1 <_{Deh} \beta_1\beta_2$.

THEOREM A.2 (Dehornoy). *The above definition for $<_{Deh}$ defines an LO on B_n .*

PROOF IDEA. We use the following properties to prove the theorem.

- Property A (Acyclicity): a σ -positive word is always nontrivial.
- Property C (Comparison): Every nontrivial braid of B_n admits an n -strand representative word that is σ -positive or σ -negative.

Write P_n for the positive braids on n -strands. We will show that P_n is a positive cone.

- (1) P_n is closed: let $\beta_1, \beta_2 \in B_n$. If β_1 is σ_i -positive, β_2 is σ_j positive for $i \leq j$. Then $\beta_1\beta_2$ is σ_i positive. For example:

$$(A.2) \quad \beta_1 = \sigma_1\sigma_2\sigma_3\sigma_2^{-1}$$

$$(A.3) \quad \beta_2 = \sigma_2\sigma_3\sigma_2\sigma_3^{-1}$$

$$(A.4) \quad \beta_1\beta_2 = \sigma_1\sigma_2\sigma_3\sigma_3\sigma_2\sigma_3^{-1}.$$

- (2) $B_n \setminus \{1\} = P_n \cup P_n^{-1}$: property A implies $1 \notin P_n$ and then property C implies this.

- (3) Disjoint union: Suppose $\beta \in P_n \cap P_n^{-1}$. Then $\beta^{-1} \in P_n$, so $\beta\beta^{-1} = 1 \in P_n$ which is a contradiction.

□

Proposition A.3. B_n for $n \geq 3$ is not BO.

PROOF. Define

$$\Delta_n = (\sigma_1 \dots \sigma_{n-1}) (\sigma_1 \dots \sigma_{n-2}) \dots (\sigma_1 \sigma_2) \sigma_1.$$

For example, see fig. 4 for Δ_4 .

CLAIM A.2. $\Delta_n \sigma_i = \sigma_{n-i} \Delta_n$.

Now suppose \prec is a BO on B_n . WLOG $\sigma_1 \prec \sigma_{n-1}$ implies

$$\underbrace{\Delta_n \sigma_1 \Delta_n^{-1}}_{\sigma_{n-1}} \prec \underbrace{\Delta_n \sigma_{n-1} \Delta_n^{-1}}_{\sigma_1}$$

so $\sigma_{n-1} \prec \sigma_1$, so

$$\Delta_n \sigma_i \Delta_n^{-1} = \sigma_{n-i} \Delta_n \Delta_n^{-1} = \sigma_{n-i}$$

which is a contradiction. □


REMARK A.2. (1) For each n , two elements of $(B_n, <_{Deh})$ can be compared in polynomial time (in the length of words).

- (2) This ordering has applications to knot theory. If $\beta \in B_n$ and $\beta < \Delta_n^{-6}$ or $\beta > \Delta_n^g$, then its closure $\hat{\beta}$ is prime.

DEFINITION A.3. (1) An LO group $(G, <)$ is *Conradian* if for all $g, h > 1$, there is some $p \in \mathbb{Z}^+$ with $h < gh^p$.

- (2) $(G, <)$ is *Archimedean* if for all $g, h > 1$, there is $p \in \mathbb{Z}^+$ with $g < h^p$.

Proposition A.4. $(B_n, <_{Deh})$ is not Conradian nor Archimedean.

FIGURE 4. The braid Δ_4 .

2. Nielsen-Thurston orderings on B_n

DEFINITION A.4. Suppose $G \curvearrowright \mathbb{R}$ by orientation preserving homeomorphisms and there is $x \in \mathbb{R}$ with $\text{Stab}_G(x) = \{1\}$. Then $(G, <_x)$ is defined by declaring $g <_x g'$ iff $g(x) <_{\mathbb{R}} g'(x)$.

REMARK A.3. (1) This is an LO since $G < \text{Homeo}^+(\mathbb{R})$.
 (2) Using $y \in \mathbb{R}$, $y \neq x$ could give a different ordering.

The goal is to get an action $B_n \curvearrowright \mathbb{R}$.

We can give D_n a hyperbolic metric. \widetilde{D}_n is a subset of \mathbb{H}^2 . Now compactify \mathbb{H}^2 by adding S^1_∞ . Compactify \widetilde{D}_n by adding in limit points of lifts of ∂D_n . This is a closed disk \widetilde{D}_n . $\partial \widetilde{D}_n$ has two types of points:

- (1) limit points, and
- (2) arcs which cover ∂D_n .

Now pick a basepoint \star . For each $b \in B_n$, take $\beta \mapsto h_\beta : D_n \curvearrowright$. Note that h_β has many lifts in \widetilde{D}_n . Pick one \tilde{h}_b that fixes the basepoint. Now since $\partial \widetilde{D}_n \setminus \{\star\} \cong \mathbb{R}$, we can restrict \tilde{h}_β to $\partial \widetilde{D}_n \setminus \{\star\}$ to get an action on \mathbb{R} . Then it turns out this is all well-defined.

DEFINITION A.5. An LO $<$ on B_n is of *Nielsen-Thurston type* if there is some $x \in \mathbb{R}$ such that for all $\beta, \beta' \in B_n$ $\beta < \beta'$ iff $\beta(x) <_{\mathbb{R}} \beta'(x)$.

FACT 5. (1) Some choices $x \in \mathbb{R}$ have non-trivial stabilizer. These cannot give an ordering.
 (2) Some choices $x \neq y \in \mathbb{R}$ give the same ordering.
 (3) Uncountably many of them are distinct.

3. Isolated orderings

Recall LO's on G correspond to positive cones.

DEFINITION A.6. An ordering $<$ in $\text{LO}(G)$ is *finitely determined* if there is a finite subset $S = \{g_1, \dots, g_k\} \subset G$ such that $<$ is the unique LO on G such that S is positive.

- EXAMPLE A.2. (1) $(\mathbb{Z}, <)$ is determined by choosing $\{1\} \subset P$.
 (2) If $P \subset G$ is finitely generated as a semi-group then the order $<$ determined by P is finitely determined.
 (3) $K = \langle a, b \mid aba^{-1} = b^{-1} \rangle$ is determined by $\{a, b\}$.

Proposition A.5. *A points in $\text{LO}(G)$ is isolated iff $<$ is finitely determined.*

PROOF. (\Leftarrow): Suppose that $< \in \text{LO}(G)$ is finitely determined by f_1, \dots, f_m . Recall $\text{LO}(G) \subset \{0, 1\}^G$. A basis for the topology is given by sets of the form:

$$(A.5) \quad B = \left\{ \left(\underbrace{g_1, \dots, g_k}_{\text{yes}}, \underbrace{h_1, \dots, h_l}_{\text{no}}, \underbrace{\dots}_{\text{whatever}} \right) \right\} \cap \text{LO}(G) .$$

Now we can impose that

- (1) The set of $g \in G$ which we say “yes” to is closed,
- (2) never say “yes” to both g and g^{-1}
- (3) never say “no” to g and g^{-1} .

Then for

$$(A.6) \quad U = \{(f_1, \dots, f_m, f_1^{-1}, \dots, f_m, \dots)\}$$

there is no other order inside U , so $<$ is isolated.

(\Rightarrow): Assume $< \in \text{LO}(G)$ is isolated. There is an open set U such that $<$ is the only element of $\text{LO}(G)$. Write $< \in B \subset U$ where B is of the form (A.5). Then

$$(A.7) \quad P \supset \{g_1, \dots, g_k, h_1^{-1}, \dots, h_l^{-1}\}$$

so $<$ is finitely determined. \square

DEFINITION A.7 ([4]). Let P_{DD} be the set of $\beta \in B_3$ such that β is σ_1 -positive or σ_2 -negative.

THEOREM A.6. P_{DD} is a positive cone, and is generated as a semigroup by $\sigma_1\sigma_2$ and σ_2^{-1} .

PROOF. We will assume that a σ_i -positive word is never trivial. We will also assume that either β is σ_1 -positive or σ_1 -negative or σ_1 -free. Note that this implies σ_1 -free braids are always σ_2 -positive or σ_2 -negative.

Now we show P_{DD} is a positive cone. Write $Q = \langle \sigma_1\sigma_2, \sigma_2^{-1} \rangle$. This is a semigroup. Write $\beta_1 = \sigma_1\sigma_2$ and $\beta_2 = \sigma_2^{-1}$. It is immediate that $Q \subset P_{DD}$. Now we show the opposite. We have two cases:

- Case 1. β or β^{-1} is σ_2 -positive: Then $\beta = \sigma_2^p$ for some $p \in \mathbb{Z} \setminus \{0\}$. For $p > 0$ we have $\beta^{-1} \in Q$, and for $p < 0$ we have $\beta \in Q^{-1}$.
- Case 2. β is σ_1 -positive: then there are $m_i \in \mathbb{Z}$, $1 \leq i \leq k$, such that

$$(A.8) \quad \beta = \sigma_2^{m_1} \sigma_1 \sigma_2^{m_2} \sigma_1 \dots \sigma_1 \sigma_2^{m_k}$$

$$(A.9) \quad = \beta_2^{P_1} \beta_1 \beta_2^{P_2} \beta_1 \dots \beta_1 \beta_2^{P_k}$$

for some $P_i \in \mathbb{Z}$. Then we have

$$(A.10) \quad \beta_2 \beta_1^2 \beta_2 = \beta_1$$

so we can cancel things and keep replacing β_1 by this, until all exponents of β_2 are positive, so $\beta \in Q$.

Case 3. β is σ_1 -negative: so β^{-1} is σ_1 -positive, so $\beta^{-1} \in Q$ by case 2.

Then this means $<_{DD}$ is an ordering on B_n , so it is isolated in $\text{LO}(G)$. \square

APPENDIX B

Orderability and knot groups

A *smooth knot* in S^3 is a (smooth) embedding $K : S^1 \hookrightarrow S^3$. The *knot complement* of K is

$$(B.1) \quad X_K := S^3 \setminus \text{int}(\nu(K)) .$$

The *knot group* of K is $\pi_1(X_K) =: \pi_1(K)$.

Proposition B.1. $H_1(X_K) \cong \mathbb{Z}$.

PROOF. The idea is to use Mayer-Vietoris with $\nu(K)$ and X_K . This gives us the sequence

$$\underbrace{H_2(S^3)}_{=0} \rightarrow \underbrace{H_1(\nu(K) \cap X_K)}_{=\mathbb{Z} \oplus \mathbb{Z}} \rightarrow \underbrace{H_1(\nu(K))}_{=\mathbb{Z}} \oplus H_1(X_K) \rightarrow \underbrace{H_1(S^3)}_{=0}$$

so the result follows from exactness. \square

THEOREM B.2. *Suppose M is a prime orientable three-manifold with $\pi_1(M)$ finitely generated. Then $\pi_1(M)$ is locally indicable iff $\text{rank } H_1(M) \geq 1$.*

Recall if $\pi_1(M)$ is BO then $\pi_1(M)$ is locally indicable, which implies $\pi_1(M)$ is LO.

Corollary B.3. *If $\text{rank } H_1(M) \geq 1$ then $\pi_1(M)$ is LO.*

Corollary B.4. *Knot groups are LO.*

1. Generalized torsion

An element g in a group G is a *generalized torsion element* if and only if

$$\alpha_1^{-1} g^{n_1} \alpha_1 \alpha_2^{-1} g^{n_2} \alpha_2 \dots \alpha_k^{-1} g^{n_k} \alpha_k = 1$$

for some $\alpha_1, \dots, \alpha_n \in G$ and $n_1, \dots, n_k \in \mathbb{Z}^+$. As it turns out, if G has a generalized torsion element, then G is not BO.

EXAMPLE B.1. Consider the Klein bottle group $\langle a, b \mid a^{-1}bab = 1 \rangle$. The element b is a generalized torsion element.

REMARK B.1. There are non BO groups without generalized torsion.

The following is open: $\pi_1(M)$ is BO iff $\pi_1(M)$ has no generalized torsion.

A *torus knot* is a knot in S^3 which embeds in a Heegaard torus. So this is some simple closed curve on the torus. We know that these are parameterized by some rational number

$$\frac{p}{q} \in \mathbb{Q} \cup \left\{ \frac{1}{0} \right\} .$$

Write $T_{p,q}$ for the associated knot. Note that $T_{p,q}$ is the unknot iff $|p| = 1$ or $|q| = 1$.

EXERCISE B.5. $\pi_1(T_{p,q}) = \langle a, b \mid a^p = b^q \rangle$.

Proposition B.6. *If $T_{p,q}$ is nontrivial, then $\pi_1(T_{p,q})$ has generalized torsion.*

PROOF. Assume $p, q > 1$. Write $[x, y] = x^{-1}y^{-1}xy$. Note the following identities:

$$\begin{aligned} \text{(B.2)} \quad & [x^n, y] = x^{-1} [x^{n-1}, y] x [x, y] \\ \text{(B.3)} \quad & [x, y^n] = [x, y] y^{-1} [x, y^{n-1}] y . \end{aligned}$$

□

EXERCISE B.7. $[a^p, b^q]$ is a product of conjugates of $[a, b]$.

$[a, b] \neq 1$, but $[a^p, b^q] = 1$ so $[a, b]$ is a generalized torsion element.

Corollary B.8. $\pi_1(T_{p,q})$ is not BO.

Corollary B.9. G locally indicable does not imply G is BO.

2. Knot groups as extensions

Let $Y := [\pi_1(K), \pi_1(K)]$. Since $H_1(X_K) \cong \mathbb{Z}$ we have a short exact sequence

$$\text{(B.4)} \quad 1 \otimes Y \rightarrow \pi_1(K) \xrightarrow{\rho} \mathbb{Z} \rightarrow 1 .$$

Let $\mu \in \rho^{-1}(1)$. Define

$$\varphi_\mu \in \text{Aut}(Y)$$

by

$$y \mapsto \mu^{-1}y\mu .$$

EXERCISE B.10. $\pi_1(K)$ is BO iff there is an order on Y invariant under φ_μ .

Proposition B.11. $\pi_1(K_1 \# K_2)$ is BO iff $\pi_1(K_1)$ is BO and $\pi_1(K)$ is BO.

The lower central series is as follows. Define $Y_1 = Y$, and

$$\text{(B.5)} \quad Y_n = [Y_{n-1}, Y]$$

for $n > 1$. Notice that Y_n/Y_{n+1} is abelian. Define $\overline{Y_n}$ to be the preimage of $\text{Tor}(Y/Y_{n+1})$ under the quotient. Write $A_n := \overline{Y_n}/\overline{Y_{n+1}}$.

FACT 6. (1) $\overline{Y_b}/\overline{Y_{n+1}}$ is a torsion free abelian group.
(2) $\overline{Y_n}$ are characteristic.

This implies that φ_M induces a well-defined

$$\varphi_n \in \text{Aut}(\overline{Y_n}/\overline{Y_{n+1}}) .$$

A group G is *nilpotent* if $G_n = \{1\}$ for some n .

EXERCISE B.12. Y is residually torsion-free nilpotent if and only if

$$\bigcap_n \overline{Y_n} = \{1\} .$$

Proposition B.13. *If Y is residually torsion-free nilpotent and there are orders $<_n$ on each quotient A_n invariant under φ_n then $\pi_1(K)$ is BO.*

PROOF. We know

$$(B.6) \quad \bigcap_n \overline{Y_n} = \{1\}$$

so for $y \in Y \setminus \{1\}$ there is a unique $n(y)$ such that $y \in \overline{Y_n}$ and $y \notin \overline{Y_{n+1}}$, so

$$[y]_{n(y)} \in A_n$$

is not 0. We have positive cones $P_n \subset A_n$ invariant under φ_n . Now write

$$(B.7) \quad P = \left\{ y \in Y \mid y \neq 1, [y]_{n(y)} \in P_{n(y)} \right\}.$$

(1) $Y = P \amalg P^{-1} \amalg \{1\}$ is clear.

(2) Let $y_1, y_2 \in P$, $n_i := n(y_i)$. If $n_1 < n_2$ then $y_1, y_2 \in \overline{Y_{n_1}}$. Then

$$[y_1, y_2]_{n_1} = [y_1]_{n_1} + \cancel{[y_2]_{n_2}} = [y_1]_{n_1} \in P_{n_1}.$$

The case that $n_1 > n_2$ is similar. If $n_1 = n_2$, then

$$[y_1, y_2]_{n_1} = [y_1]_{n_1} + [y_2]_{n_1=n_2} \in P_n$$

Therefore $y_1 y_2 \in P$. So this shows us that this is an LO. Now we want to see it is a BO.

(3) Let $p \in P$, $y \in Y$. Since $p \in \overline{Y_{n(p)}}$ we have that $[p, q] \in \overline{Y_{n(p)+1}}$. Now

$$y^{-1} p y = p p^{-1} y^{-1} p y = p [p, y]$$

so

$$[y^{-1} p y]_{n(p)} = [p]_{n(p)} + \cancel{[p^{-1} y^{-1} p y]}$$

so this is in $P_{n(p)}$, so $y^{-1} p y \in P$.

(4) We know $n(\varphi_\mu(p)) = n(p)$, so

$$[\varphi_\mu(p)]_{n(p)} = \varphi_n([p]_{n(p)}) \in P_n$$

so $\varphi_n(p) \in P$.

Therefore by an earlier proposition $\pi_1(K)$ is BO. □

So we have orders A_n invariant under φ_n . Now define

$$(B.8) \quad V_n := \mathbb{Q} \otimes_{\mathbb{Z}} A_n \quad L_n := \text{id}_{\mathbb{Q}} \otimes_{\mathbb{Z}} \varphi_n.$$

Notice that this is a vector space and a linear map on it.^{B.1}

Lemma B.14. *L_n preserves an order on V_n if and only if every irreducible factor of the characteristic polynomial $\text{ch}(L_n)$ has a real positive root.*

Lemma B.15. *There is an embedding $V_n \xhookrightarrow{\iota} V_1^{\otimes n}$ such that*

$$(B.9) \quad \begin{array}{ccc} V_n & \xhookrightarrow{\quad} & V_1^{\otimes n} \\ \downarrow L_n & & \downarrow L_1^{\otimes n} \\ V_n & \xhookrightarrow{\iota} & V_1^{\otimes n} \end{array}.$$

Proposition B.16. *If $\text{ch}(L_1)$ has all real positive roots then there are orders on the A_n invariant under the φ_n .*

^{B.1}Which is of course a mathematician's bread and butter.




FIGURE 1. The figure-eight knot.

So by definition $L_1 = \text{id}_{\mathbb{Q}} \otimes \varphi_1$, where φ_1 is the automorphism of Y/Y_2 induced by φ_μ which is conjugate to a scalar multiple of the action on $H_1(\tilde{X}, \mathbb{Q})$ induced by the meridian. Then $\text{ch}(L_1)$ is a scalar multiple of the Alexander polynomial $\Delta_K(t)$.

THEOREM B.17. *If Y is residually torsion-free nilpotent and $\Delta_K(t)$ has all real positive roots then $\pi_1(K)$ is BO.*

EXAMPLE B.2. Consider the figure-eight knot as in fig. 1. This has Alexander polynomial

$$(B.10) \quad \Delta_K(t) = t^2 - 3t + 1.$$

Then $Y_K \cong F_2$, and free groups are residually torsion-free nilpotent. So $\pi_1(K)$ is BO.

Bibliography

- [1] Steven Boyer, Dale Rolfsen, and Bert Wiest, *Orderable 3-manifold groups*, Ann. Inst. Fourier (Grenoble) **55** (2005), no. 1, 243–288. MR2141698
- [2] Adam Clay and Dale Rolfsen, *Ordered groups and topology*, Graduate Studies in Mathematics, vol. 176, American Mathematical Society, Providence, RI, 2016. MR3560661
- [3] Patrick Dehornoy, Ivan Dynnikov, Dale Rolfsen, and Bert Wiest, *Ordering braids*, Mathematical Surveys and Monographs, vol. 148, American Mathematical Society, Providence, RI, 2008. MR2463428
- [4] T. V. Dubrovina and N. I. Dubrovin, *On braid groups*, Mat. Sb. **192** (2001), no. 5, 53–64. MR1859702
- [5] Martin Hertweck, *A counterexample to the isomorphism problem for integral group rings*, Ann. of Math. (2) **154** (2001), no. 1, 115–138. MR1847590
- [6] Anton A. Klyachko, *The structure of one-relator relative presentations and their centres*, J. Group Theory **12** (2009), no. 6, 923–947. MR2582059
- [7] Hellmuth Kneser, *Geschlossene flächen in dreidimensionalen mannigfaltigkeiten.*, Jahresbericht der Deutschen Mathematiker-Vereinigung **38** (1929), 248–259.
- [8] R. H. Lagrange and A. H. Rhemtulla, *A remark on the group rings of order preserving permutation groups*, Canad. Math. Bull. **11** (1968), 679–680. MR240183
- [9] Ciprian Manolescu, *Lectures on the triangulation conjecture*, Proceedings of the Gökova Geometry-Topology Conference 2015, 2016, pp. 1–38. MR3526837
- [10] Stephen H. McCleary, *Free lattice-ordered groups represented as o -2 transitive l -permutation groups*, Trans. Amer. Math. Soc. **290** (1985), no. 1, 69–79. MR787955
- [11] J. Milnor, *A unique decomposition theorem for 3-manifolds*, Amer. J. Math. **84** (1962), 1–7. MR142125
- [12] Edwin E. Moise, *Affine structures in 3-manifolds. V. The triangulation theorem and Hauptvermutung*, Ann. of Math. (2) **56** (1952), 96–114. MR48805
- [13] C. D. Papakyriakopoulos, *On Dehn’s lemma and the asphericity of knots*, Ann. of Math. (2) **66** (1957), 1–26. MR90053
- [14] Grisha Perelman, *The entropy formula for the ricci flow and its geometric applications*, 2002.
- [15] ———, *Finite extinction time for the solutions to the ricci flow on certain three-manifolds*, 2003.
- [16] ———, *Ricci flow with surgery on three-manifolds*, 2003.
- [17] Tibor Radó, *über den begriff der riemannschen fläche*, Acta Litt. Sci. Szeged **2** (1925), 101–121.
- [18] Adam S. Sikora, *Topology on the spaces of orderings of groups*, Bull. London Math. Soc. **36** (2004), no. 4, 519–526. MR2069015
- [19] J. H. C. Whitehead, *On 2-spheres in 3-manifolds*, Bull. Amer. Math. Soc. **64** (1958), 161–166. MR103473