

Breast Cancer ER positive

Firstly, as with any cancer, get to the root cause and remove it.

Stress

Mycotoxins

Gut parasites

Toxins. What you inhale and put onto your body goes INTO your body

Spike protein

Bottled water

Processed food and refined sugars

Lack of fruit and vegetables

No exercise

No detoxing

COVID detox ongoing

The spike protein either from having the virus or from the vaccine can lead to long COVID symptoms and an inflammatory response in the body.

This inflammatory response as in the growth of new blood vessels and hyper progression, is being seen in a lot of cases in the liver.

I, Jane McLelland and Drs around the world are seeing turbo cancers and inflamed lymph nodes, especially in breast cancer patients and hyper progressions in the liver.

If you contract COVID, ensure that you focus on detox, blood clotting and inflammation. I will include information,

It is wise to add detoxing agents into the protocol.

These are:

Bromelain 500mg daily

This is an anti-coagulant. Anti-inflammatory. Proteolytic.

Curcumin 500mg 2 x daily

Antiviral. Anti-fibrotic. Anti-inflammatory.

Nattokinase 2000fu 2 x daily

Anti- atherosclerotic. Proteolytic. Thrombolytic.

Written by Professor Ian Brighthope Medical Practitioner and
Agricultural Scientist. Co-founder of Australasian Integrative
Medicine Association.

Also look at Serrapeptase 240,000 SPU and Lumbrokinase
Nutricology as directed on the label.

Ivermectin 1mg per kg 3 days on and 4 days off.

Vitamin D3 and K2

Research

Cannabis Health Radio for survivor stories. Especially co-host Corrie Yelland who cleared her scc anal cancer with cannabis oil.

Yestolife for support and their interviews on www.healthradio.com

Facebook Groups

Healing Cancer Study Support Group run by Abbey Mitchell

Maria Wessling Bachteal to connect to her breast cancer group

Patient Led Oncology Trials Group run by Mark Sean Taylor

Books.

Dr Neil McKinney's Integrative Oncology Edition 4

Dr Neil McKinney Cancer Unravelling (Latest Book). Still worth buying the above book as well.

The Metabolic Approach by Dr Nasha Winters

The Cancer Resolution by Mark Lintern

Clinics to research

Chemothermia in Istanbul, Turkey

Dr Vogl in Frankfurt. TACE chemotherapy. Having great results with multiple liver lesions.

Mediterranea Medica in Valencia

Key naturopathic breast cancer supports to consider. With kind permission of Dr Neil McKinney from his wonderful book Integrative Oncology Edition 4.

Alpha Lipoic Acid R Form.

Boosts DCA effectiveness at 300mg 2 x daily. Recycles vitamin C. Inhibits NFkB. Chelates heavy metal. (Can't be used with Artemisinin). Reduces lactate to increase apoptosis.

Artemisinin is a hormone balancer. It reduces estrogen and prolactin in breast cancer. It also induces apoptosis. See document with email for instructions. THIS IS A PROTOCOL IN ITSELF.

Boswellia Serrata

Reduces inflammation.

Co-Enzyme Q10 Ubiquinol form 100mg daily to regulate apoptosis via mitochondrial support.

Doxycycline

Targets stem cells in breast cancer by inhibiting mitochondrial ribosomes.

Fresh ground flaxseed has been shown to slow down breast cancer growth. It binds estrogen in the bowel to stop upregulation. It impedes the spread into the lymph nodes. 2 tablespoons a day is a suggested dose.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5808339/>

Gamma Linolenic Acid (GLA)

It decreases ornithine carboxylase activity in breast tumours. It is found in Evening Primrose Oil. Suggested dosing is around 2g daily. It is also a hormone regulator. It is synergistic with Paclitaxel and can increase response to Tamoxifen.

Grapeseed Extract OPCs

Natural aromatase inhibitor.

Green Tea EGCG

Reduces risk of recurrence by 50% of post-operative stage I and II breast cancer. Inhibits breast cancer cells via the aryl hydrocarbon receptor. 2000mg daily is a suggested dose and should be supported with 400iu Gamma Vitamin E with min 10% tocopherols to protect the liver and kidneys.

Indole -3 Carbinol 13C and DIM suppress estrogen activity in ER+ breast cancer. It reduces the risk of recurrence. It inhibits STAT-3 which is involved in breast cancer metastasizing to the bones. 5000iu D3 and MK-7 form of K2 200-400mcg can help to divert calcium to the bones to protect them.

Indole also inhibits urokinase uPA, which is associated with breast cancer growth and metastasis. Typical Indole dosing 400-600mg daily with food split over 2 doses. DIM is taken at bedtime 200-400mg.

Intravenous Artesunate followed by Intravenous Vitamin C Dosing as per clinic.

Iodine reduces estrogen receptor expression. I use Lugol's 2% solution. 2 drops daily in juice.

It regulates the thyroid, increases apoptosis, reduces harm from chemo and reduces resistance to chemo.

Dr Max Gerson stated that the thyroid is important for clearing toxins from the body and is therefore a candidate for regulating autophagy and clearance of dead cells. A typical therapeutic dose is 2-4mg daily. Lugol's 5% solution contains 2.5mg per drop.

Caution with breast feeding as it will pass to breast milk and can cause weight loss for the baby.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC314438/>

Low Dose Naltrexone is particularly supportive with breast cancer. Doses are titrated up to 4.5mg. The prescriber will advise on the starting dose and increments.

Dickson Chemist are specialists in LDN and offer private prescriptions subject to telephone consultation. They ship off label drugs internationally.

<https://ldnresearchtrust.org/>

<https://dicksonchemist.co.uk/new/>

Melatonin is of huge benefit in breast cancer, particularly ER+ where melatonin levels are at their lowest. It decreases production of the ER receptors and is synergistic with Tamoxifen.

Milk thistle 2 capsules or 1 dropperful tincture 3 times daily restricts the growth factor active in 85% of cancers. It heals liver injury, prevents chemical damage to the liver and is a mild detoxifier. If you can access Vitazan milk thistle, Dr McKinney says this can be a life saver in liver failure and is a great detoxifier.

<https://vitazan.com/product/milk-thistle-formulation/>

UK version

[New Roots Herbal Liver 90's – Approved Vitamins](#)

Omega 9

Squalene and Omega 9 found in cold-pressed and extra virgin olive oil are shown to protect from breast cancer. Pour neat onto salads.

Pomegranates inhibit 17-estradiol growth factor in breast cancer. They are also natural aromatase inhibitors.

Quercetin is very supportive in breast cancer treatment. It activates the receptor AhR in dependent 'breast cancer resistance protein' BCRP, making cells more responsive to treatment.

It is supported by Resveratrol, Indole -3 Carbinol 13C and curcumin. It downregulates P53 protein. It also reduces HER2 signalling, which in turn EGFR.

It also binds type 2 estrogen receptors to inhibit growth.

Reishi mushrooms suppress cell adhesion and metastases in breast cancer.

Resveratrol blocks estrogen DNA adducts responsible for breast cancer. It is a phytoestrogen, which means it binds to the estrogen receptor. Dr McKinney says that safety in post-menopausal ER+ women has yet to be determined, so please research further. There is a misconception that phytoestrogens are bad; many are hormone regulators.

Sodium Selenite 300mg daily. Repairs BRAC-2 DNA mutations. A good support for lymphedema as a preventative and a support. Yeast form of selenium is suggested, however, selenomethionine form IS NOT. Acupuncture and grapeseed extract are also good supports.

Soy Isoflavones

There is a misconception that soy should be avoided with estrogen dependent cancers. Dr McKinney's guidance is to stay at around 60mg a day, no higher, as again high doses can become estrogenic. He confirms that it is safe in both pre and post-menopausal women.

Natural soy foods are synergistic with Tamoxifen. Soy intake can reduce the risk of ER+ breast cancer by 30%.

White button mushrooms (*Agaricus Bisporus*) taken in powder form (up to 10g daily) is an aromatase inhibitor in hormone dependent breast cancer.

Zinc Regulates angiogenesis.

Bone density

Vitamin D3 with K2 to divert excess calcium to the bones will strengthen them. This is preferred by Dr Neil McKinney over calcium.

Cannabis helps with bone density.

Inhibiting STAT-3 will help to stop bone metastases.

Curcumin

Doxycycline **Prescribed**

EGCG,

Indole 3 Carbinol

Niclosamide **Prescribed**

Microcrystalline hydroxyapatite ossein complex contains bone growth factors. Take away from calcium supplements as they can reduce the absorption of each other.

Strontium is a support to help reduce fractures. Take away from calcium supplements.

Strontium Citrate 680mg at night is Dr McKinney's suggestion for hardening the bones.

Bone pain

MCHA Microcrystalline Hydroxyapatite Ossein Complex can reduce bone pain. 2-3 capsules twice daily. Pure Encapsulations is a good brand.

Aurum, K1-7 Acupuncture and Oxaloacetate can also help.

Iv Curcumin 20-40mg/kg slow infusion.

Luteolin

There are studies which suggest that Luteolin can inhibit bone pain in some cancers, for example lung cancer, where it was shown to inhibit the activation of neurons and NLRP3 inflammasomes in the spinal cord in mouse studies.

Kaempferol

Natural sources: broccoli, berries, walnuts and from quercetin found in onions.

Kaempferol is a bioflavonoid which can support bone marrow through its inhibition of bone marrow mesenchymal stem cells and stimulation of osteoblasts to support bone health.

Cissus Quadrangularis

There are studies that show that this may be helpful in bone health and helping to repair bones. Studies suggest that this herb has steroidal effects which can produce osteoporosis fighting properties as well as converting bone marrow stem cells to osteoblasts.

Bone Metastases

NFkB plays a large role in bone metastases. It increases bone expression of granulocyte-macrophage colony stimulating factor (GM-CSF). Inhibitors of NFkB are Green Tea, Indole 3 Carbinol, Quercetin, Pomegranates and Reishi.

Other pathways to target are IGF-1, TGF- β 1, Interleukins 1,3,6,11,15 and 17 and VEGF.

See pathway list of inhibitors in files.

Phytoestrogens

Good or bad?

The assumption by most oncologists and pharmacists is that ALL phytoestrogens are bad. This is not the case.

Although some *are* estrogenic, many have the capability to **block** estrogen from binding to the receptor proteins on the cell surface.

A study by Ingram et al found that both pre- and post-menopausal women who consumed phytoestrogens regularly had a reduced risk of breast cancer.

Phytoestrogens also have anti-angiogenic properties, inhibit tumour growth, act as aromatase inhibitors, and enhance the effects of radiotherapy treatment.

The main types of anti-oestrogen acting phytoestrogens are lignans, isoflavones and coumestans.

Isoflavones

The three most common isoflavones are genistein, equol and daidzein. These are largely found in soy. Genistein and daidzein can be found in foods such as currants, fava beans, soybeans, Tofu, tempeh, and miso.

Equol on the other hand cannot be found in food. It is produced when diadzein is metabolized in the intestines to create an intestinal bacterium. The right gut microbiome is necessary to produce equol, and not everyone has the ability to produce it. Seaweed consumption is a way to increase production of equol.

When choosing soy products, it is important to select the ‘healthy’ options such as miso, tempeh and tofu. The processed versions of soy, such as soy based vegetarian products and soy milk, contain much higher levels of genistein and this can have the reverse effect of raising estrogen.

Lignans

Sesame seeds and flaxseeds are known to be two of the plant foods with the highest yielding content of lignans. Flaxseeds have been shown to bind both estrogen and testosterone in the bowel and eliminate it from the body. Studies show that best results come from ground flaxseed rather than oil or whole flaxseeds. 1 tablespoon per day.

Coumestans: split peas, pinto beans, lima beans, alfalfa, brussels sprouts, spinach, and red clover; however, the latter comes with a caution, and should only be used as advised by a health practitioner. See below.

Red Clover contains biochanin A. In studies this has shown to inhibit metabolism of the carcinogen benzo(a) pyrene, which suggests it has chemo preventative properties. It is an inhibitor of CYP3A4, so be aware of that as it can increase liver toxicity if taken with any drugs metabolized using the P450 liver enzyme system.

Resveratrol is considered safe when pre-menopausal but not post menopause because of its phytoestrogen properties. There is not enough evidence to suggest it is safe, so best to avoid.

Binding ability

The key factor of all phytoestrogens is their ability to bind to the receptor known as the RBA (Relative Binding Affinity).

The electrical charge, size and shape determines the ability of the receptor to close around them, and as the majority are so weak in estradiol, binding cannot take place.

% of estradiol activity:

Estradiol has an RBA of 100%

Tamoxifen has an RBA of 80%

Biochanin A from soy 0.004

Coumestrol from red clover and alfalfa 0.7-5

Daidzein from soy 0.02-1

Genistein from soy 0.5-45

Resveratrol from Muscadine grapes 0.001-0.05

Aim for no more than 60mg soy isoflavones per day.

Blood Brain Barrier.

Breast Cancer can metastasise to the brain. MRI scans should be performed to monitor this.

The blood brain barrier consists of capillary endothelial cells and a membrane to protect the brain from toxins, which includes chemotherapies and other targeted drugs. The endothelial cells are fused together by proteins which restrict blood components from crossing through the barrier.

Molecules within the barrier are responsible for angiogenesis (blood vessel growth) and signalling.

Membrane proteins within the brain are known as uptake and efflux transporters. Downregulation of efflux transporter proteins can also create a more penetrable blood brain barrier.

Understanding how to break down the barrier and finding supports that will help drugs to be safely delivered through it, are key factors in treating brain metastases, very few chemotherapy drugs cross the blood brain barrier.

The endothelial cells contain transporters such as multi-drug resistant proteins MDRP1-9, ABCG2 (also known as BBRP) and P-Glycoprotein mediated efflux.

Histone protein acylation modulators can impact on MDRP-1 to reduce drug resistance.

Regulators of histone proteins include green tea EGCG, curcumin, quercetin, grape cyanidins, L Carnitine, sulforaphane, artichoke, silymarin from milk Thistle, Genistein and apigenin.

ABCG2 is the protein responsible from stopping drug resistance in breast cancer. It is also known as BCRP.

BCRP can be increased by indole 3 carbinol, green tea, resveratrol (not in post-menopausal women), Quercetin, and fruit and vegetables such as apples, grapes, onions, broccoli, and cabbage.

P-Glycoprotein limits the bioavailability of drugs taken orally. It does this by pumping them back into the lumen (membrane of the intestine). This promotes drug elimination into the urine and the bile and protects organs such as the brain, testes and lymphocytes.

P-Glycoprotein has been recognised as a contributing factor to resistance to drugs such as vinca alkaloids, taxanes, etoposide, lonafarnib, imatinib and topotecan.

P-Glycoprotein (P-gp) pumps cytotoxins such as chemotherapy out of the cells. Regulators of this are Genistein, Curcumin, Quercetin, Apigenin, Carnosol in Rosemary (synergistic with D3) and hypericin. These will help to keep drugs in the cells for longer.

Inhibiting P-Glycoprotein transportation has shown to increase drug levels of Paclitaxel.

Acetyl L Carnitine helps to carry meds and supplement through the BB barrier. It must be acetyl form to cross BB barrier.

Grapeseed repairs the BB barrier to help drugs to cross.

Artesunate crosses the blood/brain barrier better than Artemisinin.

DMSO 400mg 4 x daily can help penetrate the BB barrier.

Brain Metastases in Breast Cancer and How Drug Resistance Impacts of Treating Brain Mets.

The blood brain barrier consists of capillary endothelial cells and a membrane to protect the brain from toxins, which includes chemotherapies and other targeted drugs. The endothelial cells are fused together by proteins which restrict blood components from crossing through the barrier.

Molecules within the barrier are responsible for angiogenesis (blood vessel growth) and signalling.

Membrane proteins within the brain are known as uptake and efflux transporters. Downregulation of efflux transporter proteins can also create a more penetrable blood brain barrier.

Understanding how to break down the barrier and finding supports that will help drugs to be safely delivered through it, are key factors in treating brain metastases, very few chemotherapy drugs cross the blood brain barrier.

The endothelial cells contain transporters such as multi-drug resistant proteins MDRP1-9, ABCG2 (also known as BBRP) and P-Glycoprotein mediated efflux.

Histone protein acylation modulators can impact on MDRP-1 to reduce drug resistance.

Regulators of histone proteins include green tea EGCG, curcumin, quercetin, grape cyanidins, L Carnitine, sulforaphane, artichoke, silymarin from milk Thistle, Genistein and apigenin.

ABCG2 is the protein responsible from stopping drug resistance in breast cancer. It is also known as BCRP.

BCRP can be increased by indole 3 carbinol, green tea, resveratrol (not in post-menopausal women), Quercetin, and fruit and vegetables such as apples, grapes, onions, broccoli, and cabbage.

P-Glycoprotein limits the bioavailability of drugs taken orally. It does this by pumping them back into the lumen (membrane of the intestine). This promotes drug elimination into the urine and the bile and protects organs such as the brain, testes and lymphocytes.

P-Glycoprotein has been recognised as a contributing factor to resistance to drugs such as vinca alkaloids, taxanes, etoposide, lonafarnib, imatinib and topotecan.

P-Glycoprotein (P-gp) pumps cytotoxins such as chemotherapy out of the cells. Regulators of this are Genistein, Curcumin, Quercetin, Apigenin, Carnosol in Rosemary (synergistic with D3) and hypericin. These will help to keep drugs in the cells for longer.

Inhibiting P-Glycoprotein transportation has shown to increase drug levels of Paclitaxel.

Elacridar has been used in clinical trials to inhibit both P-gp and BCRP.

The Endothelial cells also contain organic anion transporters (OATS), which remove drugs and toxins from the brain. OATP1A2, OATP1B1, OATP1B3 (found in 50% of breast cancer patients) and OATP2B1 are all over expressed in breast cancer. There are variations for other cancer types.

Overexpression of OATS can increase cellular levels of hormones, such as oestrogen and androgen. They are also found in prostate, lung, pancreatic, colon and bone cancers.

Studies have shown that silencing the DNA Methylation-dependent gene regulates expression of OATS.

DNA Methylation-dependent gene inhibitors: B3 Niacin, COQ10, Vitamin B2 Riboflavin and Niacinamide.

Regulate hormones with melatonin, ground flaxseeds, Indole 3 Carbinol and DIM.

Glucose and amino acids cross the blood brain barrier.

Acetyl L Carnitine is a fat-soluble amino acid. Powder mixed as a drink can help with crossing meds over the blood brain barrier. It MUST be acetyl form as L Carnitine will not cross the blood brain barrier.

Dr Neil McKinney suggests Carnivora, which blocks tumour proteins, diluted in 20% mannitol, which will help it to cross the blood brain barrier. This may be worth researching further as a carrier.

Larger molecules cross by using receptors, transferrin, insulin, iron and lipoprotein.

Studies in breast cancer brain metastases show that COX 2 (Cyclooxygenase) increases permeability of the blood brain barrier by producing prostaglandin (PGE2).

Prostaglandins are fatty acid eicosanoids made from dietary fatty-acids via COX 2, the inflammation pathway.

Prostaglandin production (PGE2) is inhibited with NSAIDs (anti-inflammatory drugs), Quercetin, Corticosteroids, L-Glutamine, Omega 3 DHA with EPA, Bromelain, Gamma Lipoic Acid found in Primrose Oil, Echinacea, Olive Oil, Flaxseed.

70% of breast cancer cells produce aromatase. Levels are directly linked to COX2 expression, which increases prostaglandin.

COX-2 inhibitors

Curcumin

Grapeseed OPGs

Green Tea EGCG

Quercetin

Boswellia

Ginger

Resveratrol

Bilberry
Reishi
Liquorice
Garlic
Feverfew
Rosemary
Green-lipid mussel extract
Bromelain
Omega 3 oils EPA/DHA
Aloe Vera
Zeel
Propolis
Black Seed Thymoquinone

Aromatase

Quercetin
Grapeseed procyanidin B dimers
White button mushrooms
Agaricus bisporus,
Melatonin
Reishi
Green tea EGCG
Pomegranate
Iodine

Invasiveness is increased by production of EGFR (Epidermal Growth Factor), which helps breast cancer cells to stick to the epithelial cells in the brain.

EGFR inhibitors

Milk thistle

Resveratrol

Metformin **Prescribed**

Curcumin

Grapeseed Extract OPCs

Quercetin

Soy Genistein

Berberine

EGCG

Chloroquine/Hydroxychloroquine. This can overcome resistance to EGFR targeted treatments. **Prescribed**

In triple negative breast cancer cells will induce Ang-2 (Angiopoeitin-2) resulting in a leaky blood brain barrier. Ang-2 is regulated via the COX2 inflammatory pathway.

Expression of HER2 is more evident in brain metastases than in primary tumours. HER3 is also overexpressed and together with activation of MAPK pathway increases brain metastases.

HER-2 inhibitors

Quercetin

Aloe Vera

HER-3 SEE EGFR

MAPK

Grapeseed OPCs

Vit C

Ellagic Acid
Pomegranate
Milk Thistle
Oleander
Metformin

Gene testing is suggested. Guardant 360 is often available with private healthcare or at a 50% discounted rate for NHS patients. This must be requested through the oncologist.

Genes that are **over expressed** in breast cancer brain metastases, are ATAD2, BRAF, DERL1 and DNMT3B (**DNA Methyltransferases**).

MiR-372 (MicroRNA) suppresses ATAD2

Indole 3 Carbinol and DIM increase MicroRNA

BRAF

Melatonin
Metformin
Berberine
Quercetin
Curcumin
Exercise

DERL-1 over expression can be regulated via the MAPK and ERK pathways.

MAPK

Grapeseed OPCs

Vit C

Ellagic Acid

Pomegranate

Milk Thistle

Oleander

Metformin

ERK-1 &2

Milk Thistle

Metformin

Chaga Mushrooms

DNA Methyltransferases

EGCG

Genes that are **under expressed** in breast cancer brain metastases are ATM, CRYAB and HSPB2.

CRYAB is an anti-apoptotic (cell death) shock protein. Its role is to negatively regulate the BCL2 family, BAX and Caspase-3. It interacts with the mitochondria preventing apoptosis, which is the aim of chemotherapy drugs.

Whilst an under-expression lowers impact on apoptosis, in TNBC, CRYAB can be over expressed. Gene testing will confirm if you have an over-expression.

Regulation of the VEGF and MEK/ERK pathways are needed to regulate an over-expression.

MEK (Regulated by AMP Kinase)

Metformin
Melatonin
Berberine
Quercetin
Curcumin
Exercise

ERK-1 &2

Milk Thistle
Metformin
Chaga Mushrooms

Breast cancer cells adapt by neuronal changes which allow them to take up GABA and bind with NADPH (Nicotinamide adenine dinucleotide phosphate) to increase tumours cells in the microenvironment in the brain. This causes upregulation of survival genes known as GSTAT5, BCL2L1 and TWIST1 (TNBC). This protects them from the cytotoxic effects of treatments such as chemotherapy.

STAT-5A & 5B

Chaste tree berry

BCL-2

Curcumin

Delphinidins found in blueberries and bilberries

Twist 1

Cannabinoids binding to CB1

Loss of PTEN also impacts of brain metastases, particularly in triple negative breast cancer because the loss will activate the P13K/AKT/mTOR pathway.

Regulating the PTEN pathway will reduce invasion into the brain.

PTEN regulators

Indole 3 carbinol

Quercetin

Isoflavones

PI3/Akt/mTOR

Curcumin

Soy

Sulforaphanes

Green Tea EGCG

Indole 3 Carbinol 13C

Metformin prescribe
Itraconazole prescribed
Niclosamide prescribed

HER2 positive cells depend on cyclin D1 which activates cyclin-dependent kinases CDK4 and CDK6.

Cyclin Inhibitors

Curcumin

Indole 3 Carbinol

DIM

Milk Thistle

Resveratrol

Metformin

Also inhibit akt (separate list)

Promote P21 with green tea, butyrates, vitamin D, zinc and zeolite

MiR-202-3p is downregulated in brain metastases in breast cancer, which in turn impacts on MMP-1, leading to over expression. This plays a key role in cells migrating to the brain by weakening the inter-endothelial junction within the blood brain barrier.

Heavy metals upregulate MMP-1. Chelation of heavy metals could be beneficial.

MMP regulators:

Scutellaria

Baiccalein

Green tea EGCG Polyphenols

Curcumin

Metformin

Resveratrol

Zeel

Hormeel

Digestive Enzymes

Bovine Cartilage

Regulation of COX 2 inflammation pathway will also regulate MMPS.

COX2

Curcumin

Grapeseed OPGs

Green Tea EGCG

Quercetin

Boswellia

Ginger

Resveratrol

Bilberry

Reishi

Liquorice

Garlic

Feverfew
Rosemary
Green-lipid mussel extract
Bromelain
Omega 3 oils EPA/DHA
Aloe Vera
Zeel
Propolis
Black Seed Thymoquinone

Research Reference:

Anti Tumour Actions of Cannabinoids by Professor Burkhard Hinz and Robert Ramer.

Breast Cancer Resistance Protein and P-Glycoprotein in Brain Cancer: Two Gatekeepers Team Up by Sagar Agarwal, Anika M.S. Hartz and Bjorn Bauer.

Naturopathic Oncology Edition 4 by Dr Neil McKinney.

Naturopathic Oncology Edition 4 by Dr Neil McKinney.

P-glycoprotein mediated efflux limits substrate and drug uptake in a preclinical brain metastases breast cancer model by Chris E. Atkins et al.

Progression of the role in signalling pathways and cancers by JunFei Zhang, Jia Liu and LiShan Yang.

Silencing miR-202-3p increases MMP1 and promotes a brain invasive phenotype in metastatic breast cancer cells by Rania Harati, Shirin Hafezi, Aloise Mabondzo and Abdelaziz Tlili.

Tumor-Specific Expression of Organic Anion-Transporting Polypeptides: Transporters as Novel Targets for Cancer Therapy by Veronika Buxhofer-Ausch, Lena Secky and Theresia Thalhammer.

Understanding Patterns of brain metastasis in breast cancer and designing rational therapeutic strategies by Evelyn M. Brosnan and Carey K. Anders.

Cancer Feeding Pathways from fatty acids, glucose, and glutamine.

Glucose:

Glut-1 - Atorvastatin and Quercetin

Insulin - Low GI Diet, Berberine, Chromium Picolinate

PP Pathway- DHEA (Not for hormone sensitive cancers), D mannose

OXPHOS- Berberine, Doxycycline, Metformin, Niclosamide

Aerobic Glycolysis- IV Vit C infusions or 5000/6000mg Vit C Liposomal Ascorbate. Chrysin, 2DG (purchase from yourhealthbasket ship internationally), DCA.

Fatty Acids pathways:

SREBP-1- Berberine, Reishi

Acetate – Berberine

ACLY- Hydroxycitrate

FAS- Metformin, Aspirin, EGCG and Curcumin.

F.A.O. Doxycycline, Melatonin

SREBP-2 Dipyridamole, Omega 3 DHA, Luteolin

Mevlonate- Atorvastatin, Omega 3 DHA, Citrus Bergamot

Glutamine (protein) pathways:

Glutaminolysis- EGCG, Ursolic Acid, Resveratrol, Curcumin

Nucleoside Salvage - (Autophagy) Dipyridamole, Danshen

Macropinocytosis- Hydroxychloroquine, Loratadine

mTOR - Metformin, Berberine, EGCG

Gln OXPHOS - Berberine, Doxycycline, Metformin, Niclosamide.

IGF-1- Metformin, Tamoxifen, Lycopene and EGCG

Signalling pathways for breast cancer.

IDEALLY YOU NEED MORE THAN ONE ITEM PER PATHWAY
BUT I HAVE MARKED AN X IF YOU ARE STRUGGLING
FINANCIALLY.

Angiogenesis

EGCG

Quercetin X

Aromatase

Iodine

Quercetin X

White Button Mushrooms

COX2

Aspirin

CBD

Curcumin X

Grapeseed OPCs

DNA hyper-methylation

Niacinamide

Vitamin D X

E-Cadherin (with CDH1 alterations)

Metformin Prescribed X

MSM

Quercetin

EGFR

Berberine X

EGCG covers multiple pathways as you will see below (try with gamma vitamin E and detox regularly)

Milk Thistle

Quercetin

Estrogen receptors

Indole 3 Carbinol/DIM X

Ground Flaxseed

HER2

Aloe Vera

Quercetin X

IL-6

EGCG

R-Alpha Lipoic Acid with Benfotiamine X

Vitamin D3

Insulin

Berberine X

Chromium Picolinate

Metformin (pharma option)

Insulin Like Growth IGF-1

EGCG

Lycopene

Metformin (pharma option) X

NFkB

Blackseed Oil

Metformin (pharma option)

Niclosamide (pharma option) X

R Alpha Lipoic Acid

P13K/Akt/mTOR

Berberine

Curcumin X

Sulforaphanes

SRC-3

Grapeseed Extract

STAT-3

Doxycycline

EGCG X

Indole 3 Carbinol

Survivin

DHA Omega 3 Oils X

Indole 3 Carbinol

Resveratrol

TNF

Blackseed Oil X

Boswellia

uPA

EGCG

Indole 3 Carbinol (This converts to DIM in the liver) X

YB-1

Curcumin

EGCG X

Indole 3 Carbinol