

NOTES ON EXT

JACOB HEGNA
DRAFT: AUGUST 13, 2018

The following are a collection of notes¹ on basic properties of Ext. We work in the category of R -modules for a commutative unital ring R . However, at first we will introduce derived functors in a more general setting, then specialize to Ext.

CONTENTS

1. Conventions	1
2. Derived functors	1
3. Ext	4

1. CONVENTIONS

In these notes, a *category* \mathcal{A} is a collection of objects $Ob(\mathcal{A})$ and, for any two objects $A, B \in \mathcal{A}$ a collection of maps $\text{Hom}_{\mathcal{A}}(A, B)$. It is usually at this point that notes will mention that this definition often has some set-theoretic issues, and say either

- (1) we ignore all set theoretic issues
- (2) all categories are assumed to be *locally small*, which means all Hom objects are indeed sets, not proper classes

these notes will take the first approach. The Hom sets describe formal arrows from A to B ; we require that there be an identity arrow for every object, and that there is an operation \circ called *composition* which is associative. A *functor* between categories is a map $\mathcal{F} : \mathcal{A} \rightarrow \mathcal{B}$ which takes objects to objects and arrows to arrows. In particular, the mapping of objects and arrows should play nicely, in other words if $f : A \rightarrow B$ in \mathcal{A} , then $\mathcal{F}(f)$ should be a map from $\mathcal{F}(A) \rightarrow \mathcal{F}(B)$. It should finally be a homomorphism under composition.

We will often refer to the notion of an Abelian category. The definition is a bit involved, and the reader is welcome to search it out on their own, but the two most important things to remember about such categories are that

- (1) They possess all kernels, cokernels, products, coproducts, and there exists a zero object.
- (2) Any abelian category \mathcal{A} has a fully faithful embedding into a subcategory of $\text{Mod}(R)$. Note, however, that R need not be commutative, so (for instance) things like tensor products need not make sense in every abelian category. In these notes, if a tensor product is used, the reader should assume that the category in question is (essentially) modules over a commutative ring.

These categories are a standard context for homological algebra.

2. DERIVED FUNCTORS

Definition 2.1. A *cochain complex* in an abelian category \mathcal{A} is a collection of objects $\{A_i\}_{i \in \mathbb{Z}}$ where $A_i = 0$ for $i < 0$, and maps $\delta_i : A_i \rightarrow A_{i+1}$ which satisfy the *cochain condition*, $\delta^2 = 0$.

Date: August 13, 2018.

¹These notes by Peter May served as an invaluable guide to basic facts concerning the topics presented here <http://www.math.uchicago.edu/~may/MISC/TorExt.pdf>

Often it is convenient to consider the cochain complex as a \mathbb{Z}_+ -graded object $A = \bigoplus A_i$ with the obvious map δ . We say that the complex (A, δ) is *exact* if $\ker \delta = \operatorname{im} \delta$. It is *short exact* if it degenerates to the complex $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$, and *long exact* otherwise.

Given a functor \mathcal{F} and a short exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$, it is common to ask when the sequence

$$0 \longrightarrow \mathcal{F}(A) \longrightarrow \mathcal{F}(B) \longrightarrow \mathcal{F}(C) \longrightarrow 0$$

is exact. In general, it is more interesting to study those functors \mathcal{F} for which exactness fails. We say \mathcal{F} is *right exact* if the complex is exact everywhere but $\mathcal{F}(A)$, and *left exact* if it is exact everywhere but $\mathcal{F}(C)$. \mathcal{F} is *exact* if it is both left and right exact.

Example 2.2. An equivalence of categories is exact. The global section functor Γ is left exact. The functor which assigns to a G -module M the group M^G is left exact. In general, if \mathcal{F} is a right adjoint of some functor \mathcal{G} , then it is left exact and \mathcal{G} is right exact.

Remark 2.3. Actually, the previous statements are a consequence of a much deeper fact: left adjoints preserve *all* colimits, and right adjoints preserve *all* limits. This is one of the reasons that in any category enriched over **Set**, products look just like they do set-theoretically. Recall that functors $\mathcal{F} : \mathcal{A} \rightarrow \mathcal{B}$ and $\mathcal{G} : \mathcal{B} \rightarrow \mathcal{A}$ are said to be *adjoint* if

$$\operatorname{Hom}_{\mathcal{B}}(\mathcal{F}(A), B) \simeq \operatorname{Hom}_{\mathcal{A}}(A, \mathcal{G}(B))$$

For $A \in \mathcal{A}$ and $B \in \mathcal{B}$.

Studying the failure of a short exact sequence to be exact after an application of \mathcal{F} has given rise to some particularly interesting mathematics. For example, in algebraic geometry, a particular functor Γ called the “global sections” functor is left exact, and studying why it fails to be right exact gives detailed information about the geometric object underlying the functor. Before introducing derived functors, however, we require the following definition.

Definition 2.4. An object A in an abelian category is *injective* if, for all $f : G \rightarrow A$ and monomorphisms $h : G \rightarrow G'$, there is an f' for which

$$\begin{array}{ccc} & A & \\ \exists f' \nearrow & \uparrow f & \\ G' \xleftarrow{h} G & \xleftarrow{\quad} 0 & \end{array} \quad \text{commutes.}$$

In this context, we say that f' is a lift of h through f .

An abelian category has *enough injectives* if, for any object A , there is a monomorphism $i : A \rightarrow I$ where I is injective. A *monomorphism* is a map which behaves like an injection. More specifically, $i : X \rightarrow Y$ is a monomorphism if and only if commutativity of

$$A \begin{array}{c} \xrightarrow{\alpha} \\ \xrightarrow{\beta} \end{array} X \xrightarrow{i} Y$$

implies that $\alpha = \beta$. Equivalently, one could say that $\ker i = 0$ (whatever this means in an arbitrary abelian category).

Remark 2.5. Consider the exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$. Suppose A is injective. In the diagram used to define injectivity, take $G = A$, f the identity on A , and h the map from $A \rightarrow B$. Injectivity of A is precisely the statement that the short exact sequence splits. If \mathcal{F} is a covariant functor, then splitting of the sequence implies the identity $\mathcal{F}(C) \rightarrow \mathcal{F}(C)$ factors through $\mathcal{F}(B) \rightarrow \mathcal{F}(C)$. If \mathcal{F} is left-exact, then the sequence $0 \rightarrow \mathcal{F}(A) \rightarrow \mathcal{F}(B) \rightarrow \mathcal{F}(C) \rightarrow 0$ is exact.

Now, suppose we wish to develop a (co)homology theory which captures the failure of \mathcal{F} to preserve (right) exactness. We would wish that $H^0(X) = \mathcal{F}(X)$, and we would also want that short exact sequences give rise to long exact sequences in cohomology. More concretely, if $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is exact, then we would want exactness of

$$0 \longrightarrow H^0(A) \longrightarrow H^0(B) \longrightarrow H^0(C) \longrightarrow H^1(A) \longrightarrow H^1(B) \longrightarrow H^1(C) \longrightarrow \cdots$$

Finally, if $0 \rightarrow \mathcal{F}(A) \rightarrow \mathcal{F}(B) \rightarrow \mathcal{F}(C) \rightarrow 0$ is exact, then all the higher cohomology objects of A should vanish. In general, if \mathcal{F} is left exact, we desire a cohomology theory. If it is right exact, we desire a homology theory. For the purposes of these notes, we will primarily be concerned with the cohomological case.

How² could we calculate the value of these objects $H^i(-)$? By fiat, we know that $H^0(X) = \mathcal{F}(X)$. To calculate $H^1(X)$, let us assume first we have a theory which satisfies the long exact sequence of cohomology. If we embed X into an injective object I^0 , we have a short exact sequence

$$0 \longrightarrow X \longrightarrow I^0 \longrightarrow K^1 \longrightarrow 0$$

Taking the long exact sequence, we have that the higher $H^i(I^0)$ terms all vanish (by Remark 1.5), so the sequence degenerates to isomorphisms $H^i(K^1) \cong H^{i-1}(X)$ for $i > 1$. In the $i = 1$ case, we have a sequence

$$0 \longrightarrow \mathcal{F}(X) \longrightarrow \mathcal{F}(I^0) \longrightarrow \mathcal{F}(K^1) \longrightarrow H^1(X) \longrightarrow 0$$

This gives us that $H^1(X) = \mathcal{F}(K^1)/\text{im}(\mathcal{F}(I^0))$, which lets us compute $H^1(X)$ in terms of \mathcal{F} itself. But how do we calculate the higher cohomology of K^1 needed to recover the higher cohomology groups of X ? We should embed K^1 into an injective object I^1 (with cokernel K^2), generating isomorphisms $H^1(K^1) \cong \mathcal{F}(K^2)/\text{im}(\mathcal{F}(I^1))$. Repeating this process, we generate an *injective resolution* of the object X

$$\begin{array}{ccccccc} & & & K^1 & & & \\ & & \nearrow & & \searrow & & \\ 0 & \longrightarrow & X & \longrightarrow & I^0 & \xrightarrow{\quad \quad \quad} & I^1 & \xrightarrow{\quad \quad \quad} & I^2 & \longrightarrow & \cdots \\ & & & & & & \searrow & & \nearrow & & \\ & & & & & & K^2 & & & & \end{array}$$

The resolution itself is the chain complex $0 \rightarrow X \rightarrow I^\bullet$. Note that, essentially by construction, the resolution is exact. Now, calculating the higher cohomology groups is reduced to induction. By definition,

$$H^i(X) = H^{i-1}(K^1) = H^{i-2}(K^2) = \cdots = H^1(K^{i-1}) = \mathcal{F}(K^i)/\text{im}(\mathcal{F}(K^{i-1}))$$

We can simplify this further to remove reference to the K^i objects. As \mathcal{F} is left-exact, the sequence $0 \rightarrow K^i \rightarrow I^i \rightarrow I^{i+1}$ gives rise to an isomorphism $\mathcal{F}(K^i) \cong \ker(\mathcal{F}(I^i))$. Thus, after all this work, we can compute $H^i(X)$ by taking an injective resolution $0 \rightarrow X \rightarrow I^\bullet$ and computing the cohomology of the complex $0 \rightarrow I^\bullet$.

There are many things to check. Primarily, one should ask the following question: if one takes two distinct resolutions of X , does the computation of $H^i(X)$ differ? The answer is given by the following lemma, when one takes $f : X \rightarrow X$ to be the identity.

Lemma 2.6. *Let $\alpha : X \rightarrow I^\bullet$ and $\beta : Y \rightarrow J^\bullet$ be injective resolutions of X and Y , respectively. A map $f : X \rightarrow Y$ induces a map $\tilde{f} : I^\bullet \rightarrow J^\bullet$ which satisfies $\tilde{f} \circ \alpha = \beta \circ f$, and this map is unique up to chain homotopy.*

The lemma and the previous discussion motivate the following definition. Before we make it, fix once and for all an injective resolution for each object of our ambient category.

Definition 2.7. Let \mathcal{F} be a left exact functor of abelian categories, where the domain has enough injectives. The *right derived functors* of \mathcal{F} are the objects formed by taking an injective resolution $0 \rightarrow X \rightarrow I^\bullet$, chopping off X , applying \mathcal{F} to $0 \rightarrow I^\bullet$, and taking cohomology. Symbolically,

$$R^i \mathcal{F}(X) = h^i(I^\bullet) = \frac{\ker(\mathcal{F}(I^i \rightarrow I^{i+1}))}{\text{im}(\mathcal{F}(I^{i-1} \rightarrow I^i))}$$

where $0 \rightarrow X \rightarrow I^\bullet$ is the chosen injective resolution of X .

The functors $R^i \mathcal{F}(-)$ depend on the choice of resolution for each object. But, by Lemma 1.5, different choices yield isomorphic functors.

²The following motivation is inspired shamelessly by <https://mathoverflow.net/questions/1151/sheaf-cohomology-and-injective-resolutions>

Remark 2.8. In the previous work, we (crucially) used the fact if I was injective, then it has no higher cohomology. However, the same game would have worked if we had taken an \mathcal{F} -acyclic resolution J^\bullet of X , where $R^i\mathcal{F}(J^k) = 0$ for $i, k > 0$. This is particularly useful in the case of sheaf cohomology, where often we compute the cohomology groups $H^i(X, \mathcal{F})$ by taking flasque or fine resolutions.

Although we have motivated this definition, we still need to formally prove some results about it.

Theorem 2.9. *The functors $R^i\mathcal{F}(-)$ satisfy the following properties*

- (1) $R^0\mathcal{F}(X) = \mathcal{F}(X)$.
- (2) *Short exact sequences give rise to long exact sequences under $R^i\mathcal{F}$.*

Proof. The kernel of the map $\mathcal{F}(I^0 \rightarrow I^1)$ is the object $\mathcal{F}(X)$. The image of $\mathcal{F}(I^{-1} \rightarrow I^0)$, taking $I^{-1} = 0$, is 0. This proves (1).

Given a short exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$, we take injective resolutions of each object, and the Lemma induces maps between the resolutions. Diagrammatically, we have

$$\begin{array}{ccccccccc}
0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & I_A^0 & \longrightarrow & I_B^0 & \longrightarrow & I_C^0 & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & I_A^1 & \longrightarrow & I_B^1 & \longrightarrow & I_C^1 & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
& & \vdots & & \vdots & & \vdots & &
\end{array}$$

Apply the snake lemma to each row. This gives the desired long exact sequence. Naturality of the snake lemma yields naturality of the induced long exact sequence. \square

Example 2.10. The core example of these notes is the Ext functor. However, note that this definition is central to many aspects of commutative algebra. Cohomology of groups and of sheaves (an algebraic tool to keep track of local data on a topological space) are described explicitly by derived functors.

3. EXT

Let R be a commutative ring with unity. The category $\text{Mod}(R)$ of R -modules is abelian. If $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is an exact sequence of R -modules, the the sequences

$$\begin{aligned}
0 &\longrightarrow \text{Hom}_R(M, A) \longrightarrow \text{Hom}_R(M, B) \longrightarrow \text{Hom}_R(M, C) \\
0 &\longrightarrow \text{Hom}_R(C, N) \longrightarrow \text{Hom}_R(B, N) \longrightarrow \text{Hom}_R(A, N)
\end{aligned}$$

are exact for R -modules M, N . In other words, the bifunctor $\text{Hom}_R(-, -)$ is left exact in both arguments.

Definition 3.1. The functors $\text{Ext}_R^i(A, -)$ (resp. $\text{Ext}_R^i(-, B)$) are defined to be the right derived functors of $\text{Hom}_R(A, -)$ (resp. $\text{Hom}_R(-, B)$).

Note that, in the case of $\text{Hom}_R(-, B)$, taking the right derived functors amounts to taking a projective resolution instead of an injective resolution.

Concretely, the modules $\text{Ext}_R^i(M, N)$ classify extensions of the module N by M of length i . So, $\text{Ext}_R^i(M, N)$ is in bijection with the set of $\{E_j\}$'s for which

$$0 \longrightarrow N \longrightarrow E_1 \longrightarrow \cdots \longrightarrow E_j \longrightarrow \cdots \longrightarrow E_i \longrightarrow M \longrightarrow 0$$

Given extensions $\{E_i\} \in \text{Ext}_R^n(N, P)$ and $\{R_j\} \in \text{Ext}_R^m(M, N)$, one can form an extension of length $m+n$ in $\text{Ext}_R^{m+n}(M, P)$ by

$$\begin{array}{ccccccccccc}
& & & & & & N & & & & \\
& & & & & \nearrow & & \searrow & & & \\
0 & \longrightarrow & P & \longrightarrow & E_1 & \longrightarrow & \cdots & \longrightarrow & E_n & \xrightarrow{\quad\quad\quad} & R_1 & \longrightarrow & \cdots & \longrightarrow & R_m & \longrightarrow & M & \longrightarrow & 0
\end{array}$$

This (informally) describes a pairing

$$\text{Ext}_R^n(N, P) \otimes \text{Ext}_R^m(M, N) \longrightarrow \text{Ext}_R^{m+n}(M, P)$$

called the *Yoneda product*. Observe that if we take $N = M = P$, then we obtain a multiplicative structure on the graded module $\text{Ext}_R^*(M, M)$. This is somewhat analogous to the cup product in singular cohomology. We may return to this later.

Remark 3.2. Let M, N, I, P be R -modules where I is injective and P is projective. The following properties follow from general derived functor nonsense.

- (1) $\text{Ext}_R^0(M, N) = \text{Hom}_R(M, N)$.
- (2) $\text{Ext}_R^i(P, N) = 0$ for $i > 0$.
- (3) $\text{Ext}_R^i(M, I) = 0$ for $i > 0$.

We now begin the setup for the universal coefficient theorem. Note that, induced by the obvious short exact sequences, we have that the following are exact

$$0 \longrightarrow \text{Hom}(B_{n-1}, G) \longrightarrow \text{Hom}(C_n, G) \longrightarrow \text{Hom}(Z_n, G) \longrightarrow 0 \quad (3.1)$$

$$0 \longrightarrow \text{Hom}(H_n, G) \longrightarrow \text{Hom}(Z_n, G) \longrightarrow \text{Hom}(B_n, G) \longrightarrow \text{Ext}^1(H_n, G) \longrightarrow 0 \quad (3.2)$$

Theorem 3.3 (Universal Coefficients). *For a PID R , the sequence*

$$0 \longrightarrow \text{Ext}_R^1(H_{n-1}, G) \xrightarrow{j} H^n(\text{Hom}_R(C_*, G)) \xrightarrow{h} \text{Hom}_R(H_n, G) \longrightarrow 0$$

of R -modules is exact. It splits, although not naturally with respect to the grading of δ .

In the proof, we have omitted the subscripts on Hom and Ext for clarity. Note that the general case of universal coefficients is a spectral sequence, but if R is a PID, then it degenerates to the above short exact sequence.

Proof. Consider the following commutative diagram.³ The middle row is the sequence (2.1), the first column is the sequence (2.2) in degree $n-1$, the third column is the sequence (2.2) in degree n . The top and bottom rows are each sequence (2.1), in degrees n and $n-1$, respectively.

$$\begin{array}{ccccccc}
& & 0 & & & & \\
& & \uparrow & & & & \\
& & \text{Ext}^1(H_{n-1}, G) & & \text{Hom}(C_{n+1}, G) & \xleftarrow{\delta} & \text{Hom}(B_n, G) \xleftarrow{\quad} 0 \\
& & \uparrow & \nearrow j & \uparrow \delta & \nwarrow i^* & \uparrow i^* \\
0 & \longrightarrow & \text{Hom}(B_{n-1}, G) & \xrightarrow{\delta} & \text{Hom}(C_n, G) & \xrightarrow{i^*} & \text{Hom}(Z_n, G) \longrightarrow 0 \\
& & \uparrow i^* & & \uparrow \delta & \nwarrow h & \uparrow q^* \\
0 & \longleftarrow & \text{Hom}(Z_{n-1}, G) & \xleftarrow{i^*} & \text{Hom}(C_{n-1}, G) & & \text{Hom}(H_n, G) \\
& & & & & & \uparrow \\
& & & & & & 0
\end{array}$$

The map j is defined by taking an element $x \in \text{Ext}^1(H_{n-1}, G)$, lifting it to $x_0 \in \text{Hom}(B_{n-1}, G)$, and setting $j(x) = \delta(x_0)$. It is well-defined on cohomology classes because if we had lifted x to x_1 , then $x_0 - x_1$ is in the image of the map from $\text{Hom}(Z_{n-1}, G)$ and thus is 0 in cohomology.

³Proof taken from <http://ckottke.ncf.edu/docs/exttoruct.pdf>

The map h is defined by taking an element $y \in \text{Hom}(C_n, G)$ then taking the preimage of $i^*(y)$ under q^* . It exists, because $i^*(y)$ is in the kernel of i^* by chasing the commutative square.

The composition $h \circ j = 0$ because it takes the two arrows in the middle exact sequence. $\ker h = \text{im } j$ because $h(y) = 0$ implies $i^*(y) = 0$ so we can lift y to $\text{Hom}(B_{n-1}, G)$ and project it onto $\text{Ext}^1(H_{n-1}, G)$. We leave the reader to ponder exactness at the endpoints of the sequence.

The splitting of the sequence is due to the splitting of the middle row. It is not natural because of the failure of naturality of the middle row (it splits between gradings of the chain complex). \square

Remarkably, none of the proof of Universal Coefficients relies on the properties of Hom or Ext outside of general properties of derived functors. Thus, we can rewrite the statement of the theorem to be:

Theorem 3.4. *Let R be a PID, and C_\bullet a chain complex in $\text{Mod}(R)$. If $\mathcal{F}, \mathcal{G} : \text{Mod}(R) \rightarrow \mathcal{A}$ are functors such that \mathcal{F} is contravariant and left exact, and \mathcal{G} is covariant and right exact, then*

$$0 \longrightarrow R^1\mathcal{F}(H_{n-1}(C_\bullet)) \longrightarrow H^n(\mathcal{F}(C_\bullet)) \longrightarrow \mathcal{F}(H_n(C_\bullet)) \longrightarrow 0$$

$$0 \longrightarrow \mathcal{G}(H_n(C_\bullet)) \longrightarrow H_n(\mathcal{G}(C_\bullet)) \longrightarrow L^1\mathcal{G}(H_{n-1}(C_\bullet)) \longrightarrow 0$$

are exact. In particular, if the homology groups of C_\bullet are free, then \mathcal{G} and H_n commute over C_\bullet .