

HOMEWORK 01

MATH 8300

JACOB HEGNA

Problem 1.

- (1) Describe all the isomorphism classes of representations of $\mathbb{C}[x]$ of dimension 1. How many are there?

Proof. The modules $\mathbb{C}[x]/(x - z)$ for $z \in \mathbb{C}$ are all of dimension one (they are isomorphic to \mathbb{C}) and they are not mutually isomorphic for $(x - z) \neq (x - s)$, as the action of x differs between them. Appealing to the structure theorem of modules over a PID, we see that these are indeed the only possible modules of dimension 1.

There are uncountably many such modules of distinct isomorphism classes when considered as $\mathbb{C}[x]$ -modules. Base changing to \mathbb{C} collapses these to a single isomorphism class. \square

- (2) Describe also the isomorphism classes of representations of $\mathbb{C}[X]$ of dimension 2. Can they all be generated by a single element? If not, identify the representations that can be generated by a single element. Are any of these representations of dimension 2 simple?

Proof. Once again, an appeal to the structure theorem says that any such module must be of the form

$$\mathbb{C}[x] \bigoplus_i \frac{\mathbb{C}[x]}{I}.$$

Indeed, such modules of dimension two are necessarily of the form $\mathbb{C}[x]/(x - a) \oplus \mathbb{C}[x]/(x - b)$ for $a, b \in \mathbb{C}$ or $\mathbb{C}[x]/(x - a)^2$ for $a \in \mathbb{C}$. By the Chinese remainder theorem, $\mathbb{C}[x]/(x - a) \oplus \mathbb{C}[x]/(x - b) \cong \mathbb{C}/((x - a)(x - b))$ for a, b distinct. Thus, we reduce our classification to the modules which can be generated by a single element and those which cannot. The former class is of the form $\mathbb{C}[x]/(f)$ for f degree 2. The latter is of the form $\mathbb{C}[x]/(x - a) \oplus \mathbb{C}[x]/(x - a)$. To see this cannot be generated by a single element, suppose that there was such a generator a . Then, \square

Problem 2.

- (1) Let $f \in \mathbb{Q}[x]$ be an irreducible polynomial. Show that every finitely generated module for the ring $A = \mathbb{Q}[x]/(f^r)$ is a direct sum of modules isomorphic to $V_s := \mathbb{Q}[x]/(f^s)$, where $1 \leq s \leq r$. Show that A has only one simple module up to isomorphism. When $r = 5$, calculate $\dim \operatorname{Hom}_A(V_2, V_4)$ and $\dim \operatorname{Hom}(V_4, V_2)$.

Proof. Of course, that every finitely generated module over A is a direct sum of V_s comes immediately from the structure theorem, as we have that the module must be of the form

$$\frac{\mathbb{Q}[x]}{f^r} \bigoplus_i \frac{\mathbb{Q}[x]/(f^r)}{(f^s)/(f^r)},$$

which is isomorphic to $\bigoplus_i V_{s_i}$ for $s_i \leq r$.

That A has only one simple module comes from a more general fact that any (commutative, Noetherian) local ring (R, m) has a unique simple module isomorphic to R/m —to demonstrate this fact, consider the map $A \rightarrow M$ given by fixing a nonzero element a of the simple module M and taking the map to be multiplication by a . Surjectivity is implied by simplicity of M , and thus M is only of length 1 if the kernel of this map is m .

Let $r = 5$. $\operatorname{Hom}_A(V_2, V_4)$ is the collection of maps $A \rightarrow V_4$ which annihilate f^2 . This only occurs if 1 is mapped into (f^2) . The length of $\operatorname{Hom}_A(V_2, V_4)$ is given by considering that the submodules

are classified by which ideal 1 is sent to in the underlying map $A \rightarrow V_4$, of which there are three choices, $0 \subset (f^3) \subset (f^2)$. This gives the desired length: 2. The module $\text{Hom}_A(V_4, V_2)$ is similarly classified, and 1 may be mapped into either of the ideals $0 \subset (f) \subset V_2$, which gives a length of 2 as well. \square

Problem 2. Show that $\mathbb{Q}[x]/((x-1)^5) \cong \mathbb{Q}[x]/((x-2)^5)$ as algebras.

Proof. As \mathbb{Q} -algebras, they are isomorphic by the coordinate change map sending

They are not isomorphic as $\mathbb{Q}[x]$ -algebras, however, as there is no \mathbb{Q} -linear arrow making the following diagram commute:

$$\begin{array}{ccc} & \mathbb{Q}[x] & \\ \swarrow & & \searrow \\ \frac{\mathbb{Q}[x]}{(x-1)^5} & & \frac{\mathbb{Q}[x]}{(x-2)^5} \end{array}$$

This can be seen by \square

Problem 3. Let A be a ring and let V be an A -module.

- (1) Show that V is simple if and only if for all nonzero $x \in V$, x generates V .

Proof. Suppose there is a nonzero element x which does not generate V . What it generates is a nonzero submodule strictly contained in V which is absurd as V is simple. In the other direction, there can be so nonzero submodules of V as all of their elements generate the entirety of V . \square

- (2) Show that V is simple if and only if V is isomorphic to A/I for some maximal left ideal I .

Proof. Fix a nonzero $v \in V$ and consider the map $A \rightarrow V$ which sends a to av . This is surjective by the prior question. The kernel of this map must be maximal, otherwise the maximal ideal containing the kernel would yield a nonzero strict submodule of V as a quotient of A . Thus $V \cong A/I$ for I maximal. \square

- (3) Show that if A is a finite dimensional algebra over a field then every simple A -module is a composition factor of the free rank 1 module ${}_A A$, and hence that a finite dimensional algebra only has finitely many isomorphism classes of simple modules.

Proof. Fix a simple module $M \cong A/m$ for m a maximal left ideal. We write a composition series for ${}_A A$ which begins with the inclusion $m \rightarrow A$ (we will drop the left-module subscript notation, e.g. ${}_A A$, as from now on everything in sight is acted on the left by A), this begins a composition series as $A/m \cong M$ is simple. We may extend this to 0 by observing that ideals are vector subspaces of A , and thus we may inductively choose a maximal subideal of each element of the composition series, which terminates as each step decreases the dimension by 1. By Jordan-Hölder, there are only finitely many such (isomorphism classes of) simple modules, as the composition series is unique up to reordering of the quotient modules. \square

Problem 4.. Let K be a field, and let $Q_2 = y \bullet \xleftarrow{\beta} \bullet x$ be the quiver in the notes with representations $S_x = 0 \xleftarrow{0} K$, $S_y = K \xleftarrow{1} 0$, and $V = K \xleftarrow{1} K$.

- (1) Compute $\dim \text{Hom}_{K(F(Q_2))}(S_x, V)$, $\dim \text{Hom}_{K(F(Q_2))}(V, S_x)$ and $\dim \text{Hom}_{K(F(Q_2))}(V, V)$.

Proof. We consider the following diagram

$$\begin{array}{ccc} K & \xrightarrow{\alpha} & K \\ \downarrow 0 & & \downarrow 1 \\ 0 & \xrightarrow{\beta} & K \end{array}$$

2

The only choice for β is 0. To make things commute, we require $\alpha = 0$ as well, which implies the dimension of the Hom-module is 0. Similarly, we may consider

$$\begin{array}{ccc} K & \xrightarrow{\alpha} & K \\ \downarrow 1 & & \downarrow 0 \\ K & \xrightarrow{\beta} & 0 \end{array}$$

β must be 0, and α can be anything. Thus, the dimension of $\text{Hom}(V, S_x)$ is 1. Finally, we may consider

$$\begin{array}{ccc} K & \xrightarrow{\alpha} & K \\ \downarrow 1 & & \downarrow 1 \\ K & \xrightarrow{\beta} & K \end{array}$$

For which we require $\alpha = \beta$ for commutativity, which yields a dimension of 1 for $\text{Hom}(V, V)$. \square

- (2) Determine whether or not the path algebra $K(F(Q_2))$ is isomorphic to either $K[x]/(x^2)$ or $K[x]/(x^3)$.

Proof. \square

Problem 5. Show that the path algebras of the two quivers $Q_1 = \bullet \rightarrow \bullet \leftarrow \bullet$ and $Q_2 = \bullet \leftarrow \bullet \rightarrow \bullet$ over R are isomorphic to the algebras of 3×3 matrices over R of the form

$$R_1 = \begin{bmatrix} a & b & 0 \\ 0 & c & 0 \\ 0 & d & e \end{bmatrix} \quad \text{and} \quad R_2 = \begin{bmatrix} a & 0 & 0 \\ b & c & d \\ 0 & 0 & e \end{bmatrix}$$

Proof. I claim that $K(F(Q_1)) = R_2$ and $K(F(Q_2)) = R_1$. To see this, we write out the multiplication tables. Write $Q_1 = x \xrightarrow{\alpha} y \xleftarrow{\beta} z$. The multiplication tables are:

$$\begin{array}{c|ccccc} K(F(Q_1)) & 1_x & 1_y & 1_z & \alpha & \beta \\ \hline 1_x & 1_x & 0 & 0 & 0 & 0 \\ 1_y & 0 & 1_y & 0 & \alpha & \beta \\ 1_z & 0 & 0 & 1_z & 0 & 0 \\ \alpha & \alpha & 0 & 0 & 0 & 0 \\ \beta & 0 & 0 & \beta & 0 & 0 \end{array} \quad \text{and} \quad \begin{array}{c|ccccc} R_2 & a & c & e & b & d \\ \hline a & a & 0 & 0 & 0 & 0 \\ c & 0 & c & 0 & b & d \\ e & 0 & 0 & e & 0 & 0 \\ b & b & 0 & 0 & 0 & 0 \\ d & 0 & 0 & d & 0 & 0 \end{array},$$

where we write $i \in \{a, b, c, d, e\}$ for the matrix with 1 in the i th place, where i refers to the coordinate of R_1 indicated in the statement of the question. These algebras are necessarily now the same, as mapping obvious elements together, it is immediate that addition is preserved at $1 = 1_x + 1_y + 1_z \mapsto 1 = a + c + e$.

It is not too hard to see that transposing this multiplication table corresponds to transposing R_1 to R_2 and similarly reversing the direction of α, β (as transposing the multiplication table simply conjugates the endpoints of the path). However, for clarity, we write the multiplication tables below:

$$\begin{array}{c|ccccc} K(F(Q_2)) & 1_x & 1_y & 1_z & \alpha & \beta \\ \hline 1_x & 1_x & 0 & 0 & \alpha & 0 \\ 1_y & 0 & 1_y & 0 & 0 & 0 \\ 1_z & 0 & 0 & 1_z & 0 & \beta \\ \alpha & 0 & \alpha & 0 & 0 & 0 \\ \beta & 0 & \beta & 0 & 0 & 0 \end{array} \quad \text{and} \quad \begin{array}{c|ccccc} R_1 & a & c & e & b & d \\ \hline a & a & 0 & 0 & b & 0 \\ c & 0 & c & 0 & 0 & 0 \\ e & 0 & 0 & e & 0 & d \\ b & 0 & b & 0 & 0 & 0 \\ d & 0 & d & 0 & 0 & 0 \end{array}.$$

\square

Problem 6. Let k be a field. Show that the space of column vectors k^n is a simple module for $M_n(k)$. Show that, as a left module, $M_n(k)$ is the direct sum of n modules each isomorphic to k^n . Show that, up to isomorphism, $M_n(k)$ has only one simple module.

Proof. Let $v \in k^n$ be a nonzero column vector. We consider k^n a vector space with standard basis. As v is nonzero, it has a nonzero coordinate i . Indeed, as k is a field, we may choose a matrix which maps v to the i th basis vector. There exists a linear transformation which sends this basis vector to any other basis vector

in k^n of our liking, which is expressed as an element of $M_n(k)$. This implies that v generates all of k^n and thus k^n is simple.

Let $N_i \subset M_n(k)$ refer to the submodule of matrices which are nonzero outside of the i th column. This is a submodule as it is a subset which inherits the addition and action of $M_n(k)$, and it is closed under this action by the definition of matrix multiplication. There is an obvious isomorphism of $N_i \cong k^n$ by identifying k^n with the i th column of any given matrix. Of course, we have a surjection $M_n(k) \rightarrow \bigoplus_i^n N_i$ which projects the i th column to N_i , and its kernel is zero by definition.

$M_n(k)$ has only one simple module (up to isomorphism) as the composition series

$$0 \subset N_0 \subset N_0 \oplus N_1 \subset \cdots \subset \bigoplus_{i=0}^{n-1} N_i = M_n(k)$$

has composition factors N_i and the i th place, which is simple as they agree with k^n . By Jordan-Hölder, we conclude. \square

Problem 7. Let C_n denote the category with n objects labeled a_1, \dots, a_n and where there is a unique homomorphism $a_i \rightarrow a_j$ for every ordered pair of numbers (i, j) . Show that the category algebra RC_n is isomorphic to the algebra of $n \times n$ matrices $M_n(R)$.

Proof. This is simply unpacking the definitions. The algebra RC_n has, as a generating set, the arrows $a_i \rightarrow a_j =: \beta_{ij}$. The multiplication is defined to be:

$$X = \left(\sum_{i \leq j} b_{ij} \beta_{ij} \right) \cdot \left(\sum_{i \leq j} c_{ij} \beta_{ij} \right),$$

where the coefficient of β_{ij} in X is exactly the sum $\sum_{k=1}^n b_{ik} c_{kj}$, because the product is zero if the morphisms are not composable and the R -inherited product otherwise. If we consider the map of sets $RC_n \rightarrow M_n(R)$ sending β_{ij} to the unit in the ij th coordinate of $M_n(R)$ then extending R -linearly, we find this is precisely the definition of matrix multiplication. \square

Problem 8. Let x be an object of a finite category \mathcal{C} .

- (1) Show that the subset $RC \cdot 1_x$ of the category algebra RC is the span of the morphisms whose domain is x , and that $1_x \cdot RC$ is the span of the morphisms whose codomain is x .

Proof. Of course, this is just the definition of the multiplication in the category algebra. For any $r\beta \in RC$, we have that $(r\beta) \cdot 1_x$ is 0 if $\text{od}(\beta) \neq x$ and $r\beta$ otherwise. This is the definition of the desired span. For the other direction, the argument is the same, but we consider $t(\beta)$ as the multiplication is on the other side. \square

- (2) Show that $RC = \bigoplus_{x \in \text{Ob } \mathcal{C}} RC \cdot 1_x$ as left RC -modules.

Proof. Define a map $\phi : RC \rightarrow \bigoplus RC \cdot 1_x$ by projecting a formal linear sum of morphisms to the direct summand of its domain. By part (1), this is surjective. The only element mapped to the identity is $\sum_{x \in \text{Ob } \mathcal{C}} 1_x$, so it is a bijection. The projection, of course, preserves the addition. So, we only have to check that the map is stable under the action of the ring. Left multiplication by RC preserves the domain of a morphism, so the map is a bijective RC -module homomorphism. \square

- (3) Let $R\text{Hom}_{\mathcal{C}}(x, -)$ denote the functor $\mathcal{C} \rightarrow R\text{-mod}$ that sends an object y to the free R -module with the set of homomorphisms $\text{Hom}_{\mathcal{C}}(x, y)$ as a basis. Under the correspondence between representations of \mathcal{C} over R and RC -modules, show that the functor $R\text{Hom}(x, -)$ corresponds to the left RC -module $RC \cdot 1_x$ and that $R\text{Hom}(-, x)$ corresponds to the right RC -module $1_x \cdot RC$.

Proof. Under the Mitchell-correspondence, the module formed from the representation (functor) given by $\text{Hom}(x, -)$ is precisely the direct sum over the objects x in \mathcal{C} , e.g. $\bigoplus_{y \in \text{Ob } \mathcal{C}} R\text{Hom}(x, y)$. This is, by definition, the R -linear span of all morphisms whose domain is x . By (1), this is $RC \cdot 1_x$. The only thing to check is that the action of RC agrees on each. Of course, by definition, $R\text{Hom}(x, y)$ is acted on by the basis elements of RC by composition, which is the same action of $RC \cdot 1_x$. The same procedure shows that $R\text{Hom}(-, x)$ corresponds to $1_x \cdot RC$. \square