

Abstract Algebra

Module 4

Section 24 : Noncommutative Examples

June 14, 2021

$End(A)$: Noncommutative Ring

$\langle End(A), +, \circ \rangle$ is well-defined only if A is an abelian group.

Let $\theta, \phi, \psi \in End(A)$, $x \in A$

1. Addition is associative. $\theta + (\phi + \psi) = (\theta + \phi) + \psi$

$End(A)$: Noncommutative Ring

$\langle End(A), +, \circ \rangle$ is well-defined only if A is an abelian group.

Let $\theta, \phi, \psi \in End(A)$, $x \in A$

1. Addition is associative. $\theta + (\phi + \psi) = (\theta + \phi) + \psi$
2. Addition is commutative. $\theta + \phi = \phi + \theta$

$End(A)$: Noncommutative Ring

$\langle End(A), +, \circ \rangle$ is well-defined only if A is an abelian group.

Let $\theta, \phi, \psi \in End(A)$, $x \in A$

1. Addition is associative. $\theta + (\phi + \psi) = (\theta + \phi) + \psi$
2. Addition is commutative. $\theta + \phi = \phi + \theta$
3. Existence of Additive Identity.

$$\forall \theta \in End(A), \exists 0 \in End(A), \theta + 0 = \theta$$

$End(A)$: Noncommutative Ring

$\langle End(A), +, \circ \rangle$ is well-defined only if A is an abelian group.

Let $\theta, \phi, \psi \in End(A)$, $x \in A$

1. Addition is associative. $\theta + (\phi + \psi) = (\theta + \phi) + \psi$
2. Addition is commutative. $\theta + \phi = \phi + \theta$
3. Existence of Additive Identity.

$$\forall \theta \in End(A), \exists 0 \in End(A), \theta + 0 = \theta$$

4. Existence of Additive Inverses.

$$\forall \theta \in End(A), \exists \theta' \in End(A), \theta + \theta' = 0$$

$End(A)$: Noncommutative Ring

$\langle End(A), +, \circ \rangle$ is well-defined only if A is an abelian group.

Let $\theta, \phi, \psi \in End(A)$, $x \in A$

1. Addition is associative. $\theta + (\phi + \psi) = (\theta + \phi) + \psi$
2. Addition is commutative. $\theta + \phi = \phi + \theta$
3. Existence of Additive Identity.
 $\forall \theta \in End(A), \exists 0 \in End(A), \theta + 0 = \theta$
4. Existence of Additive Inverses.
 $\forall \theta \in End(A), \exists \theta' \in End(A), \theta + \theta' = 0$
5. Multiplication is associative. $\theta(\phi\psi) = (\theta\phi)\psi$

$End(A)$: Noncommutative Ring

$\langle End(A), +, \circ \rangle$ is well-defined only if A is an abelian group.

Let $\theta, \phi, \psi \in End(A)$, $x \in A$

1. Addition is associative. $\theta + (\phi + \psi) = (\theta + \phi) + \psi$
2. Addition is commutative. $\theta + \phi = \phi + \theta$
3. Existence of Additive Identity.

$$\forall \theta \in End(A), \exists 0 \in End(A), \theta + 0 = \theta$$

4. Existence of Additive Inverses.

$$\forall \theta \in End(A), \exists \theta' \in End(A), \theta + \theta' = 0$$

5. Multiplication is associative. $\theta(\phi\psi) = (\theta\phi)\psi$
6. Multiplication is distributive over Addition.

- 6.1 Multiplication is left-distributive over Addition.

$$\theta(\phi + \psi) = \theta\phi + \theta\psi$$

- 6.2 Multiplication is right-distributive over Addition.

$$(\phi + \psi)\theta = \phi\theta + \psi\theta$$

$End(A)$: Noncommutative Ring

$\langle End(A), +, \circ \rangle$ is well-defined only if A is an abelian group.

Let $\theta, \phi, \psi \in End(A)$, $x \in A$

1. Addition is associative. $\theta + (\phi + \psi) = (\theta + \phi) + \psi$

2. Addition is commutative. $\theta + \phi = \phi + \theta$

3. Existence of Additive Identity.

$$\forall \theta \in End(A), \exists 0 \in End(A), \theta + 0 = \theta$$

4. Existence of Additive Inverses.

$$\forall \theta \in End(A), \exists \theta' \in End(A), \theta + \theta' = 0$$

5. Multiplication is associative. $\theta(\phi\psi) = (\theta\phi)\psi$

6. Multiplication is distributive over Addition.

6.1 Multiplication is left-distributive over Addition.

$$\theta(\phi + \psi) = \theta\phi + \theta\psi$$

6.2 Multiplication is right-distributive over Addition.

$$(\phi + \psi)\theta = \phi\theta + \psi\theta$$

7. Existence of Multiplicative Identity.

$$\forall \theta \in End(A), \exists I \in End(A), \theta I = \theta = I\theta$$

$End(A)$: Noncommutative Ring

$\langle End(A), +, \circ \rangle$ is well-defined only if A is an abelian group.

Let $\theta, \phi, \psi \in End(A)$, $x \in A$

1. Addition is associative. $\theta + (\phi + \psi) = (\theta + \phi) + \psi$

2. Addition is commutative. $\theta + \phi = \phi + \theta$

3. Existence of Additive Identity.

$$\forall \theta \in End(A), \exists 0 \in End(A), \theta + 0 = \theta$$

4. Existence of Additive Inverses.

$$\forall \theta \in End(A), \exists \theta' \in End(A), \theta + \theta' = 0$$

5. Multiplication is associative. $\theta(\phi\psi) = (\theta\phi)\psi$

6. Multiplication is distributive over Addition.

6.1 Multiplication is left-distributive over Addition.

$$\theta(\phi + \psi) = \theta\phi + \theta\psi$$

6.2 Multiplication is right-distributive over Addition.

$$(\phi + \psi)\theta = \phi\theta + \psi\theta$$

7. Existence of Multiplicative Identity.

$$\forall \theta \in End(A), \exists I \in End(A), \theta I = \theta = I\theta$$

8. Multiplication is non-commutative. $\theta\phi \neq \phi\theta$

Functions - Graduate Level Notions

well-defined function

Let $f : A \rightarrow B$ be a function. Then,

1. $\forall x \in A, f(x) \in B$
2. If $x_1 \xrightarrow{f} y_1$, and $x_2 \xrightarrow{f} y_2$, then $x_1 = x_2 \implies y_1 = y_2$

Example

► $f(x) = \sqrt{x}$

Functions - Graduate Level Notions

well-defined function

Let $f : A \rightarrow B$ be a function. Then,

1. $\forall x \in A, f(x) \in B$
2. If $x_1 \xrightarrow{f} y_1$, and $x_2 \xrightarrow{f} y_2$, then $x_1 = x_2 \implies y_1 = y_2$

Example

- ▶ $f(x) = \sqrt{x}$
- ▶ $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = \sqrt{x}$ is not well-defined. Since, $f(-1) \notin \mathbb{R}$

Functions - Graduate Level Notions

well-defined function

Let $f : A \rightarrow B$ be a function. Then,

1. $\forall x \in A, f(x) \in B$
2. If $x_1 \xrightarrow{f} y_1$, and $x_2 \xrightarrow{f} y_2$, then $x_1 = x_2 \implies y_1 = y_2$

Example

- ▶ $f(x) = \sqrt{x}$
- ▶ $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = \sqrt{x}$ is not well-defined. Since, $f(-1) \notin \mathbb{R}$
- ▶ $f : \mathbb{R}^+ \rightarrow \mathbb{R}, f(x) = \sqrt{x}$ is not well-defined. Since, $f(1) = ?$

Functions - Graduate Level Notions

well-defined function

Let $f : A \rightarrow B$ be a function. Then,

1. $\forall x \in A, f(x) \in B$
2. If $x_1 \xrightarrow{f} y_1$, and $x_2 \xrightarrow{f} y_2$, then $x_1 = x_2 \implies y_1 = y_2$

Example

- ▶ $f(x) = \sqrt{x}$ is not appreciated.
- ▶ $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+, f(x) = \sqrt{x}$ is preferred.
- ▶ $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = \sqrt{x}$ is not well-defined. Since, $f(-1) \notin \mathbb{R}$
- ▶ $f : \mathbb{R}^+ \rightarrow \mathbb{R}, f(x) = \sqrt{x}$ is not well-defined. Since, $f(1) = ?$
- ▶ $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+, f(x) = \sqrt{x}$ is well-defined.

Algebraic Structure is well-defined ?

well-defined ring $\langle R, +, * \rangle$

1. R is a set. — **usually true**

Algebraic Structure is well-defined ?

well-defined ring $\langle R, +, * \rangle$

1. R is a set. — **usually true**
2. The addition operation $+ : R \times R \rightarrow R$ is well-defined.

Algebraic Structure is well-defined ?

well-defined ring $\langle R, +, * \rangle$

1. R is a set. — **usually true**
2. The addition operation $+: R \times R \rightarrow R$ is well-defined.
 - 2.1 $\forall x, y \in R, +(x, y) = x + y \in R$ — **closure property**

Algebraic Structure is well-defined ?

well-defined ring $\langle R, +, * \rangle$

1. R is a set. — **usually true**
2. The addition operation $+ : R \times R \rightarrow R$ is well-defined.
 - 2.1 $\forall x, y \in R, +(x, y) = x + y \in R$ — **closure property**
 - 2.2 $(x_1, y_1) \xrightarrow{+} z_1, (x_2, y_2) \xrightarrow{+} z_2$. OR $x_1 + y_1 = z_1, x_2 + y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**

Algebraic Structure is well-defined ?

well-defined ring $\langle R, +, * \rangle$

1. R is a set. — **usually true**
2. The addition operation $+: R \times R \rightarrow R$ is well-defined.
 - 2.1 $\forall x, y \in R, +(x, y) = x + y \in R$ — **closure property**
 - 2.2 $(x_1, y_1) \xrightarrow{+} z_1, (x_2, y_2) \xrightarrow{+} z_2$. OR $x_1 + y_1 = z_1, x_2 + y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**

Algebraic Structure is well-defined ?

well-defined ring $\langle R, +, * \rangle$

1. R is a set. — **usually true**
2. The addition operation $+: R \times R \rightarrow R$ is well-defined.
 - 2.1 $\forall x, y \in R, +(x, y) = x + y \in R$ — **closure property**
 - 2.2 $(x_1, y_1) \xrightarrow{+} z_1, (x_2, y_2) \xrightarrow{+} z_2$. OR $x_1 + y_1 = z_1, x_2 + y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**
3. The multiplication operation $*: R \times R \rightarrow R$ is well-defined.

Algebraic Structure is well-defined ?

well-defined ring $\langle R, +, * \rangle$

1. R is a set. — **usually true**
2. The addition operation $+: R \times R \rightarrow R$ is well-defined.
 - 2.1 $\forall x, y \in R, +(x, y) = x + y \in R$ — **closure property**
 - 2.2 $(x_1, y_1) \xrightarrow{+} z_1, (x_2, y_2) \xrightarrow{+} z_2$. OR $x_1 + y_1 = z_1, x_2 + y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**
3. The multiplication operation $*: R \times R \rightarrow R$ is well-defined.
 - 3.1 $\forall x, y \in R, *(x, y) = x * y \in R$ — **closure property**

Algebraic Structure is well-defined ?

well-defined ring $\langle R, +, * \rangle$

1. R is a set. — **usually true**
2. The addition operation $+: R \times R \rightarrow R$ is well-defined.
 - 2.1 $\forall x, y \in R, +(x, y) = x + y \in R$ — **closure property**
 - 2.2 $(x_1, y_1) \xrightarrow{+} z_1, (x_2, y_2) \xrightarrow{+} z_2$. OR $x_1 + y_1 = z_1, x_2 + y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**
3. The multiplication operation $*: R \times R \rightarrow R$ is well-defined.
 - 3.1 $\forall x, y \in R, *(x, y) = x * y \in R$ — **closure property**
 - 3.2 $(x_1, y_1) \xrightarrow{*} z_1, (x_2, y_2) \xrightarrow{*} z_2$. OR $x_1 * y_1 = z_1, x_2 * y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**

Algebraic Structure is well-defined ?

well-defined ring $\langle R, +, * \rangle$

1. R is a set. — **usually true**
2. The addition operation $+: R \times R \rightarrow R$ is well-defined.
 - 2.1 $\forall x, y \in R, +(x, y) = x + y \in R$ — **closure property**
 - 2.2 $(x_1, y_1) \xrightarrow{+} z_1, (x_2, y_2) \xrightarrow{+} z_2$. OR $x_1 + y_1 = z_1, x_2 + y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**
3. The multiplication operation $*: R \times R \rightarrow R$ is well-defined.
 - 3.1 $\forall x, y \in R, *(x, y) = x * y \in R$ — **closure property**
 - 3.2 $(x_1, y_1) \xrightarrow{*} z_1, (x_2, y_2) \xrightarrow{*} z_2$. OR $x_1 * y_1 = z_1, x_2 * y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**

Algebraic Structure is well-defined ?

well-defined ring $\langle R, +, * \rangle$

1. R is a set. — **usually true**
 2. The addition operation $+: R \times R \rightarrow R$ is well-defined.
 - 2.1 $\forall x, y \in R, +(x, y) = x + y \in R$ — **closure property**
 - 2.2 $(x_1, y_1) \xrightarrow{+} z_1, (x_2, y_2) \xrightarrow{+} z_2$. OR $x_1 + y_1 = z_1, x_2 + y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**
 3. The multiplication operation $*: R \times R \rightarrow R$ is well-defined.
 - 3.1 $\forall x, y \in R, *(x, y) = x * y \in R$ — **closure property**
 - 3.2 $(x_1, y_1) \xrightarrow{*} z_1, (x_2, y_2) \xrightarrow{*} z_2$. OR $x_1 * y_1 = z_1, x_2 * y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**
- $\langle \mathbb{R}, +, * \rangle, x * y = \sqrt{xy}$ is not well-defined.

Algebraic Structure is well-defined ?

well-defined ring $\langle R, +, * \rangle$

1. R is a set. — **usually true**
 2. The addition operation $+: R \times R \rightarrow R$ is well-defined.
 - 2.1 $\forall x, y \in R, +(x, y) = x + y \in R$ — **closure property**
 - 2.2 $(x_1, y_1) \xrightarrow{+} z_1, (x_2, y_2) \xrightarrow{+} z_2$. OR $x_1 + y_1 = z_1, x_2 + y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**
 3. The multiplication operation $*: R \times R \rightarrow R$ is well-defined.
 - 3.1 $\forall x, y \in R, *(x, y) = x * y \in R$ — **closure property**
 - 3.2 $(x_1, y_1) \xrightarrow{*} z_1, (x_2, y_2) \xrightarrow{*} z_2$. OR $x_1 * y_1 = z_1, x_2 * y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**
- $\langle \mathbb{R}, +, \star \rangle, x \star y = \sqrt{xy}$ is not well-defined.
- Ring multiplication is not closed, $-1 \star 1 \notin \mathbb{R}$

Algebraic Structure is well-defined ?

well-defined ring $\langle R, +, * \rangle$

1. R is a set. — **usually true**
 2. The addition operation $+: R \times R \rightarrow R$ is well-defined.
 - 2.1 $\forall x, y \in R, +(x, y) = x + y \in R$ — **closure property**
 - 2.2 $(x_1, y_1) \xrightarrow{+} z_1, (x_2, y_2) \xrightarrow{+} z_2$. OR $x_1 + y_1 = z_1, x_2 + y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**
 3. The multiplication operation $*: R \times R \rightarrow R$ is well-defined.
 - 3.1 $\forall x, y \in R, *(x, y) = x * y \in R$ — **closure property**
 - 3.2 $(x_1, y_1) \xrightarrow{*} z_1, (x_2, y_2) \xrightarrow{*} z_2$. OR $x_1 * y_1 = z_1, x_2 * y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**
- $\langle \mathbb{R}, +, * \rangle, x * y = \sqrt{xy}$ is not well-defined.
- Ring multiplication is not closed, $-1 * 1 \notin \mathbb{R}$
 - Ring multiplication is not well-defined, $1 * 1 = ?$

Algebraic Structure is well-defined ?

well-defined ring $\langle R, +, * \rangle$

1. R is a set. — **usually true**
 2. The addition operation $+: R \times R \rightarrow R$ is well-defined.
 - 2.1 $\forall x, y \in R, +(x, y) = x + y \in R$ — **closure property**
 - 2.2 $(x_1, y_1) \xrightarrow{+} z_1, (x_2, y_2) \xrightarrow{+} z_2$. OR $x_1 + y_1 = z_1, x_2 + y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**
 3. The multiplication operation $*: R \times R \rightarrow R$ is well-defined.
 - 3.1 $\forall x, y \in R, *(x, y) = x * y \in R$ — **closure property**
 - 3.2 $(x_1, y_1) \xrightarrow{*} z_1, (x_2, y_2) \xrightarrow{*} z_2$. OR $x_1 * y_1 = z_1, x_2 * y_2 = z_2$
If $x_1 = x_2, y_1 = y_2$, then $z_1 = z_2$ — **well-defined**
- $\langle \mathbb{R}, +, \star \rangle, x \star y = \sqrt{xy}$ is not well-defined.
- Ring multiplication is not closed, $-1 \star 1 \notin \mathbb{R}$
 - Ring multiplication is not well-defined, $1 \star 1 = ?$
- $\langle \mathbb{R}^+, +, \star \rangle, x \star y = \sqrt{xy}$ is well-defined.

$\langle \text{End}(A), +, \circ \rangle$ is well-defined ?

Given $\langle A, * \rangle$ is an abelian group with identity e .

1. $\text{End}(A)$ is a non-empty set, since $0 \in \text{End}(A)$
where $0 : A \rightarrow A$ is defined by the relation $0(x) = e$.
2. Addition is well-defined.
 - 2.1 $\phi + \psi$ is an endomorphism of A
 - 2.2 $(\phi + \psi)(x)$ is uniquely defined for every $x \in A$
3. Multiplication/composition is well-defined.
 - 3.1 $\phi\psi$ is an endomorphism of A
 - 3.2 $(\phi\psi)(x)$ is uniquely defined for every $x \in A$.

Note : An endomorphism of A is a homomorphism from A into A .

Step 1 : $\phi + \psi \in \text{End}(A)$

Let $\phi, \psi \in \text{End}(A)$, and $x \in A$.

1. function $\phi + \psi : A \rightarrow A$ is well-defined,
 $(\phi + \psi)(x) = \phi(x) + \psi(x) \in A$

1.1 ϕ, ψ are well-defined. $\phi(x), \psi(x) \in A$

1.2 Group addition is closed.

2. $\phi + \psi : A \rightarrow A$ is a homomorphism

$$(\phi + \psi)(x + y) = (\phi + \psi)(x) + (\phi + \psi)(y)$$

$$(\phi + \psi)(x + y) = \phi(x + y) + \psi(x + y) \quad (1)$$

$$= (\phi(x) + \phi(y)) + (\psi(x) + \psi(y)) \quad (2)$$

$$= \phi(x) + ((\phi(y) + \psi(x)) + \psi(y)) \quad (3)$$

$$= \phi(x) + ((\psi(x) + \phi(y)) + \psi(y)) \quad (4)$$

$$= (\phi(x) + \psi(x)) + (\phi(y) + \psi(y)) \quad (5)$$

$$= (\phi + \psi)(x) + (\phi + \psi)(y) \quad (6)$$

Step 2 : $(\phi + \psi)(x)$ is uniquely defined

Let $\phi, \psi \in \text{End}(A)$, and $x \in A$.

1. $\phi(x), \psi(x)$ are uniquely defined in A .
 $\forall x \in A, \forall \phi \in \text{End}(A), \exists \text{ unique } \phi(x) \in A$.
2. Group addition is uniquely defined in A .
 $\phi(x), \psi(x) \in A \implies \exists \text{ unique } \phi(x) + \psi(x) \in A$

Step 3 : $\phi\psi \in \text{End}(A)$

Let $\phi, \psi \in \text{End}(A)$, and $x \in A$.

1. function $\phi\psi : A \rightarrow A$ is well-defined, $\phi\psi(x) = \phi(\psi(x))$

1.1 $\phi(\psi(x)) = \phi(y) \in A$ where $y = \psi(x) \in A$

1.2 $\psi(x), \phi(y) \in A$ where $y = \psi(x) \in A$.

2. $\phi\psi = \phi \circ \psi$ is a homomorphism.

$$(\phi\psi)(x + y) = \phi(\psi(x + y)) \quad (7)$$

$$= \phi(\psi(x) + \psi(y)) \quad (8)$$

$$= \phi(\psi(x)) + \phi(\psi(y)) \quad (9)$$

$$= \phi\psi(x) + \phi\psi(y) \quad (10)$$

Step 4 : $(\phi\psi)(x)$ is uniquely defined

Let $\phi, \psi \in \text{End}(A)$, and $x, y \in A$.

1. $\psi(x) = y$ is uniquely defined in A .
2. $\psi(x), \phi(y)$ are uniquely defined in A .

$\forall x \in A, \exists \text{ unique } (\phi\psi)(x) = \phi(\psi(x)) \in A$