

MATH 8660 Sec.1: Homework No.11

Due: Tuesday, November 19 at 8:05am.

No.1. (8 pts.) Let H_1 , H_2 , $B(\cdot, \cdot)$ and F be as described in the Brezzi-Nečas-Babuška Theorem. Additionally, let $H_{1,h}$ and $H_{2,h}$ be closed subspaces of H_1 and H_2 , respectively.

(i) Does it immediately follow from the Brezzi-Nečas-Babuška Theorem that there exists a unique $u_h \in H_{1,h}$ satisfying $B(u_h, v_h) = F(v_h)$, $\forall v_h \in H_{2,h}$?

If not, what additional assumptions are needed??

(ii) Follow the proof of Céa's Lemma to establish an error bound for $\|u - u_h\|_{H_1}$.

No.2. (8 pts.) Consider the problem of determining u satisfying

$$\mathcal{L}(u)(x) := -\nabla \cdot k(\mathbf{x}) \nabla u(\mathbf{x}) + \mathbf{b}(\mathbf{x}) \cdot \nabla u(\mathbf{x}) + e(\mathbf{x})u(\mathbf{x}) = f(\mathbf{x}), \quad \mathbf{x} \in \Omega, \quad (0.1)$$

$$\text{subject to } \begin{cases} u(\mathbf{x}) = 0, & \mathbf{x} \in \Gamma \subset \partial\Omega, \\ -k(x) \frac{\partial u(\mathbf{x})}{\partial \mathbf{n}} = g_2(\mathbf{x}), & \mathbf{x} \in \partial\Omega \setminus \Gamma \end{cases} \quad (0.2)$$

Give the weak formulation for (0.1), (0.2).

No.3. (8 pts.) Consider the boundary value problem: *Given f and g determine u satisfying*

$$-\nabla \cdot (k(\mathbf{x}) \nabla u(\mathbf{x})) + \mathbf{b}(\mathbf{x}) \cdot \nabla u(\mathbf{x}) + c(\mathbf{x})u(\mathbf{x}) = f \quad \text{in } \Omega, \quad (0.3)$$

$$\text{subject to } \frac{\partial u(\mathbf{x})}{\partial \mathbf{n}} = g(\mathbf{x}), \quad \mathbf{x} \in \partial\Omega. \quad (0.4)$$

(i) Derive the weak formulation for the solution of (0.3), (0.4), $B(u, v) = F(v)$.

(ii) Identify $\mathcal{L} : X \rightarrow Y$ such that $\mathcal{L}(w)(\cdot) = B(w, \cdot)$.

(iii) Determine the adjoint operator, \mathcal{L}^* , of \mathcal{L} , and give an associated adjoint problem to (0.3), (0.4).
