

Perturbation of Semigroups

Jacob Manning and Cameron Spiess

Clemson University

December 5, 2024

3.1 Perturbations by Bounded Linear Operators

Section 3.1 Perturbations by Bounded Linear Operators

3.1 Perturbations by Bounded Linear Operators

Definition

- If A is the infinitesimal generator of $T(t)$ (a semigroup of linear operators) then we say A generates $\{T(t) : t \geq 0\}$.
- C_0 semigroup $T(t)$ is the semigroup of strongly continuous operators that are bounded on X .

Theorem (3.1.1)

Let X be a Banach space and let A be the infinitesimal generator of a C_0 semigroup $T(t)$ on X , satisfying $\|T(t)\| \leq Me^{\omega t}$. If B is a bounded linear operator on X then $A + B$ is the infinitesimal generator of a C_0 semigroup $S(t)$ on X , satisfying $\|S(t)\| \leq Me^{(\omega + M\|B\|)t}$.

3.1 Perturbations by Bounder Linear Operators

Remark

- An equivalent definition of infinitesimal generator of $\{T(t) : t \geq 0\}$ is, let $A : D(A) \subset X \rightarrow X$ is defined by $D(A) = \{x \in X : \lim_{t \downarrow 0} \frac{1}{t}(T(t)x - x) \text{ exists}\}$ and $Ax = \lim_{t \downarrow 0} \frac{1}{t}(T(t)x - x)$
- The initial bound of $\|T(t)\| \leq Me^{\omega t}$ is given to the true generator A . However, there will exist some error when computing this generating numerically. The numerically computed generator could be $\tilde{A} = A + \Delta A$ where ΔA is a bounder linear operator.

3.1 Perturbations by Bounded Linear Operators

We are interested in relations between the semigroup $T(t)$ generated by A and the semigroup $S(t)$ generated by $A + B$. Consider the following operator,

$$H(s) = T(t - s)S(s)$$

for $x \in D(A) = D(A + B)$ such that $s \rightarrow H(s)x$. Notice that $H(s)$ is differentiable.

$$H'(s)x = T(t - s)BS(s)x \text{ for } x \in D(A)$$

Integrating this from 0 to t , we can find that

$$S(t)x = T(t)x + \int_0^t T(t - s)BS(s)x ds$$

Both sides are bounded, meaning $S(t)$ is a solution.

3.1 Perturbations by Bounded Linear Operators

Theorem (3.1.2)

Let $T(t)$ be a C_0 semigroup satisfying $\|T(t)\| \leq Me^{\omega t}$. Let B be a bounded operator on X . Then there exists a unique family $V(t)$, $t \geq 0$ of bounded operators on X such that $t \rightarrow V(t)x$ is continuous on $[0, \infty)$ for every $x \in X$ and

$$V(t)x = T(t)x + \int_0^t T(t-s)BV(s)x ds \quad x \in X.$$

Remark

We can immediately gain an explicit representation of $S(t)$ in terms of $T(t)$ such that

$$S(t) = \sum_{n \in \mathbb{N}} S_{n-1}(t)$$

where $S_0(t) = T(t)$ and $S_n(t) = \int_0^t T(t-s)BS_{n-1}(s)x ds$ with $x \in X$. The series converges uniformly.

3.1 Perturbations by Bounded Linear Operators

Corollary (3.1.3)

Let A be the infinitesimal generator of a C_0 semigroup $T(t)$ satisfying $\|T(t)\| \leq Me^{\omega t}$. Let B be a bounded operator and let $S(t)$ be the C_0 semigroup generated by $A + B$. Then

$$\|S(t) - T(t)\| \leq Me^{\omega t}(e^{M\|B\|t} - 1).$$

Proof.

From our theorems 3.1.1 and 3.1.2 we gain the following

$$\begin{aligned} \|S(t)x - T(t)x\| &\leq \int_0^t \|T(t-s)\| \|B\| \|S(s)\| \|x\| ds \\ &\leq M^2 e^{\omega t} \|B\| \int_0^t e^{M\|B\|s} \|x\| ds \\ &= Me^{\omega t} (e^{M\|B\|t} - 1) \|x\| \end{aligned}$$

3.1 Perturbations by Bounded Linear Operators

Remark

The big takeaway of this section is theorem 3.1.1, which shows that we can add a bounded linear operator to an infinitesimal operator and it does not destroy the property.

Follow up question

What other properties can a semigroup $T(t)$ generated by an infinitesimal operator A carry over?

It turns out our perturbation $(A + B)$ will be compact or analytic if our infinitesimal operator A is compact or analytic.

3.1 Perturbations by Bounded Linear Operators

Proposition (3.1.4)

Let A be the infinitesimal generator of a compact C_0 semigroup $T(t)$. Let B be a bounded operator, then $A + B$ is the infinitesimal generator of a compact C_0 semigroup $S(t)$.

Proposition (3.2.1)

Let A be the infinitesimal generator of an analytic semigroup. Let B be a closed linear operator satisfying $D(B) \supset D(A)$ and

$$\|Bx\| \leq a\|Ax\| + b\|x\|$$

for $x \in D(A)$ There exists a positive number δ such that if $0 \leq a \leq \delta$ then $A + B$ is the infinitesimal generator of an analytic semigroup

3.1 Perturbations by Bounded Linear Operators

Remark

Not all of the properties of the semigroup of $T(t)$ are preserved by a bounded perturbation of its infinitesimal generator.

Ex

Let A be the infinitesimal generator of a semigroup $T(t)$ which is continuous in the uniform operator topology for $t \geq t_0 \geq 0$, or is differentiable for $t \geq t_0 \geq 0$, or is compact for $t \geq t_0 \geq 0$. Then $S(t)$, the semigroup generated by $A + B$ where B is a bounded operator need not have the corresponding property.

3.3 Perturbations of Infinitesimal Generators of Contraction Semigroups

3.3 Perturbations of Infinitesimal Generators of Contraction Semigroups

3.3 Perturbations of Infinitesimal Generators of Contraction Semigroups

Recall

Theorem (1.4.3 Lumer-Phillips)

- If A is dissipative and there is a $\lambda_0 > 0$ such that $R(\lambda_0 I - A) = X$, then A is the infinitesimal generator of a C_0 semigroup of contractions on X .
- If A is the infinitesimal generator of a C_0 semigroup of contractions on X then $R(\lambda I - A) = X$ for all $\lambda > 0$ and A is dissipative.

3.3 Perturbations of Infinitesimal Generators of Contraction Semigroups

Definition (3.3.1)

- A linear operator A is called dissipative if $\operatorname{Re}\langle Ax, x \rangle \leq 0 \quad \forall x \in D(A)$
- A dissipative operator A such that $R(I - A) = X$ is called a maximal dissipative operator.
- An operator A is called closable if there exists a closed operator \bar{A} such that $G(A) \subset G(\bar{A})$.

Remark

If A is dissipative, for any $c > 0$, cA is also dissipative.

Theorem (Lumer-Phillips)

A densely defined operator A is the infinitesimal generator of a C_0 semigroup of contractions iff A is maximal dissipative.

3.3 Perturbations of Infinitesimal Generators of Contraction Semigroups

Theorem (3.3.2)

Let A and B be linear operators in X such that $D(B) \supset D(A)$ and $A + tB$ is dissipative for $0 \leq t \leq 1$. If

$$\|Bx\| \leq \alpha \|Ax\| + \beta \|x\| \quad x \in D(A)$$

where $0 \leq \alpha < 1$, $\beta \geq 0$ and for some $0 \leq t_0 \leq 1$, $A + t_0B$ is maximal dissipative then $A + tB$ is maximal dissipative for all $0 \leq t \leq 1$.

This theorem is often used with the following corollary.

3.3 Perturbations of Infinitesimal Generators of Contraction Semigroups

Corollary (3.3.3)

Let A be the infinitesimal generator of a C_0 semigroup of contractions. Let B be dissipative and satisfy $D(B) \supset D(A)$ and

$$\|Bx\| \leq \alpha \|Ax\| + \beta \|x\| \quad x \in D(A)$$

where $0 \leq \alpha < 1$, $\beta \geq 0$. Then $A + B$ is the infinitesimal generator of a C_0 semigroup of contractions.

Remark

If $A + tB$ is dissipative for $0 \leq t \leq 1$, $D(B) \supset D(A)$, $\overline{D(A)} = X$, and

$$\|Bx\| \leq \alpha \|Ax\| + \beta \|x\| \quad x \in D(A)$$

where $0 \leq \alpha < 1$, $\beta \geq 0$. Then either both A and $A + B$ are maximal dissipative or neither.

3.3 Perturbations of Infinitesimal Generators of Contraction Semigroups

$\alpha = 1$ can be an issue since $A + B$ may not be closed and thus cannot be the infinitesimal generator of a C_0 semigroup.

3.3 Perturbations of Infinitesimal Generators of Contraction Semigroups

$\alpha = 1$ can be an issue since $A + B$ may not be closed and thus cannot be the infinitesimal generator of a C_0 semigroup.

Ex

Let iA be a self adjoint operator in a Hilbert space. It follows that A and $-A$ are infinitesimal generators of C_0 semigroups of contractions. Taking $\alpha = 1$ and $\beta = 0$. It follows

$$\| -Ax \| \leq 1 * \| Ax \| + 0 * \| x \| = \| Ax \|^2$$

However $\overline{A + (-A)} \Big|_{D(A)}$ is not closed.

3.3 Perturbations of Infinitesimal Generators of Contraction Semigroups

What conditions can we set $\alpha = 1$?

3.3 Perturbations of Infinitesimal Generators of Contraction Semigroups

What conditions can we set $\alpha = 1$?

Theorem (3.3.4)

Let A be the infinitesimal generator of a C_0 semigroup of contractions. Let B be dissipative and satisfy $D(B) \supset D(A)$ and

$$\|Bx\| \leq \|Ax\| + \beta \|x\| \quad x \in D(A)$$

where $\beta \geq 0$. If B^ is densely defined, then the closure $\overline{A+B}$ of $A+B$ is the infinitesimal generator of a C_0 semigroup of contractions.*

3.3 Perturbations of Infinitesimal Generators of Contraction Semigroups

If X is a reflexive Banach space and T is closable and densely defined, we know that T^* is closed and $D(T^*)$ is dense in X^* . It follows

Corollary (3.3.5)

Let X be a reflexive Banach space and let A be the infinitesimal generator of a C_0 semigroup of contractions in X . Let B be dissipative such that $D(B) \supset D(A)$ and

$$\|Bx\| \leq \|Ax\| + \beta \|x\| \quad x \in D(A)$$

where $\beta \geq 0$. Then $\overline{A+B}$, the closure of $A+B$, is the infinitesimal generator of a C_0 semigroup of contractions in X .

Amnon Pazy. "Perturbations and Approximations." Semigroups of Linear Operators and Applications to Partial Differential Equations. Springer Science & Business Media, 6 Dec. 2012.