

C_0 Semigroups

Jacob Manning

Clemson University

April 16, 2025

This presentation is based on three books, namely "Topics in functional analysis and applications" by Srinivasan Kesavan[1], "Introductory functional analysis with applications" by Erwin Kreyszig[2], and "Semigroups of linear operators and applications to partial differential equations" by Amnon Pazy[3].

Definition

- A linear operator $A : D(A) \subseteq X \rightarrow Y$ is said to be **bounded** if there exists a $C > 0$ such that

$$\|Au\|_Y \leq C\|u\|_X, \quad \text{for every } u \in D(A)$$

Otherwise it is said to be **unbounded**.

Definition

- A linear operator $A : D(A) \subseteq X \rightarrow Y$ is said to be **bounded** if there exists a $C > 0$ such that

$$\|Au\|_Y \leq C\|u\|_X, \quad \text{for every } u \in D(A)$$

Otherwise it is said to be **unbounded**.

- A linear operator $A : D(A) \subseteq X \rightarrow Y$ is said to be **densely defined** if $\overline{D(A)} = X$

Definition

- A linear operator $A : D(A) \subseteq X \rightarrow Y$ is said to be **bounded** if there exists a $C > 0$ such that

$$\|Au\|_Y \leq C\|u\|_X, \quad \text{for every } u \in D(A)$$

Otherwise it is said to be **unbounded**.

- A linear operator $A : D(A) \subseteq X \rightarrow Y$ is said to be **densely defined** if $\overline{D(A)} = X$
- A linear operator $A : D(A) \subseteq X \rightarrow Y$ is said to be **closed** if the **graph**

$$G(A) = \{(u, Au) \mid u \in D(A)\} \subseteq X \times Y$$

is closed as a subspace of $X \times Y$

Definition

Let X be a Banach space with dual space X' . Denote $x' \in X'$ at $x \in X$ by $\langle x', x \rangle$ or $\langle x, x' \rangle$. Define the following set $F(x) \subseteq X'$ as

$$F(x) = \{x' \mid \langle x', x \rangle = \|x\|^2 = \|x'\|^2\}$$

(This set is non-empty by the Hahn-Banach theorem.)

Definition

Let X be a Banach space with dual space X' . Denote $x' \in X'$ at $x \in X$ by $\langle x', x \rangle$ or $\langle x, x' \rangle$. Define the following set $F(x) \subseteq X'$ as

$$F(x) = \{x' \mid \langle x', x \rangle = \|x\|^2 = \|x'\|^2\}$$

(This set is non-empty by the Hahn-Banach theorem.)

Definition (Dissipativity)

A linear operator A is dissipative if for every $x \in D(A)$ there is a $x' \in F(x)$ such that $\operatorname{Re} \langle Ax, x' \rangle \leq 0$

Definition

Let X be a Banach space with dual space X' . Denote $x' \in X'$ at $x \in X$ by $\langle x', x \rangle$ or $\langle x, x' \rangle$. Define the following set $F(x) \subseteq X'$ as

$$F(x) = \{x' \mid \langle x', x \rangle = \|x\|^2 = \|x'\|^2\}$$

(This set is non-empty by the Hahn-Banach theorem.)

Definition (Dissipativity)

A linear operator A is dissipative if for every $x \in D(A)$ there is a $x' \in F(x)$ such that $\operatorname{Re} \langle Ax, x' \rangle \leq 0$

Definition (Maximal Dissipativity)

A linear operator A is called maximally dissipative if it is dissipative and $R(I - A) = X$.

Definition

Let $X \neq \{0\}$ be a complex normed space and $A : D(A) \subseteq X \rightarrow X$ be a linear operator. With A we associate the operator

$$A_\lambda = \lambda I - A$$

where λ is a complex number and I is the identity operator on $D(A)$. If A_λ has an inverse, we denote it by $R_\lambda(A)$ and call it the resolvent operator of A or, simply, the **resolvent** of A . If it is clear which operator we are discussing, we will write R_λ .

Definition

Let $X \neq \{0\}$ be a complex normed space and $A : D(A) \subseteq X \rightarrow X$ be a linear operator. A regular value λ of A is a complex number such that

- R_λ exists,
- R_λ is bounded,
- R_λ is densely defined.

The resolvent set $\rho(A)$ of A is the set of all regular values λ of A . Its complement $\sigma(A) = \mathbb{C} \setminus \rho(A)$ in the complex plane \mathbb{C} is called the spectrum of A , and a $\lambda \in \sigma(A)$ is called a spectral value of A .

Definition

Recall the limit definition of the derivative

$$u_t = \lim_{h \downarrow 0} \frac{u(t+h) - u(t)}{h}$$

Definition

Recall the limit definition of the derivative

$$u_t = \lim_{h \downarrow 0} \frac{u(t+h) - u(t)}{h}$$

Example

Consider the following linear first order ODE

$$\begin{cases} u_t = -\alpha u & t > 0, \alpha \in \mathbb{R} \\ u(0) = u_0 \end{cases}$$

Lemma

$$\log(1+x) = x \quad \text{as} \quad x \downarrow 0$$

Motivation

Lemma

$$\log(1+x) = x \quad \text{as } x \downarrow 0$$

Proof

Let $y = \log(1+x)$.

Note $y \rightarrow 0$ as $x \rightarrow 0$

Thus,

$$\begin{aligned} 1+x &= e^y \\ &= 1+y+o(y) \\ &\rightarrow 1+y \end{aligned}$$

Now let's get back to the example

Example (Continued)

$$-\alpha = \frac{u_t}{u}$$

Example (Continued)

$$\begin{aligned} -\alpha &= \frac{u_t}{u} \\ &= \frac{1}{u(t)} \lim_{h \downarrow 0} \frac{u(t+h) - u(t)}{h} \end{aligned}$$

Example (Continued)

$$\begin{aligned} -\alpha &= \frac{u_t}{u} \\ &= \frac{1}{u(t)} \lim_{h \downarrow 0} \frac{u(t+h) - u(t)}{h} \\ &= \lim_{h \downarrow 0} \frac{1}{h} \log \left| 1 + \frac{u(t+h) - u(t)}{u(t)} \right| \end{aligned}$$

Example (Continued)

$$\begin{aligned} -\alpha &= \frac{u_t}{u} \\ &= \frac{1}{u(t)} \lim_{h \downarrow 0} \frac{u(t+h) - u(t)}{h} \\ &= \lim_{h \downarrow 0} \frac{1}{h} \log \left| 1 + \frac{u(t+h) - u(t)}{u(t)} \right| \\ &= \lim_{h \downarrow 0} \frac{1}{h} \log \left| \frac{u(t+h)}{u(t)} \right| \end{aligned}$$

Example (Continued)

$$\begin{aligned} -\alpha &= \frac{u_t}{u} \\ &= \frac{1}{u(t)} \lim_{h \downarrow 0} \frac{u(t+h) - u(t)}{h} \\ &= \lim_{h \downarrow 0} \frac{1}{h} \log \left| 1 + \frac{u(t+h) - u(t)}{u(t)} \right| \\ &= \lim_{h \downarrow 0} \frac{1}{h} \log \left| \frac{u(t+h)}{u(t)} \right| \\ &= \lim_{h \downarrow 0} \frac{\log |u(t+h)| - \log |u(t)|}{h} \end{aligned}$$

Example (Continued)

$$\begin{aligned} -\alpha &= \frac{u_t}{u} \\ &= \frac{1}{u(t)} \lim_{h \downarrow 0} \frac{u(t+h) - u(t)}{h} \\ &= \lim_{h \downarrow 0} \frac{1}{h} \log \left| 1 + \frac{u(t+h) - u(t)}{u(t)} \right| \\ &= \lim_{h \downarrow 0} \frac{1}{h} \log \left| \frac{u(t+h)}{u(t)} \right| \\ &= \lim_{h \downarrow 0} \frac{\log |u(t+h)| - \log |u(t)|}{h} \\ &= \frac{d}{dt} \log |u(t)| \end{aligned}$$

Example (Continued)

$$\begin{aligned} -\alpha &= \frac{u_t}{u} \\ &= \frac{1}{u(t)} \lim_{h \downarrow 0} \frac{u(t+h) - u(t)}{h} \\ &= \lim_{h \downarrow 0} \frac{1}{h} \log \left| 1 + \frac{u(t+h) - u(t)}{u(t)} \right| \\ &= \lim_{h \downarrow 0} \frac{1}{h} \log \left| \frac{u(t+h)}{u(t)} \right| \\ &= \lim_{h \downarrow 0} \frac{\log |u(t+h)| - \log |u(t)|}{h} \\ &= \frac{d}{dt} \log |u(t)| \end{aligned}$$

Therefore $u(t) = u_0 e^{-\alpha t}$

How can we generalize this idea?

Definition (C_0 Semigroups)

Let X be a Banach space and $\{S(t)\}_{t \geq 0}$ be a family of bounded linear operators on X . $\{S(t)\}_{t \geq 0}$ is said to be a C_0 semigroup if the following are true:

How can we generalize this idea?

Definition (C_0 Semigroups)

Let X be a Banach space and $\{S(t)\}_{t \geq 0}$ be a family of bounded linear operators on X . $\{S(t)\}_{t \geq 0}$ is said to be a C_0 semigroup if the following are true:

- $S(0) = I$, the identity of X

How can we generalize this idea?

Definition (C_0 Semigroups)

Let X be a Banach space and $\{S(t)\}_{t \geq 0}$ be a family of bounded linear operators on X . $\{S(t)\}_{t \geq 0}$ is said to be a C_0 semigroup if the following are true:

- $S(0) = I$, the identity of X
- $S(t + s) = S(t)S(s)$, for all $t, s \geq 0$

How can we generalize this idea?

Definition (C_0 Semigroups)

Let X be a Banach space and $\{S(t)\}_{t \geq 0}$ be a family of bounded linear operators on X . $\{S(t)\}_{t \geq 0}$ is said to be a C_0 semigroup if the following are true:

- $S(0) = I$, the identity of X
- $S(t + s) = S(t)S(s)$, for all $t, s \geq 0$
- For every $u \in X$

$$S(t)u \rightarrow u \quad \text{as } t \downarrow 0$$

Definition

Let $\{S(t)\}_{t \geq 0}$ be a C_0 semigroup on X . The **infinitesimal generator** of the semigroup is a linear operator A given by

$$D(A) = \left\{ u \in X \mid \lim_{t \downarrow 0} \frac{S(t)u - u}{t} \text{ exists} \right\}$$
$$Au = \lim_{t \downarrow 0} \frac{S(t)u - u}{t}, \quad u \in D(A)$$

Some C_0 Semigroup Properties

What are some properties of these semigroups?

Some C_0 Semigroup Properties

What are some properties of these semigroups?

Theorem

Let $\{S(t)\}_{t \geq 0}$ be a C_0 -semigroup on X . Then there exists $M \geq 1$ and ω such that

$$\|S(t)\| \leq Me^{\omega t}, \quad \text{for all } t \geq 0$$

Some C_0 Semigroup Properties

What are some properties of these semigroups?

Theorem

Let $\{S(t)\}_{t \geq 0}$ be a C_0 -semigroup on X . Then there exists $M \geq 1$ and ω such that

$$\|S(t)\| \leq Me^{\omega t}, \quad \text{for all } t \geq 0$$

This is a direct result of the Uniform Boundedness Principle. Thus we have the following definition.

Some C_0 Semigroup Properties

What are some properties of these semigroups?

Theorem

Let $\{S(t)\}_{t \geq 0}$ be a C_0 -semigroup on X . Then there exists $M \geq 1$ and ω such that

$$\|S(t)\| \leq Me^{\omega t}, \quad \text{for all } t \geq 0$$

This is a direct result of the Uniform Boundedness Principle. Thus we have the following definition.

Definition

*If $M = 1$ and $\omega = 0$, so that $\|S(t)\| \leq 1$ for all $t \geq 0$, we say that $\{S(t)\}$ is a **contraction semigroup**.*

How does this help us?

How does this help us?

Theorem

Let $\{S(t)\}_{t \geq 0}$ be a C_0 semigroup and let A be its infinitesimal generator. Let $u \in D(A)$. Then

$$S(t)u \in C^1([0, \infty); X) \cap C([0, \infty); X)$$

and

$$\frac{d}{dt}(S(t)u) = AS(t)u = S(t)Au$$

Proof

Let $u \in D(A)$. Then

Proof

Let $u \in D(A)$. Then

$$AS(t)u = \lim_{h \downarrow 0} \left(\frac{S(h) - I}{h} \right) S(t)u = \lim_{h \downarrow 0} S(t) \left(\frac{S(h) - I}{h} \right) = S(t)Au$$

Proof

Let $u \in D(A)$. Then

$$\begin{aligned} AS(t)u &= \lim_{h \downarrow 0} \left(\frac{S(h) - I}{h} \right) S(t)u = \lim_{h \downarrow 0} S(t) \left(\frac{S(h) - I}{h} \right) = S(t)Au \\ AS(t)u &= S(t)Au = D^+ S(t)u. \end{aligned}$$

Proof

Let $u \in D(A)$. Then

$$\begin{aligned} AS(t)u &= \lim_{h \downarrow 0} \left(\frac{S(h) - I}{h} \right) S(t)u = \lim_{h \downarrow 0} S(t) \left(\frac{S(h) - I}{h} \right) = S(t)Au \\ AS(t)u &= S(t)Au = D^+ S(t)u. \end{aligned}$$

Next consider

$$\frac{S(t)u - S(t-h)u}{h} = S(t-h) \frac{S(h)u - u}{h}$$

Proof

Let $u \in D(A)$. Then

$$\begin{aligned} AS(t)u &= \lim_{h \downarrow 0} \left(\frac{S(h) - I}{h} \right) S(t)u = \lim_{h \downarrow 0} S(t) \left(\frac{S(h) - I}{h} \right) = S(t)Au \\ AS(t)u &= S(t)Au = D^+ S(t)u. \end{aligned}$$

Next consider

$$\frac{S(t)u - S(t-h)u}{h} = S(t-h) \frac{S(h)u - u}{h}$$

$$\begin{aligned} \frac{S(t)u - S(t-h)u}{h} - S(t)Au &= S(t-h) \left(\frac{S(h)u - u}{h} - Au \right) \\ &\quad + (S(t-h) - S(t))Au \end{aligned}$$

Proof (Proof Continued)

$$\left\| S(t-h) \left(\frac{S(h)u - u}{h} - Au \right) \right\| \leq Me^{\omega t} \left\| \frac{S(h)u - u}{h} - Au \right\|$$

Proof (Proof Continued)

$$\left\| S(t-h) \left(\frac{S(h)u - u}{h} - Au \right) \right\| \leq Me^{\omega t} \left\| \frac{S(h)u - u}{h} - Au \right\|$$

$\rightarrow 0 \quad \text{as } h \downarrow 0$

Proof (Proof Continued)

$$\left\| S(t-h) \left(\frac{S(h)u - u}{h} - Au \right) \right\| \leq Me^{\omega t} \left\| \frac{S(h)u - u}{h} - Au \right\|$$
$$\rightarrow 0 \quad \text{as } h \downarrow 0$$
$$\| (S(t-h) - S(t))Au \| \rightarrow 0 \quad \text{as } h \downarrow 0$$

Proof (Proof Continued)

$$\left\| S(t-h) \left(\frac{S(h)u - u}{h} - Au \right) \right\| \leq Me^{\omega t} \left\| \frac{S(h)u - u}{h} - Au \right\|$$
$$\rightarrow 0 \quad \text{as } h \downarrow 0$$
$$\|(S(t-h) - S(t))Au\| \rightarrow 0 \quad \text{as } h \downarrow 0$$

Thus,

$$D^- S(t)u = S(t)Au = D^+ S(t)u$$

Proof (Proof Continued)

$$\left\| S(t-h) \left(\frac{S(h)u - u}{h} - Au \right) \right\| \leq Me^{\omega t} \left\| \frac{S(h)u - u}{h} - Au \right\|$$
$$\rightarrow 0 \quad \text{as } h \downarrow 0$$
$$\|(S(t-h) - S(t))Au\| \rightarrow 0 \quad \text{as } h \downarrow 0$$

Thus,

$$D^- S(t)u = S(t)Au = D^+ S(t)u$$

and

$$\frac{d}{dt}(S(t)u) = AS(t)u = S(t)Au$$

Proof (Proof Continued)

$$\left\| S(t-h) \left(\frac{S(h)u - u}{h} - Au \right) \right\| \leq M e^{\omega t} \left\| \frac{S(h)u - u}{h} - Au \right\|$$
$$\rightarrow 0 \quad \text{as } h \downarrow 0$$
$$\|(S(t-h) - S(t))Au\| \rightarrow 0 \quad \text{as } h \downarrow 0$$

Thus,

$$D^- S(t)u = S(t)Au = D^+ S(t)u$$

and

$$\frac{d}{dt}(S(t)u) = AS(t)u = S(t)Au$$

Similarly, by the boundedness of $S(t)$, the map $t \mapsto S(t)Au$ is continuous so $S(t)u \in C^1([0, \infty); X)$.

What does that tell us?

What does that tell us?

Remark

If A is the infinitesimal generator of a C_0 semigroup $\{S(t)\}$ then we know by the above theorem that

$$u(t) = S(t)u_0$$

defines the unique solution of the initial value problem

$$\begin{cases} \frac{du(t)}{dt} = Au(t), & t \geq 0 \\ u(0) = u_0 \end{cases}$$

Hille-Yosida Theorem

When can we guarantee that an operator is an infinitesimal generator of a contraction semigroup?

Hille-Yosida Theorem

When can we guarantee that an operator is an infinitesimal generator of a contraction semigroup?

Theorem

A linear unbounded operator A on a Banach space X is the infinitesimal generator of a contraction semigroup if and only if

When can we guarantee that an operator is an infinitesimal generator of a contraction semigroup?

Theorem

A linear unbounded operator A on a Banach space X is the infinitesimal generator of a contraction semigroup if and only if

- *A is closed*

When can we guarantee that an operator is an infinitesimal generator of a contraction semigroup?

Theorem

A linear unbounded operator A on a Banach space X is the infinitesimal generator of a contraction semigroup if and only if

- *A is closed*
- *A is densely defined*

When can we guarantee that an operator is an infinitesimal generator of a contraction semigroup?

Theorem

A linear unbounded operator A on a Banach space X is the infinitesimal generator of a contraction semigroup if and only if

- *A is closed*
- *A is densely defined*
- *For every $\lambda > 0$, $R_\lambda(A)$ is a bounded linear operator and*

$$\|R_\lambda(A)\| \leq \frac{1}{\lambda}$$

When can we guarantee that an operator is an infinitesimal generator of a contraction semigroup?

Theorem

A linear unbounded operator A on a Banach space X is the infinitesimal generator of a contraction semigroup if and only if

- *A is closed*
- *A is densely defined*
- *For every $\lambda > 0$, $R_\lambda(A)$ is a bounded linear operator and*

$$\|R_\lambda(A)\| \leq \frac{1}{\lambda}$$

This is a more general result and will not be shown.

In Hilbert spaces it might be easier to use the following theorem.

In Hilbert spaces it might be easier to use the following theorem.

Theorem

- If A is dissipative and there is a $\lambda_0 > 0$ such that $R(A_{\lambda_0}) = X$, then A is the infinitesimal generator of a C_0 semigroup of contractions on X .
- If A is the infinitesimal generator of a C_0 semigroup of contractions on X then $R(A_\lambda) = X$ for all $\lambda > 0$ and A is dissipative.

Proof

Let $\lambda > 0$, the dissipativeness of A implies that

$$\|A_\lambda x\| = \|\lambda x - Ax\| \geq \lambda \|x\| \quad \text{for every } \lambda > 0 \text{ and } x \in D(A).$$

Since $R(A_{\lambda_0}) = X$, it follows when $\lambda = \lambda_0$ that R_{λ_0} is a bounded linear operator and thus closed. This implies A is closed. If $R(A_\lambda) = X$ for every $\lambda > 0$ then $\rho(A) \supseteq (0, \infty)$ and $\|R_\lambda\| \leq \lambda^{-1}$. It follows by the Hille-Yosida theorem that A is the infinitesimal generator of a C_0 semigroup of contractions on X .

Proof

Let $\lambda > 0$, the dissipativeness of A implies that

$$\|A_\lambda x\| = \|\lambda x - Ax\| \geq \lambda \|x\| \quad \text{for every } \lambda > 0 \text{ and } x \in D(A).$$

Since $R(A_{\lambda_0}) = X$, it follows when $\lambda = \lambda_0$ that R_{λ_0} is a bounded linear operator and thus closed. This implies A is closed. If $R(A_\lambda) = X$ for every $\lambda > 0$ then $\rho(A) \supseteq (0, \infty)$ and $\|R_\lambda\| \leq \lambda^{-1}$. It follows by the Hille-Yosida theorem that A is the infinitesimal generator of a C_0 semigroup of contractions on X . Consider the set

$$\Lambda = \{\lambda \mid 0 < \lambda < \infty, R(A_\lambda) = X\}.$$

Let $\lambda \in \Lambda$. By the previous inequality, $\lambda \in \rho(A)$. Since $\rho(A)$ is open, the intersection of $B_r(\lambda) \cap \mathbb{R} \subseteq \Lambda$ and thus Λ is open.

Proof (Proof Continued)

On the other hand, let $\{\lambda_n\} \subseteq \Lambda$ and $\lambda_n \rightarrow \lambda > 0$. For every $y \in X$ there exists an $x_n \in D(A)$ such that

$$A_\lambda x_n = \lambda_n x_n - Ax_n = y$$

Proof (Proof Continued)

On the other hand, let $\{\lambda_n\} \subseteq \Lambda$ and $\lambda_n \rightarrow \lambda > 0$. For every $y \in X$ there exists an $x_n \in D(A)$ such that

$$A_\lambda x_n = \lambda_n x_n - Ax_n = y$$

From the inequality it follows that $\|x_n\| \leq \lambda_n^{-1} \|y\| \leq C$ for some $C > 0$. Now,

$$\begin{aligned}\lambda_m \|x_n - x_m\| &\leq \|\lambda_m(x_n - x_m) - A(x_n - x_m)\| \\ &= |\lambda_n - \lambda_m| \|x_n\| \\ &\leq C |\lambda_n - \lambda_m| \rightarrow 0\end{aligned}$$

We see $\{x_n\}$ is Cauchy. Let $x_n \rightarrow x$. It follows $Ax_n \rightarrow \lambda x - y$. Since A is closed, $x \in D(A)$ and $A_\lambda x = y$. Thus $R(A_\lambda) = X$ and $\lambda \in \Lambda$ which implies that Λ is closed.

Proof (Proof Continued)

On the other hand, let $\{\lambda_n\} \subseteq \Lambda$ and $\lambda_n \rightarrow \lambda > 0$. For every $y \in X$ there exists an $x_n \in D(A)$ such that

$$A_\lambda x_n = \lambda_n x_n - Ax_n = y$$

From the inequality it follows that $\|x_n\| \leq \lambda_n^{-1} \|y\| \leq C$ for some $C > 0$. Now,

$$\begin{aligned}\lambda_m \|x_n - x_m\| &\leq \|\lambda_m(x_n - x_m) - A(x_n - x_m)\| \\ &= |\lambda_n - \lambda_m| \|x_n\| \\ &\leq C |\lambda_n - \lambda_m| \rightarrow 0\end{aligned}$$

We see $\{x_n\}$ is Cauchy. Let $x_n \rightarrow x$. It follows $Ax_n \rightarrow \lambda x - y$. Since A is closed, $x \in D(A)$ and $A_\lambda x = y$. Thus $R(A_\lambda) = X$ and $\lambda \in \Lambda$ which implies that Λ is closed. By assumption, $\Lambda \neq \emptyset$, therefore $\Lambda = (0, \infty)$

Proof (Proof Continued)

If A is the infinitesimal generator of a C_0 semigroup of contractions, $S(t)$, on X , then by the Hille-Yosida theorem $\rho(A) \supseteq (0, \infty)$ and therefore $R(A_\lambda) = X$ for all $\lambda > 0$.

Proof (Proof Continued)

If A is the infinitesimal generator of a C_0 semigroup of contractions, $S(t)$, on X , then by the Hille-Yosida theorem $\rho(A) \supseteq (0, \infty)$ and therefore $R(A_\lambda) = X$ for all $\lambda > 0$. Furthermore if $x \in D(A)$, $x' \in F(x)$ then

$$|\langle S(t)x, x' \rangle| \leq \|S(t)x\| \|x'\| \leq \|x\|^2$$

Proof (Proof Continued)

If A is the infinitesimal generator of a C_0 semigroup of contractions, $S(t)$, on X , then by the Hille-Yosida theorem $\rho(A) \supseteq (0, \infty)$ and therefore $R(A_\lambda) = X$ for all $\lambda > 0$. Furthermore if $x \in D(A)$, $x' \in F(x)$ then

$$|\langle S(t)x, x' \rangle| \leq \|S(t)x\| \|x'\| \leq \|x\|^2$$

and therefore,

$$\operatorname{Re} \langle S(t)x - x, x' \rangle = \operatorname{Re} \langle S(t)x, x' \rangle - \|x\|^2 \leq 0.$$

By dividing the previous line by $t > 0$ and letting $t \downarrow 0$ yields

$$\operatorname{Re} \langle Ax, x' \rangle \leq 0.$$

By using our definition of maximally dissipative operators we can rewrite the Lumer-Phillips theorem as follows

By using our definition of maximally dissipative operators we can rewrite the Lumer-Phillips theorem as follows

Theorem

A densely defined operator A is the infinitesimal generator of a C_0 semigroup of contractions if and only if it is maximal dissipative.

- [1] Srinivasan Kesavan. *Topics in functional analysis and applications*. Wiley Eastern Ltd., 1989.
- [2] Erwin Kreyszig. *Introductory functional analysis with applications*. John Wiley & Sons, 1991.
- [3] Amnon Pazy. *Semigroups of linear operators and applications to partial differential equations*. Vol. 44. Springer Science & Business Media, 2012.