

Representation Varieties of Stacks and Trace Maps

Jacob Erlikhman

Abstract

We define a derived stack $\mathcal{R}ep_n(X)$ which generalizes the assignment $A \leadsto \mathrm{Rep}_n(A)$ to an algebra of its derived GL_n -representation variety of [BKR13] from algebras A to perfect stacks X over characteristic 0 fields. In the case of a quasi-projective classical scheme X , we show that $\mathcal{R}ep_n(X)$ admits a subfunctor $\mathrm{Quot}_{\mathcal{O}_X^n}^{n,\mathrm{fr}}(X) \subset \mathcal{R}ep_n(X)$, which is in fact represented by a derived scheme almost of finite type. We further construct a Fourier-Mukai integral transform between the derived categories of quasi-coherent sheaves on X and $\mathrm{Quot}_{\mathcal{O}_X^n}^{n,\mathrm{fr}}(X)$ which induces a trace map at the level of Hochschild homology generalizing the trace morphism constructed in [BKR13] (for finitely presented commutative algebras). We show that the subfunctor $\mathrm{Quot}_{\mathcal{O}_X^n}^{n,\mathrm{fr}}(X)$ is a derived enhancement of the framed locus of the Quot scheme of points and that this derived scheme is a GL_n -torsor over the stack of coherent length n torsion sheaves. Hence, this stack is an analog of the derived character stack for quasi-projective schemes, and we show that for smooth, Calabi-Yau X , it inherits a shifted symplectic structure in the sense of [Pan+11] from the one constructed in [BD18] on the moduli stack of perfect complexes with proper support. This structure is shown to give a generalization of the classical symplectic structure on character varieties of surfaces of genus $g \leq 1$ constructed by Goldman [Gol84].

Contents

1	Introduction	2
1.1	Motivation	2
1.2	Overview	3
1.3	Contents	4
1.4	Notation and Conventions	7
1.5	Glossary of Categories	8
1.6	Acknowledgments	8
2	Domain Walls	8
2.1	Dualizability	8
2.2	Trace Maps I	10
3	Extension to Stacks	15
3.1	Definition	15
3.2	Examples	18

4	Trace Maps II	20
4.1	Coherent Representation Moduli and Derived Quot Schemes	21
4.2	The Universal Representation	26
4.3	The Universal Representation Defines a Domain Wall	29
5	The Shifted Symplectic Structure	31
A	Comparison with Existing Constructions	33
A.1	Quot Schemes	33
	References	35

1 Introduction

1.1 Motivation

Given a topological surface Σ , its (SL_2) -representation and character varieties have been of classical interest since the beginning of the 20th century (see the introduction to [Gol09]). Classically, one fixes a (connected, reductive) Lie group G , and studies either representations of the fundamental group

$$\mathrm{Hom}_{\mathrm{Gp}}(\pi_1(\Sigma), G)$$

or these up to conjugacy

$$\mathrm{Hom}_{\mathrm{Gp}}(\pi_1(\Sigma), G)/G,$$

where G acts by the conjugation action. In the case of $G = \mathrm{GL}_n$, one can give representation varieties the structure of an affine algebraic variety by defining a representable functor [LM85, Proposition 1.2]

$$\begin{aligned} \mathrm{Rep}_n(\pi_1(\Sigma)) : \mathrm{Aff}^{\mathrm{op}} &\rightarrow \mathrm{Set} \\ \mathrm{Spec } B &\mapsto \mathrm{Hom}_{\mathrm{Gp}}(\pi_1(\Sigma), \mathrm{GL}_n(B)), \end{aligned}$$

where $\mathrm{Aff}^{\mathrm{op}}$ denotes the opposite of the category of affine $\mathbb{C}$ -schemes. By the adjunction

$$\mathrm{Gp} \begin{array}{c} \xrightarrow{\mathbb{C}[-]} \\ \xleftarrow{(-)^\times} \end{array} \mathrm{Alg}_{\mathbb{C}}$$

and the observation that $M_n(\mathbb{C})^\times = \mathrm{GL}_n(\mathbb{C})$, where $M_n(\mathbb{C})$ is the $\mathbb{C}$ -algebra of $(n \times n)$ -matrices with complex entries, one can reformulate this functor as

$$\mathrm{Rep}_n(\pi_1(\Sigma)) = \mathrm{Rep}_n(A)^{\mathrm{cl}} : \mathrm{Spec } B \mapsto \mathrm{Hom}_{\mathrm{Alg}_{\mathbb{C}}}(A, M_n(B)),$$

where $M_n(B)$ denotes $(n \times n)$ -matrices with entries in the $\mathbb{C}$ -algebra B . Similarly, the GL_n **character stack** is defined as

$$\mathrm{Rep}_n(\pi_1(\Sigma))/\mathrm{GL}_n = \mathrm{Rep}_n(A)^{\mathrm{cl}}/\mathrm{GL}_n.$$

On the other hand, the associated GL_n **character variety** is the GIT quotient $\mathrm{Rep}_n(A)^{\mathrm{cl}} // \mathrm{GL}_n$. In the algebraic setting, the derived enhancements of these objects, denoted $\mathrm{Rep}_n(A)$ and $\mathrm{Rep}_n(A) // \mathrm{GL}_n$, respectively, are already known in the literature, first constructed in [BKR13]. In this paper, we study the derived and stacky versions of these constructions, so that there is a correspondence

$$\mathrm{Rep}_n(A) / \mathrm{GL}_n \leftarrow (\mathrm{Rep}_n(A) / \mathrm{GL}_n)^{\mathrm{cl}} \rightarrow \mathrm{Rep}_n(A)^{\mathrm{cl}} // \mathrm{GL}_n \rightarrow \mathrm{Rep}_n(A) // \mathrm{GL}_n.$$

In the 1980's, Goldman defined in [Gol84] a symplectic structure on

$$\mathrm{Hom}(\pi_1(\Sigma), G) // G,$$

and its quantization has been of interest in knot theory, see e.g. [Tur91; AMR96; AMR98] for the classical story and [BBJ18; Jor23] for a modern formulation.

When A is finitely presented and commutative (so that $\mathrm{Spec} A$ is a quasi-projective scheme), the functor

$$B \mapsto \mathrm{Hom}_{\mathrm{Alg}_{\mathbb{C}}}(A, M_n(B))$$

is also the open subfunctor of the functor of points of the Quot scheme parametrizing length n quotients

$$q : (A \otimes_{\mathbb{C}} B)^n \rightarrow M$$

for which the induced map $B^n \rightarrow M$ is an isomorphism. We call this the framed locus and explain this in detail in Section 4.1. Like the specific case of Hilbert schemes, Quot schemes are a basic example of moduli problems in algebraic geometry and thus have been of interest since Grothendieck's original proof of their representability [Gro61; Nit05; HL10]. They are used as covers of the stack of coherent sheaves [HL10]; thus, they are also important in the theory of cohomological hall algebras [KV22; PS22].

1.2 Overview

This paper generalizes results of [BKR13] and [Yeu18] in several directions. We begin by placing a trace map of [BKR13] in a trace formalism context [Lur09b; HSS17; BN19]. Next, we generalize the study of such representation and character varieties for $G = \mathrm{GL}_n$ from $A \leadsto \mathrm{Rep}_n(A)$ to $X \leadsto \mathrm{Rep}_n(X)$, where X is now any perfect stack (see [BFN10] for what this means). In fact, our construction takes as input a category $\mathcal{C}$ of the form $\mathrm{QCoh}(X)$ or $A\text{-mod}$ equipped with the natural functor to $k\text{-mod}$ given by push-forward/restriction; in the latter case $\mathcal{C} = A\text{-mod}$, the choice of this restriction functor is equivalent to choosing a perfect generator, so this construction is not Morita invariant. In contrast, the closely related stack $\mathcal{M}_{\mathcal{C}}$ of [TV07] is Morita invariant. In the process, we construct a derived enhancement of an open subfunctor of the Quot scheme of n points which parametrizes quotients $q : \mathcal{O}_X^n \twoheadrightarrow \mathcal{E}$, where $\mathcal{E}$ is a S -flat family of length n torsion sheaves and the adjoint map to q , $\mathcal{O}_S^n \rightarrow p_{S*}\mathcal{E}$, is an isomorphism. Such derived Quot schemes have been constructed before, for projective X in [BKS19] and for quasi-projective X in [Adh22] (see also [CK99]); however, our construction differs significantly from theirs, as it uses a fiber of the stack of coherent sheaves as input instead of constructing the functor of points directly (see Section 4.1, in particular, Corollary 4.6, for details). On their common domains of definition, we expect all three constructions to agree. We check this explicitly for the construction in [Adh22] for smooth quasi-projective schemes in the appendix.

We show that in the case of a quasi-projective scheme X , there is a Fourier-Mukai integral transform between quasi-coherent sheaves on this derived Quot scheme and $\mathrm{QCoh}(X)$. This functor induces a trace

(in the sense of [HSS17; BN19]) map at the level of Hochschild homology which generalizes the trace map of [BKR13] from finitely presented commutative algebras to arbitrary quasi-projective schemes, and gives it an especially natural characterization: It arises due to the existence of a universal almost perfect complex on (the noncommutative analog of) $\mathrm{Spec} A \times \mathrm{Rep}_n(A)$, see Sections 2.2 and 4.2.

We next construct a generalization (in the sense of [Pan+11]) of the Goldman symplectic structure to (derived) character stacks equipped with a Calabi-Yau structure whose underlying classical truncation is of the form

$$\mathrm{Hom}(A, M_n)/\mathrm{GL}_n,$$

where A is a commutative smooth algebra. In [Pan+11], the authors construct a shifted symplectic structure on the derived moduli stack of maps $\mathrm{Map}(X_B, BG)$, where for a (compact, oriented) topological manifold X , X_B is the constant derived stack with value the singular complex of X . This moduli is the derived character stack of X ; moreover, in the case that X is a compact oriented 2-manifold, the shifted symplectic structure becomes 0-shifted, see [Pan+11, Theorem 2.5] and the examples following it. Further, this canonical symplectic structure arises from the choice of a symmetric bilinear form on the Lie algebra $\mathfrak{g}$ of G . This symplectic structure on the derived character variety of X recovers the Goldman symplectic structure of [Gol84] by pulling back to the classical character stack and then restricting to the appropriate smooth, classical locus [Pan+11; Saf16]. In Section 2.3 of [Pan+11], the authors also construct a symplectic structure on the moduli stack of perfect complexes on a smooth and proper Calabi-Yau scheme. It turns out that the properness assumption can be dropped [BD18], [BCS24, Theorem D and Lemma 3.7]. We use these foundations to construct our symplectic structure by pulling back this already existing one. Symplectic structures on derived character varieties have been constructed before in [Yeu18]. Therein, a generalization of derived representation varieties to categories is considered, where algebras are viewed as categories with one object. We did not take this approach, instead viewing algebras as categories (noncommutative spaces) via their categories of modules. Our constructions are functorial with respect to that view (as in e.g. [TV07]). We now give a more detailed overview of the results below.

1.3 Contents

Let k be a characteristic 0 field, fixed for the rest of this introduction (in fact, for the rest of this paper). In [BKR13], the authors construct and study a trace map

$$\mathrm{HC}_*(A) \rightarrow \mathbb{L}(A)_n$$

between the cyclic homology of an associative k -algebra A and its “representation homology,” $\mathbb{L}(A)_n$. This latter homology is by definition derived global functions on the derived n -dimensional representation variety of A , defined as follows (see Section 1.4 for context). As earlier in this introduction, let $\mathrm{Rep}_n(A)^{\mathrm{cl}}$ be the affine variety which represents the functor

$$B \mapsto \mathrm{Hom}_{\mathrm{Alg}_k}(A, M_n(B)),$$

where $M_n : \mathrm{CAlg}_k \rightarrow \mathrm{Alg}_k$ takes a commutative k -algebra B to the algebra of $(n \times n)$ -matrices with entries in B . This scheme is called the $(n$ -dimensional) **representation variety** of A . In particular, functions on it are the left adjoint of the functor M_n . All these constructions can now be derived as in [BKR13], to obtain a derived representation variety and a left derived functor $\mathbb{L}(-)_n$, and this latter functor is called the

representation homology. When $A = k[\pi_1(\Sigma)]$ for a surface Σ , this is a derived enhancement of the cohomology of the classical representation variety of A .

In Section 2, we place the above into a trace formalism context, (see [HSS17; BN19] for the trace formalism we use). Given algebras $A, \mathbb{L}(A)_n$, we show that the categories of modules over these algebras admit a natural “domain wall,” i.e., a Fourier-Mukai integral transform with a continuous right adjoint (explained in Sections 2 and 4.2), between them. This point of view gives a natural reason for why the trace map of [BKR13] should exist: It follows from the existence of a well-behaved integral kernel (Lemma 2.4 and Proposition 4.16). Indeed, by [BN19, Section 3.4], this integral transform gives rise to a map

$$\mathrm{HH}_*(A) \rightarrow \mathrm{HH}_*(\mathbb{L}(A)_n).$$

Because $\mathbb{L}(A)_n$ is a commutative algebra, it admits a map

$$\mathrm{HH}_*(\mathbb{L}(A)_n) \rightarrow \mathbb{L}(A)_n.$$

Moreover, the composition

$$\mathrm{HH}_*(A) \rightarrow \mathbb{L}(A)_n$$

is equivariant with respect to the natural S^1 -action on $\mathrm{HH}_*(A)$ and the trivial S^1 -action on $\mathbb{L}(A)_n$ [HSS17, Theorem 2.14]. Since S^1 -coinvariants form an adjunction [NS18, Section I.1]

$$(-)_{S^1} = \mathrm{colim}_{BS^1} : \mathrm{Vect}^{S^1} \rightarrow \mathrm{Vect} : \mathrm{triv},$$

where triv equips a vector space with the trivial S^1 -action and $\mathrm{Vect}^{S^1} := \mathrm{Map}(BS^1, \mathrm{Vect})$ denotes vector spaces with S^1 -actions, we obtain a map

$$\mathrm{HC}_*(A) \rightarrow \mathbb{L}(A)_n.$$

Theorem 1.1 (Theorem 2.8 in the text). If A is a connective associative algebra, then the induced map

$$\mathrm{HC}_*(A) \rightarrow \mathbb{L}(A)_n$$

is the trace map of [BKR13].

In Section 3, we extend this definition of derived representation varieties from algebras A to perfect stacks X (see [BFN10] for what this means). Let $\mathcal{X}_{A\text{-mod}}$ denote the stack which represents the functor $\mathrm{Spec} B \mapsto \widehat{(A \otimes B)\text{-mod}}$, where $\widetilde{\mathcal{C}}$ denotes the maximal subgroupoid of a category $\mathcal{C}$. Similarly, let $\mathcal{X}_{\mathrm{QCoh}(X)}$ for a perfect stack X denote the stack of quasi-coherent complexes on X , defined in Section 3. We equip both of these stacks with natural maps to the stack of quasi-coherent complexes on a point, $\mathrm{Vect} := k\text{-mod}$, defined by pushforward (Lemma 3.3). Using either of these categories together with their maps to Vect , we define a derived stack

$$\mathcal{R}ep_n(\mathcal{C}) := \mathcal{X}_{\mathcal{C}} \times_{\mathrm{Vect}} \{k^n\},$$

where the map $\{k^n\} \rightarrow \mathrm{Vect}$ classifies k^n . Because the map $\mathcal{X}_{\mathcal{C}} \rightarrow \mathrm{Vect}$ is part of the construction of this moduli, this construction is *not* Morita invariant. We show (Theorem 3.7) that the extension, which we call

the “ n -dimensional representation moduli,” agrees with representation varieties in the case of $\mathcal{C} = A\text{-mod}$. We compute this stack in the case that $\mathcal{C} = \text{QCoh}(X)$ for $X = \mathbb{P}^1$ and $X = BG$ with G a reductive algebraic group.

Further, we show in Section 4 that for $\mathcal{C} = \text{QCoh}(X)$ with X a classical quasi-projective scheme, this stack admits a substack of “coherent representations,” denoted $\mathcal{R}ep_n^{\text{coh}}(X)$. In Theorem 4.5, we prove that this substack is 1-Artin, locally almost of finite type, and with affine diagonal. Finally, we show in Corollary 4.6 that the stack of coherent representations admits a substack

$$\text{Quot}_{\mathcal{O}_X^n}^{n,\text{fr}}(X) \subset \mathcal{R}ep_n^{\text{coh}}(X) \subset \mathcal{R}ep_n(\text{QCoh}(X))$$

which is a derived scheme almost of finite type and with affine diagonal whose classical truncation is the open subfunctor of the Quot scheme of points which parametrizes quotients $q : \mathcal{O}_{X \times S}^n \rightarrow \mathcal{E}$ for which the induced map $\mathcal{O}_S^n \rightarrow p_{S*}\mathcal{E}$ is an isomorphism. On affine schemes of finite presentation, all of these stacks agree.

Theorem 1.2 (Proposition 4.8 and Corollary 4.10 in the text). Let $X = \text{Spec } A$ be an affine scheme of finite presentation. Then

$$\text{Quot}_{\mathcal{O}_X^n}^{n,\text{fr}}(X) \cong \mathcal{R}ep_n^{\text{coh}}(X) \cong \mathcal{R}ep_n(\text{QCoh}(X)) \cong \text{Rep}_n(A).$$

Remark 1.3. By the above theorem, this Quot scheme on smooth and affine $X = \text{Spec } A$ is equivalent to the corresponding open subfunctor of the derived enhancement of this Quot scheme defined in [Adh22] (see Example 4.1, Proposition 4.2, and Definition 3.3 of *op. cit.*). In fact, these also match when X is smooth quasi-projective by Proposition A.1.

Now, let X be a (not necessarily smooth) quasi-projective scheme. We show that the scheme $\text{Quot}_{\mathcal{O}_X^n}^{n,\text{fr}}(X) \times X$ admits an almost perfect complex $\mathcal{R}$ called the “universal representation.” This complex is a generalization of the domain wall which defines the map of Theorem 1.1.

Theorem 1.4 (Proposition 4.16 in the text). The sheaf $\mathcal{R}$ defines a **domain wall** between the partially extended field theories defined by

$$\text{QCoh}(X) \quad \text{and} \quad \text{QCoh}(\text{Quot}_{\mathcal{O}_X^n}^{n,\text{fr}}(X)),$$

meaning that the Fourier-Mukai integral transform it defines,

$$\begin{aligned} \Phi_{\mathcal{R}} : \text{QCoh}(X) &\rightarrow \text{QCoh}(\text{Quot}_{\mathcal{O}_X^n}^{n,\text{fr}}(X)) \\ \mathcal{F} &\mapsto p_{\text{Quot}_{\mathcal{O}_X^n}^{n,\text{fr}}(X)}^*(p_X^* \mathcal{F} \otimes \mathcal{R}) \end{aligned}$$

has a continuous right adjoint. Here, $p_X, p_{\text{Quot}_{\mathcal{O}_X^n}^{n,\text{fr}}(X)}$ are the respective projections from $X \times \text{Quot}_{\mathcal{O}_X^n}^{n,\text{fr}}(X)$.

In particular, this theorem implies that in addition to functors between the respective categories, $\mathcal{R}$ induces a map on Hochschild homology by a theorem of [BN19], which is moreover S^1 -equivariant.

Theorem 1.5 (Theorem 4.17 in the text). If $X = \text{Spec } A$ is a finitely presented affine scheme, then the domain wall defined by $\mathcal{R}$ precisely recovers the domain wall which gives rise to the trace map of [BKR13].

Thus, this generalizes the story of [BKR13] by initiating the study of analogous trace maps for arbitrary classical quasi-projective schemes.

Let $\mathrm{Coh}_0^n(X)$ denote the derived stack of length n torsion sheaves on a classical quasi-projective scheme X (as defined in [PS22] and reviewed in §4). There is a canonical projection map

$$\mathrm{Quot}_{\mathcal{O}_X^n}^{n,\mathrm{fr}}(X) \rightarrow \mathrm{Coh}_0^n(X),$$

which informally can be described as sending a quotient $\mathcal{O}_X^n \twoheadrightarrow \mathcal{E}$, where $\mathcal{E}$ is length n torsion, to $\mathcal{E}$ itself. We show (Proposition 5.1) that this map is a GL_n -torsor; thus, $\mathrm{Coh}_0^n(X)$ is the analog of the derived character variety $\mathrm{Rep}_n(A)/\mathrm{GL}_n$ for non-affine schemes. When X is furthermore smooth, this stack admits a formally étale map

$$f : \mathrm{Coh}_0^n(X) \rightarrow \mathrm{Perf}_{\mathrm{prop}}(X)$$

to the stack of perfect complexes with proper support. In the case that X is Calabi-Yau of dimension d , this latter stack admits a $(2 - d)$ -shifted symplectic structure by [BD18].

Theorem 1.6 (Theorem 5.2 in the text). Let X be a smooth classical quasi-projective scheme equipped with a trivialization $\mathcal{O}_X \cong K_X$ of its canonical bundle. Then $\mathrm{Coh}_0^n(X)$ admits a $(2 - d)$ -shifted symplectic structure induced from the one on $\mathrm{Perf}_{\mathrm{prop}}(X)$ by pullback along f .

This theorem has been shown for projective X in [BKS19]; as far as we know, the extension to quasi-projective X in the context of framed representation moduli is new. Moreover, the above extension to quasi-projective schemes thus defines a shifted symplectic structure on the derived character variety

$$\mathrm{Rep}_n(A)/\mathrm{GL}_n$$

when A is a smooth commutative algebra of finite presentation and the stack is equipped with a Calabi-Yau structure. We expect that this symplectic structure agrees with the structure obtained in [Yeu18] on derived character varieties.

1.4 Notation and Conventions

All categories considered are without further comment ∞ -categories and all functors derived functors/functors between ∞ -categories [Lur09a]. For example, Hom denotes the derived functor $\mathbb{R}\mathrm{Hom}$, $\otimes$ denotes the derived functor $\otimes^{\mathbb{L}}$, etc. Schemes are likewise assumed derived (see [GR17] for an introduction). Any object, map, or functor will be explicitly called “classical” if it is supposed to be the underived version. We always work over a fixed field k of characteristic 0, and our gradings will always be cohomological. All algebras are assumed to be unital.

Our running assumptions on stacks is that they are locally almost of finite type (laft) with affine diagonal. The exception is in Section 3, where we use the big stacks of quasi-coherent complexes (which are not laft). If X is a scheme or stack, we denote by X^{cl} the classical truncation [GR17, Chapter 2]. We use sans serif font for names of categories; e.g., $\mathrm{Perf}(X)$ denotes the $(\infty, 1)$ -category of perfect complexes on a scheme or stack X , and we use ordinary font for names of stacks; e.g., $\mathrm{Perf}(X)$ is the stack of perfect complexes on X .

1.5 Glossary of Categories

Since k is assumed to be a characteristic 0 field throughout, the category $\mathbf{CAlg}_k$ (often denoted simply $\mathbf{CAlg}$) of commutative differential graded k -algebras is our model for (the opposite to) the category of derived affine schemes, and we write $\mathbf{Aff}$ for its opposite category.

We denote by $\mathbf{Alg}_k$ the category of associative differential graded k -algebras.

For $A \in \mathbf{Alg}_k$, we let $\mathbf{LMod}_A$ denote the stable category of left A -modules. Similarly, $\mathbf{RMod}_A$ denotes the stable category of right A -modules. In particular, if $A \in \mathbf{CAlg}_k \subset \mathbf{Alg}_k$, then we denote $\mathbf{LMod}_A =: A\text{-mod}$ and $\mathbf{QCoh}(\mathrm{Spec} A) := A\text{-mod}$. The subcategory of compact objects is denoted by $\mathbf{LPerf}_A$; in the case that A is commutative, we write $\mathbf{Perf}(\mathrm{Spec} A) := A\text{-perf}$. The category $A\text{-mod}$ comes equipped with a canonical t -structure defined in [GR17, Chapter 2 Section 1.2].

For a prestack X , we let $\mathbf{QCoh}(X) := \lim_{\mathrm{Spec} A \rightarrow X} A\text{-mod}$, where the limit is taken over all derived affine schemes mapping to X . We also have the category $\mathbf{Perf}(X) := \lim_{\mathrm{Spec} A \rightarrow X} A\text{-perf}$.

We denote by $\mathbf{Mor}_k$ the Morita $(\infty, 2)$ -category of k -algebras, bimodules, and intertwiners of [Hau17]. Informally, objects of this category are algebras, 1-morphisms $A \rightarrow B$ are (A, B) -bimodules, and 2-morphisms are intertwining bimodule-homomorphisms.

We denote by $\mathbf{Pr}^{\mathrm{L}}$ the $(\infty, 2)$ -category of presentable $(\infty, 1)$ -categories, colimit preserving functors, and natural transformations as in [Lur17].

We let $\mathbf{Cat}$ denote the $(\infty, 2)$ -category of all $(\infty, 1)$ -categories, colimit preserving functors, and natural transformations.

The category of spaces, equivalently, of ∞ -groupoids, is denoted $\mathbf{Spc}$. This category admits a natural inclusion $\mathbf{Spc} \hookrightarrow \mathbf{Cat}$.

1.6 Acknowledgments

I would like to thank my advisor, David Nadler, for helping me with the TQFT portion of this project and for his continued support, encouragement, and advice during this project. This paper would not have been completed without his support. I would also like to thank Yuri Berest for suggesting the portion on extending representation varieties to stacks, as it led to the most interesting (in my opinion) results. Ansuman Bardalai taught me much of the basic (and not so basic) algebraic geometry that I used during the completion of this project and answered many of my algebro-geometric confusions related to this project. I also benefited from conversations with Daigo Ito, Kabir Kapoor, and Theo Johnson-Freyd related to this project.

2 Domain Walls

2.1 Dualizability

Consider the Morita $(\infty, 2)$ -category of k -algebras, bimodules, and intertwiners, $\mathbf{Mor}_k$, which is defined in e.g. [Hau17]. We will say simply “2-category” in the sequel, but we always mean “ $(\infty, 2)$ -category.” We now record some basic results implied by [GS18, Theorem 5.1].

Proposition 2.1 (Gwilliam-Scheimbauer). Every object $A \in \mathbf{Mor}_k$ is dualizable with dual A^{op} , coevaluation ${}_k A_{A \otimes A^{\mathrm{op}}}$, and evaluation ${}_{A^{\mathrm{op}} \otimes A} A_k$.

Since we're in a 2-category, it is natural to ask for dualizable 1-morphisms in addition to objects. For this, we require unit and counit 2-morphisms which exhibit a given 1-morphism as dualizable [Lur09b]. For example, a 1-morphism $M : A \rightarrow B$ is right dualizable with right dual M^R if there exist unit and counit maps, $\eta : A \rightarrow M \otimes_B M^R$ and $\varepsilon : M^R \otimes_A M \rightarrow B$, such that the following compositions compose to id_M and id_{M^R} , respectively [Lur17, Definition 4.6.1.1].

$$\begin{aligned} M &= A \otimes_A M \xrightarrow{\eta \otimes \text{id}_M} M \otimes_B M^R \otimes_A M \xrightarrow{\text{id}_M \otimes \varepsilon} M \otimes_B B = M \\ M^R &= M^R \otimes_A A \xrightarrow{\text{id}_{M^R} \otimes \eta} M^R \otimes_A M \otimes_B M^R \xrightarrow{\varepsilon \otimes \text{id}_{M^R}} B \otimes_B M^R = M^R. \end{aligned}$$

Lemma 2.2. Let M be an (A, B) -bimodule. Then there are canonical evaluation maps

$$\begin{aligned} \text{Hom}_B(M, B) \otimes_A M &\rightarrow B \\ M \otimes_B \text{Hom}_A(M, A) &\rightarrow A. \end{aligned}$$

Proof. The functor

$$- \otimes_A M : \text{RMod}_A \rightarrow \text{RMod}_B$$

has a right adjoint $\text{Hom}_B(M, -)$, and the counit of the adjunction evaluated at B is precisely

$$\text{Hom}_B(M, B) \otimes_A M \rightarrow B.$$

The proof of the existence of the evaluation

$$M \otimes_B \text{Hom}_A(M, A) \rightarrow A$$

is similar. □

Proposition 2.3 (Lurie). Let M be an (A, B) -bimodule viewed as a 1-morphism in Mor_k . Then M has a left (resp. right) dual if and only if it is a perfect A -module (resp. B -module) with left (resp. right) dual $M^L = \text{Hom}_A(M, A)$ (resp. $M^R = \text{Hom}_B(M, B)$).

Proof. Propositions 4.6.2.1 and 4.6.2.13 of [Lur17] state that M is left dualizable as an (A, B) -bimodule if and only if the restriction of M to an A -module is left dualizable. We then observe that dualizable A -modules are precisely the perfect A -modules by [Lur17, Proposition 7.2.4.2]. The existence of the left dual implies that the functor

$$- \otimes_A M : \text{RMod}_A \rightarrow \text{RMod}_B$$

has a left adjoint given by $- \otimes_B M^L$ by Proposition 4.6.2.1 of *op. cit.* Since M is perfect as a left A -module, then (the dual to) Proposition 7.2.4.4 of *op. cit.* shows that

$$- \otimes_A M^L : \text{RMod}_A \rightarrow k\text{-mod}$$

is given by the functor corepresented by M , i.e. by $\text{Hom}_A(M, -)$. This identifies

$$A \otimes_A M^L \cong M^L = \text{Hom}_A(M, A).$$

The proof for M^R is similar. □

2.2 Trace Maps I

By [BN19, Section 3.4], any continuous (that is, colimit-preserving) morphism $M : A \rightarrow B$ of dualizable objects in Mor_k which is furthermore equipped with a continuous right adjoint defines a map

$$\text{HH}_*(A) \rightarrow \text{HH}_*(B)$$

between the Hochschild homologies of A, B . We call such a morphism M , which is a dualizable (A, B) -bimodule, a **domain wall** between the categories $A\text{-mod}$ and $B\text{-mod}$; i.e., M defines a colimit preserving functor between the categories $A\text{-mod}$ and $B\text{-mod}$ which moreover has a colimit preserving right adjoint. For example, we have a functor

$$\begin{aligned} M_{\text{pt}} : A\text{-mod} &\rightarrow B\text{-mod} \\ K &\mapsto K \otimes_A M. \end{aligned}$$

The map

$$M_{S^1} : \text{HH}_*(A) \rightarrow \text{HH}_*(B)$$

is defined as follows. (Right) dualizability ensures we have unit and counit maps

$$\begin{aligned} \eta : A &\rightarrow M \otimes_B M^R \\ \varepsilon : M^R \otimes_A M &\rightarrow B, \end{aligned}$$

where $M^R = \text{Hom}_B(M, B)$ is the right dual of M . We thus obtain a map of k -vector spaces

$$\text{HH}_*(A) \rightarrow A \underset{A \otimes A^{\text{op}}}{\otimes} M \otimes_B M^R = B \underset{B \otimes B^{\text{op}}}{\otimes} M^R \otimes_A M \rightarrow \text{HH}_*(B). \quad (2.1)$$

We resolve A and B by cofibrant resolutions $QA \rightarrow A$ and $QB \rightarrow B$. Since quasi-isomorphisms induce Morita equivalences (via extension of scalars), we can consider an (A, B) -bimodule M to be a (QA, QB) -bimodule with a canonical bimodule structure. Thus, the above maps descend to maps in the homotopy category of k -algebras, hAlg :

$$\text{HH}_*(QA) \rightarrow QA \underset{QA \otimes QA^{\text{op}}}{\otimes} M \otimes_{QB} M^R = QB \underset{QB \otimes QB^{\text{op}}}{\otimes} M^R \otimes_{QA} M \rightarrow \text{HH}_*(QB),$$

where now $M^R = \text{Hom}_{QB}(M, QB)$. Moreover, this map admits a canonical S^1 -equivariant refinement by [HSS17, Theorem 2.14].

We claim that the trace map studied in [BKR13] exactly arises from such a domain wall construction.

Lemma 2.4. Let $A \in \text{Alg}$ be an associative algebra over k , and let $QA \rightarrow A$ be a semi-free dg, i.e. cofibrant, resolution of A . Denote by $B = \mathbb{L}(A)_n$ its derived n -dimensional representation homology, i.e. the left derived functor of $(-)_n$ applied to A , and let $M = B^n$. Then M is a (QA, B) -bimodule.

Proof. The unit of the adjunction $((-)_n, M_n)$ is a map of algebras $QA \rightarrow M_n(B) = \text{End}_B(B^n)$, which is a (QA, B) -bimodule structure on B^n . $\square$

Remark 2.5. Since the quasi-isomorphism $QA \rightarrow A$ induces a Morita equivalence, we see that M has an (A, B) -bimodule structure.

Since M is a perfect B -module, it is right dualizable and hence defines a domain wall between the B -models associated to QA and B . By Proposition 2.3 it has a right dual given by $M^R = \text{Hom}_B(M, B)$, and

$$M \otimes_B M^R \cong \text{End}_B(M) \cong M_n(B).$$

Proposition 2.6. In the setting of Lemma 2.4, the map

$$QA \otimes_{QA \otimes QA^{\text{op}}} QA \rightarrow QA \otimes_{QA \otimes QA^{\text{op}}} M \otimes_B M^R$$

induced by the domain wall given by M is exactly $\text{id}_{QA} \otimes \eta$, where η is the unit of the adjunction $((-)_n, M_n)$.

Proof. M is right dualizable with right dual M^R . Thus, M^R , viewed as a 1-morphism in the Morita 2-category, has unit given by

$$QA \rightarrow M \otimes_B M^R,$$

and this is the unit of the adjunction

$$- \otimes_{QA} M \dashv - \otimes_B M^R.$$

Under the isomorphism

$$M \otimes_B M^R \cong \text{End}_B(M),$$

this unit is the map

$$QA \rightarrow \text{End}_B(M)$$

which defines the left QA -module structure on $\text{End}_B(M)$. On the other hand, the left QA -action on

$$\text{End}_B(M) \cong M_n((QA)_n)$$

is induced by the unit of the adjunction $((-)_n, M_n)$,

$$\eta_{QA} : QA \rightarrow M_n((QA)_n) \cong \text{End}_B(M).$$

Thus, we see that the units of the two adjunctions in question are the same, so the map in the statement of the Proposition is $\text{id}_{QA} \otimes \eta_{QA}$. $\square$

Let $B \in \mathbf{CAlg}$ be a commutative algebra, and let $\text{Map}(S^1, \text{Spec } B) =: \mathcal{L} \text{Spec } B$ be the loop space of $X = \text{Spec } B$. There is a canonical inclusion map $X \rightarrow \mathcal{L} X$ given by constant loops, which corresponds to a map of algebras

$$\text{HH}_*(B) = \mathcal{O}(\mathcal{L} \text{Spec}(B)) \rightarrow B. \quad (2.2)$$

Further, this map is equivariant with respect to the S^1 -actions on $\mathcal{L} X$ and X , where S^1 acts on X trivially.

Proposition 2.7. Let $\text{Alg}^{\leq 0} \subset \text{Alg}$ denote the full subcategory of connective associative algebras. Consider the functors

$$\text{HC}_*, \mathbb{L}(-)_n : \text{Alg}^{\leq 0} \rightarrow \text{Vect},$$

where in this case $\mathbb{L}(A)_n$ denotes the underlying vector space of the connective commutative algebra. Suppose given natural transformations

$$\alpha, \beta : \text{HC}_* \Rightarrow \mathbb{L}(-)_n$$

such that for every finite-dimensional vector space V with tensor algebra $\mathcal{T}(V) := \bigoplus_n V^{\otimes n} \in \text{Alg}^{\leq 0}$, the maps

$$H^0(\text{HC}_*(\mathcal{T}(V))) \rightarrow (\mathcal{T}(V))_n$$

induced by $\alpha_{\mathcal{T}(V)}$ and $\beta_{\mathcal{T}(V)}$ agree. Then

$$\alpha \cong \beta.$$

Proof. Both HC_* and $\mathbb{L}(-)_n$ preserve sifted colimits. Indeed, Hochschild homology is computed via the geometric realization of the cyclic bar construction, whose composition preserves sifted colimits. Similarly, S^1 -coinvariants is a colimit; hence, HC_* preserves sifted colimits. On the other hand, the functor $\mathbb{L}(-)_n$ is a left adjoint, and the forgetful functor $\text{CAlg}^{\leq 0} \rightarrow \text{Vect}$ also preserves sifted colimits.

Let $\text{Alg}^0 \subset \text{Alg}^{\leq 0}$ denote the subcategory spanned by algebras of the form $\mathcal{T}(V)$ with V a finite-dimensional vector space concentrated in degree 0. By the remarks after Corollary 7.1.4.17 of [Lur17], the category $\text{Alg}^{\leq 0} \cong \mathcal{P}_{\Sigma}(\text{Alg}^0)$, which means that every connective associative algebra can be expressed as a sifted colimit of algebras of the form $\mathcal{T}(V)$. Thus, it is sufficient to show that $\alpha \cong \beta$ on this full subcategory. Since $\mathcal{T}(V)$ is cofibrant for any V , we see that

$$\mathbb{L}(\mathcal{T}(V))_n = (\mathcal{T}(V))_n.$$

Moreover, the underlying vector space of this algebra is in the heart of Vect . On the other hand, $\text{HC}(\mathcal{T}(V))$ is connective, so that we can identify

$$\text{Hom}(\text{HC}_*(\mathcal{T}(V)), (\mathcal{T}(V))_n) \cong \text{Hom}^0(H^0(\text{HC}_*(\mathcal{T}(V))), (\mathcal{T}(V))_n),$$

where the RHS is the inner Hom in $\text{Vect}^{\heartsuit}$; in particular, it is a discrete space. Since we also have

$$H^0(\text{HC}_*(\mathcal{T}(V))) = \mathcal{T}(V)/[\mathcal{T}(V), \mathcal{T}(V)],$$

we see that a point of $\text{Hom}(\text{HC}_*(\mathcal{T}(V)), (\mathcal{T}(V))_n)$ is uniquely determined by a map

$$\mathcal{T}(V)/[\mathcal{T}(V), \mathcal{T}(V)] \rightarrow (\mathcal{T}(V))_n.$$

Now, the space of natural transformations between HC_* and $\mathbb{L}(-)_n$ is a limit of the mapping spaces $\text{Hom}(\text{HC}_*(\mathcal{T}(V)), \mathcal{T}(W))$ which are all discrete because the source is connective and the target is in the heart. Thus, the limit is 0-truncated, so the above identification of points is sufficient to identify the natural transformations. Since by assumption the maps induced by $\alpha_{\mathcal{T}(V)}$ and $\beta_{\mathcal{T}(V)}$ agree and are of this form, we conclude that

$$\alpha \cong \beta.$$

□

Let $A \in \text{Alg}^{\leq 0}$ be a connective associative algebra, and set

$$B := \mathbb{L}(A)_n.$$

Denote the unit of the adjunction $(\mathbb{L}(-)_n, M_n)$ by $\eta : A \rightarrow M_n(B)$, and denote $M := B^n$ the (A, B) -bimodule of Lemma 2.4. Since M is perfect as a right B -module, it is right dualizable and hence induces, by [HSS17, Theorem 2.14], a canonical S^1 -equivariant map

$$\text{HH}_*(A) \rightarrow \text{HH}_*(B).$$

Composing this with the canonical map $\text{HH}_*(B) \rightarrow B$ given by (2.2), we obtain an S^1 -equivariant map

$$\text{HH}_*(A) \rightarrow \text{triv}(B), \tag{2.3}$$

where $\text{triv}(B)$ denotes the underlying vector space of B equipped with the trivial S^1 -action.

Theorem 2.8. Under the adjunction

$$(-)_{S^1} : \text{Vect}^{S^1} \rightarrow \text{Vect} : \text{triv},$$

the map

$$\text{HC}_*(A) \rightarrow B$$

induced by (2.3) is the trace map of [BKR13].

Proof. We first claim that the map

$$t_A : \text{HC}_*(A) \rightarrow B$$

induces a natural transformation

$$t : \text{HC}_* \Rightarrow \mathbb{L}(-)_n.$$

Indeed, by applying to η the functor $\text{Alg} \rightarrow \text{Mor}$ which sends maps to base change bimodules, we see that the assignment $A \rightsquigarrow (A \xrightarrow{M} \mathbb{L}(A)_n)$ assembles, by functoriality of extension of scalars, into a coherent family of 1-morphisms $A \rightarrow M_n(B)$, because in Mor , we have the Morita equivalence $M_n(B) \cong B$ which is natural in B (since $M_n(B) \cong M_n(k) \otimes B$). Furthermore, by naturality of the trace functor of [HSS17, Theorem 2.14], we thus obtain an S^1 -equivariant natural transformation

$$\text{HH}_* \rightarrow \mathbb{L}(-)_n,$$

which thus induces the natural transformation t . The trace map θ of [BKR13] is also a natural transformation by the remarks around Equations 4.3 and 4.4 of *op. cit.*; moreover, these are natural transformations with the same source and target, since by [HSS17, Proposition 4.24 and Remark 6.12], the S^1 -structure on the sources are the same, functorially in A . Thus, by Proposition 2.7, it suffices to show that the two natural transformations t and θ agree on zeroth cyclic homology of algebras of the form $A = \mathcal{T}(V)$ with V a finite-dimensional vector space. Letting $B = (\mathcal{T}(V))_n = \mathbb{L}(\mathcal{T}(V))_n$ and $\eta : \mathcal{T}(V) \rightarrow M_n(B)$, we have

the associated bimodule $M = B^n$. We can now work element by element. Let $e_1, \dots, e_n$ be the standard basis of M with dual basis $e_1^\vee, \dots, e_n^\vee$ in $M^R = \text{Hom}_B(M, B)$. The map

$$\mathcal{T}(V)/[\mathcal{T}(V), \mathcal{T}(V)] \rightarrow (\mathcal{T}(V))_n$$

induced by M is explicitly

$$\bar{a} \mapsto \sum e_i^\vee(\eta(a)e_i),$$

which is just the ordinary trace

$$\sum e_i^\vee(\eta(a)e_i) = \text{tr}(\eta(a)).$$

Indeed, we can calculate this at the chain level as follows. Resolve M as an (A, B) -bimodule by a cofibrant resolution $K \rightarrow M$. We need only identify the map in question in degree 0. There, the first map sends

$$\begin{aligned} A/[A, A] &\rightarrow H^0(A \otimes_{A \otimes A^{\text{op}}} M_n(B)) = M_n(B)/[A, M_n(B)] \\ \bar{a} &\mapsto \bar{a} \cdot 1. \end{aligned}$$

This is then sent to

$$\begin{aligned} M_n(B)/[A, M_n(B)] &\rightarrow A \otimes_{A \otimes A^{\text{op}}} K^{-m} \otimes_B \text{Hom}_B^m(K, B) \\ \bar{a} &\mapsto 1 \otimes \bar{a}k \otimes \varphi, \end{aligned}$$

where $k \otimes \varphi \in K^{-m} \otimes_B \text{Hom}_B^m(K, B)$ corresponds to id_K under the isomorphism

$$M \otimes_B M^R \cong \text{End}_B(M).$$

Now, under the isomorphism

$$A \otimes_{A \otimes A^{\text{op}}} M \otimes_B M^R \cong M^R \otimes_A M \otimes_{B \otimes B} B,$$

we see that $1 \otimes \bar{a}k \otimes \varphi$ is sent to

$$\begin{aligned} A \otimes_{A \otimes A^{\text{op}}} K^m \otimes_B \text{Hom}_B^{-m}(K, B) &\rightarrow \text{Hom}_B^{-m}(K, B) \otimes_A K^m \otimes_{B \otimes B} \mathcal{B}^0 \\ 1 \otimes \bar{a}k \otimes \varphi &\mapsto \varphi \otimes \bar{a}k \otimes 1, \end{aligned}$$

where $\mathcal{B} \rightarrow B$ is a cofibrant resolution of B as a $(B \otimes B)$ -module. Finally, this is sent via $\varepsilon \otimes \text{id}_B$, where ε is the counit duality datum for M , to the element

$$\varphi(\bar{a}k) \in B.$$

Now, since $M = B^n$ is free over B , $\text{End}_B(K) = \text{End}_B(M) = M_n(B)$. Next, the endomorphism of K which corresponds to $\bar{a}k \otimes \varphi \in K^m \otimes_B \text{Hom}_B^{-m}(K, B)$ is $\bar{a}k\varphi$, and this has trace by definition given by $\varphi(\bar{a}k)$.

Moreover, since B is commutative, $B/[B, B] = B$, so the constant loop map on zeroth Hochschild homology is the identity. Thus, $t_{\mathcal{T}(V)}$ induces the map

$$\bar{a} \mapsto \text{tr}(\eta(a)),$$

which is precisely the definition of the Berest-Khachatryan-Ramadoss trace, as in e.g. [BR22, Section 3]. $\square$

3 Extension to Stacks

3.1 Definition

Let $\mathcal{C}$ be a category. Note that $\mathcal{C}$ defines a moduli functor $\mathcal{X}_{\mathcal{C}}$, which is the composition of the functor

$$\begin{aligned} \text{Aff}^{\text{op}} &\rightarrow \text{Cat} \\ S &\mapsto \mathcal{C} \otimes \text{QCoh}(S) \end{aligned}$$

with the right adjoint to the inclusion $\text{Spc} \hookrightarrow \text{Cat}$. Informally, this functor can be described as sending a map of affines $f : S' \rightarrow S$ to

$$\widetilde{\mathcal{C} \otimes \text{QCoh}(S)} \xrightarrow{\text{id}_{\mathcal{C}} \otimes f^*} \widetilde{\mathcal{C} \otimes \text{QCoh}(S')},$$

where for a category $\mathcal{A}$, $\widetilde{\mathcal{A}}$ denotes the maximal subgroupoid of $\mathcal{A}$.

Remark 3.1. This functor has as a subfunctor the perhaps more familiar moduli functor of pseudo-perfect complexes $\mathcal{M}_{\mathcal{C}}$ of [TV07].

Proposition 3.2. If $\mathcal{C} = \text{QCoh}(X)$, the category of quasi-coherent sheaves on a prestack X , or $\mathcal{C} = A\text{-mod}$ is modules over an associative algebra A , then $\mathcal{X}_{\mathcal{C}}$ is a stack.

Proof. Let A be an associative algebra and $\mathcal{C} = A\text{-mod}$. Given a faithfully flat map of commutative algebras $B \rightarrow C$, we can form its Amitsur complex $(C/B)_{\bullet}$, which is by definition the underlying cosimplicial algebra of the Čech nerve of $\text{Spec } C \rightarrow \text{Spec } B$. To prove descent for $\mathcal{X}_{A\text{-mod}}$, it thus suffices to provide an equivalence

$$\lim(A\text{-mod} \otimes (C/B)_{\bullet}\text{-mod}) \cong A\text{-mod} \otimes (\lim(C/B)_{\bullet})\text{-mod}.$$

By faithfully flat descent, we have

$$(\lim(C/B)_{\bullet})\text{-mod} \cong \lim((C/B)_{\bullet}\text{-mod}).$$

Hence, it suffices to show that the Lurie tensor product with $A\text{-mod}$ commutes with this limit. By construction, this tensor can be identified with colimit preserving functors between stable categories, since $A\text{-mod}$ is dualizable in Pr^{L} (with dual $A\text{-mod}^{\vee} = A^{\text{op}}\text{-mod}$):

$$A\text{-mod} \otimes (\lim(C/B)_{\bullet})\text{-mod} \cong \text{Map}_{\text{Pr}^{\text{L}}}((A\text{-mod})^{\vee}, (\lim(C/B)_{\bullet})\text{-mod}).$$

Since the Map functor commutes with limits in the second variable, we have the result.

If X is a prestack, then for $\mathcal{C} = \text{QCoh}(X)$, $\mathcal{X}_{\mathcal{C}}$ is given by

$$S \mapsto \widetilde{\text{QCoh}(X \times S)},$$

which is the composition of the functor $\widetilde{\text{QCoh}(-)}$ with $X \times -$ by [GR17, Chapter 3 Proposition 3.5.3]. Now, given a cover $S' \rightarrow S$, $X \times -$ applied to the Čech nerve of this map is also a Čech nerve (of the map $X \times S' \rightarrow X \times S$). Since the functor $\text{QCoh}(-)$ satisfies descent and passing to the maximal subgroupoid is a right adjoint, the composite $\mathcal{X}_{\mathcal{C}}$ satisfies descent, too. $\square$

Lemma 3.3. Let $\text{Vect} := \mathcal{X}_{k\text{-mod}}$ denote the stack of quasi-coherent complexes, so $\text{Vect}(S) = \text{QCoh}(S)$. If $\mathcal{C} = \text{QCoh}(X)$ is the category of quasi-coherent sheaves on a perfect stack or if $\mathcal{C} = A\text{-mod}$ is modules over an associative algebra $A \in \text{Alg}_k$, we have a natural transformation

$$\mathcal{X}_{\mathcal{C}} \rightarrow \text{Vect}.$$

Remark 3.4. In the case of $\mathcal{C} = A\text{-mod}$, this natural transformation *depends on the presentation* of $\mathcal{C}$ as a module category. When we write $\mathcal{C} = A\text{-mod}$, we always implicitly choose such a presentation.

Proof. Let $\mathcal{C} = A\text{-mod}$ be modules over an associative algebra. We claim restriction of modules defines a natural transformation

$$\mathcal{X}_{\mathcal{C}} \rightarrow \text{Vect}.$$

Given a map $B \rightarrow C$ of algebras, we need to show that the following diagram commutes, where the horizontal arrows are restriction and the vertical arrows base change.

$$\begin{array}{ccc} (A \otimes B)\text{-mod} & \longrightarrow & B\text{-mod} \\ \downarrow & & \downarrow \\ (A \otimes C)\text{-mod} & \longrightarrow & C\text{-mod} \end{array}.$$

Indeed, this follows from the isomorphism

$$M \otimes_{A \otimes B} A \otimes C \cong M \otimes_B C.$$

Now, let $\mathcal{C} = \text{QCoh}(X)$ for X a perfect stack. Consider the assignment

$$\begin{aligned} \mathcal{X}_{\mathcal{C}}(S) &= \text{QCoh}(X \times S) \rightarrow \text{Vect}(S) = \text{QCoh}(S) \\ \mathcal{F} &\mapsto p_* \mathcal{F}, \end{aligned}$$

where $p : X \times S \rightarrow S$ is the projection. We claim that the assignment $\mathcal{F} \mapsto p_* \mathcal{F}$ is natural in S . Explicitly, given $f : S \rightarrow S'$ a map of affines, we need to check that the following square commutes

$$\begin{array}{ccc} \mathcal{X}_{\mathcal{C}}(S') & \longrightarrow & \mathcal{X}_{\mathcal{C}}(S) \\ \downarrow & & \downarrow \\ \text{QCoh}(S') & \longrightarrow & \text{QCoh}(S) \end{array}.$$

This is equivalent to checking base change for the square

$$\begin{array}{ccc} X \times S & \xrightarrow{\text{id}_X \times f} & X \times S' \\ \downarrow p & & \downarrow p' \\ S & \xrightarrow{f} & S' \end{array}.$$

Since S' and $X \times S'$ are perfect, so is the map p' . Hence, base change holds by [BFN10, Proposition 3.10]. $\square$

Remark 3.5. If X is a qcqs scheme, then, as in [Toe06, §8.3], we can present

$$\mathcal{X}_{\mathrm{QCoh}(X)}(S) \cong \widetilde{\mathrm{QCoh}(X \times S)} \cong \mathrm{Map}(\mathrm{QCoh}(X), \mathrm{QCoh}(S)),$$

and a sheaf $\mathcal{E} \in \mathrm{QCoh}(X \times S)$ corresponds to the map

$$\varphi_{\mathcal{E}} : \mathcal{F} \mapsto p_{S*}(p_X^* \mathcal{F} \otimes \mathcal{E}).$$

In particular, by applying $\varphi_{\mathcal{E}}$ to the structure sheaf $\mathcal{O}_X$, we recover the map defined in the lemma above. In fact, by [BFN10, Theorem 4.14], this is the case for any perfect stack X .

We henceforth only consider categories $\mathcal{C}$ of the type considered in Lemma 3.3. For a category $\mathcal{C}$ of the form $\mathrm{QCoh}(X)$ with X a perfect stack or $A\text{-mod}$ with A an associative algebra, we define its **n -dimensional representation moduli** as

$$\mathcal{R}ep_n(\mathcal{C}) := \mathcal{X}_{\mathcal{C}} \times_{\mathrm{Vect}} \{k^n\},$$

where the map $\mathcal{X}_{\mathcal{C}} \rightarrow \mathrm{Vect}$ is the map of Lemma 3.3 and $\mathrm{pt} \rightarrow \mathrm{Vect}$ classifies k^n .

Remark 3.6. This definition depends not only on the abstract category $\mathcal{C}$, but also on the map to Vect defined either by Lemma 3.3 or by the presentation $\mathcal{C} = A\text{-mod}$ (with its natural restriction functor to Vect). In particular, this definition is *not* Morita invariant.

Theorem 3.7. If $\mathcal{C} = A\text{-mod}$ with A an associative algebra, then $\mathcal{R}ep_n(A) \cong \mathrm{Rep}_n(A)$ is the derived representation variety of A .

Proof. Let $QA \xrightarrow{\sim} A$ be a cofibrant replacement of A in associative algebras. Since $A\text{-mod} \cong QA\text{-mod}$, we also have

$$\widetilde{(A \otimes B)\text{-mod}} \cong \widetilde{(QA \otimes B)\text{-mod}}.$$

Moreover, this equivalence respects pushforward of modules [Hin97, Theorem 3.3.1]; thus, we are free to assume $A = QA$ is cofibrant.

In [Lur17, §4.7.1], Lurie shows that given a category $\mathcal{C}$ and an object $M \in \mathcal{C}$, there is an equivalence

$$\mathrm{Map}_{\mathrm{Alg}(\mathcal{C})}(A, \mathrm{End}(M)) \cong \mathrm{LMod}_A(\mathcal{C}) \times_{\mathcal{C}} \{M\}$$

for any $A \in \mathrm{Alg}(\mathcal{C})$, where $\mathrm{End}(M)$ is the “universal endomorphism algebra of M .” By construction, this is natural with respect to colimit preserving functors $F : \mathcal{C} \rightarrow \mathcal{D}$. Now, consider the categories

$$\begin{aligned} \mathcal{C} &= \widetilde{B\text{-mod}} \\ \mathcal{D} &= \widetilde{C\text{-mod}}, \end{aligned}$$

for B, C k -algebras, the object $M = B^n$, the algebra $A \otimes B \in \mathrm{Alg}(\mathcal{C}) = \mathrm{Alg}_{B/}$, and the functor $F = f^* : \mathcal{C} \rightarrow \mathcal{D}$ for a map of algebras $f : B \rightarrow C$. By construction, the universal endomorphism algebra is just usual endomorphisms in this case. Thus, Lurie’s equivalence becomes an identification

$$\mathrm{Hom}_{\mathrm{Alg}}(A, M_n(B)) \cong \mathrm{Map}_{\mathrm{Alg}(\mathcal{C})}(A \otimes B, \mathrm{End}(B^n)) \cong \mathcal{X}_{A\text{-mod}}(\mathrm{Spec} B) \times_{\mathrm{Vect}(\mathrm{Spec} B)} \{B^n\},$$

where the LHS of his equivalence is identified with $\text{Rep}_n(A)(\text{Spec } B)$ by the tensor-hom adjunction. It remains to check that under tensor-hom, the functor f^* on mapping spaces identifies with the map

$$M_n(f) : \text{Hom}_{\text{Alg}}(A, M_n(B)) \rightarrow \text{Hom}_{\text{Alg}}(A, M_n(C)).$$

In other words, we just need to check that tensor-hom is natural with respect to maps $f : B \rightarrow C$. This would provide an identification of the S -points of both $\text{Rep}_n(A)$ and $\mathcal{R}ep_n(\mathcal{X})$, together with naturality in the test affine S . Indeed, given

$$\varphi : A \otimes B \rightarrow M_n(B),$$

$f^*\varphi$ is sent, under tensor-hom, to $f^*\varphi(- \otimes 1_C)$. On the other hand, the canonical isomorphism $M_n(B) \otimes_B C = M_n(C)$ is obtained by sending $M \otimes c \mapsto (f(M_{ij})c)$, where $M \in M_n(B)$ is a matrix and $M_{ij} \in B$ are the entries. Thus, under tensor-hom, $f^*\varphi(M)$ is sent to the matrix $(f(M_{ij})1_C) = (f(M_{ij})) = M_n(f)(M)$. $\square$

3.2 Examples

Example 3.8. Let $\mathcal{C} = \text{QCoh}(\mathbb{P}^1)$. Then points of $\mathcal{R}ep_n(\mathcal{C})(k)$ are pairs of sheaves $\mathcal{F} \in \text{QCoh}(X)$ and isomorphisms

$$\varphi : \Gamma(\mathbb{P}^1, \mathcal{F}) \xrightarrow{\cong} k^n.$$

If $\mathcal{F}$ is coherent, then we can write

$$\mathcal{F} = \bigoplus_i \mathcal{O}(d_i) \oplus \bigoplus_j \mathcal{O}(e_j)[1] \oplus T \oplus \bigoplus_k \mathcal{O}(-1)[m_k],$$

where $d_i \geq 0$, $e_j \leq -2$, T is a torsion sheaf of length ℓ , $\sum(d_i + 1) + \sum(-e_j - 1) + \ell = n$, and the m_k are arbitrary. For families, points of $\mathcal{R}ep_n(\mathcal{C})(S)$ are pairs of a sheaf $\mathcal{F} \in \text{QCoh}(\mathbb{P}_S^1)$ and an isomorphism

$$\varphi : p_{S*}\mathcal{F} \xrightarrow{\cong} \mathcal{O}_S^n,$$

where $p_S : \mathbb{P}_S^1 \rightarrow S$ is the projection. Over geometric points p of S and $\mathcal{F}$ coherent, $\mathcal{F}|_{\mathbb{P}_{S,p}^1}$ is as before, except the sheaves $\mathcal{O}(d_i)$, $\mathcal{O}(e_j)$, $\mathcal{O}(-1)$ are relative to S and T is finite flat of rank ℓ over S . Note that due to the appearance of arbitrarily many summands of $\mathcal{O}(-1)$ in $\mathcal{F}$, there is no reason to expect $\mathcal{R}ep_n(\mathcal{C})$ to be quasi-compact.

Example 3.9. In this example, we will assume that k is algebraically closed. Let $\mathcal{C} = \text{QCoh}(BT)$ for T a torus. Then points of $\mathcal{R}ep_n(\mathcal{C})(k)$ are arbitrary T -representations V together with isomorphisms $\varphi : V^T \xrightarrow{\cong} k^n$, where V^T denotes T -invariants of the T -representation V . Recall that $\text{QCoh}(BT)$ decomposes as

$$\text{QCoh}(BT) = \text{Rep}(BT) \cong \bigoplus_{\lambda \in X^*(T)} \text{Vect}_\lambda,$$

where $X^*(T)$ is the character lattice and Vect_λ denotes representations V where T acts by the character λ . In particular, given a representation V ,

$$V = \bigoplus_{\lambda \in X^*(T)} V_\lambda,$$

and $V^T = V_0$ is the 0-weight space. Thus, k -points are given by T -representations V with the $\lambda = 0$ weight having multiplicity n together with a point of $\text{GL}_n(k)$ which identifies $V_0 \xrightarrow{\cong} k^n$.

When we pass to S -points, we now have families of T -representations $\mathcal{F} \in \text{QCoh}(BT \times S)$ with a trivialization $p_{S*}\mathcal{F} \xrightarrow{\cong} \mathcal{O}_S^n$, where $p_S : BT \times S \rightarrow S$, i.e. an element of $\text{GL}_n(S)$. As in the case $S = \text{pt}$, we have a decomposition

$$\text{QCoh}(BT \times S) \cong \text{QCoh}(BT) \otimes \text{QCoh}(S) \cong \bigoplus_{\lambda \in X^*(T)} \text{QCoh}(S)_\lambda,$$

where $\text{QCoh}(S)_\lambda := \text{Vect}_\lambda \otimes \text{QCoh}(S)$ denotes sheaves on S whose underlying vector space is a T -representation with weight λ . Thus, we see that an S -point is given by a sheaf $\mathcal{F} \in \text{QCoh}(S)$ with a decomposition

$$\mathcal{F} \cong \bigoplus_{\lambda \in X^*(T)} \mathcal{F}_\lambda,$$

together with an isomorphism $\mathcal{F}_0 \xrightarrow{\cong} \mathcal{O}_S^n$, which is an element of $\text{GL}_n(S)$. If we restrict to coherent, rather than quasi-coherent, sheaves (cf. Section 4.1), then we see that the classical truncation of the coherent moduli is given by

$$\coprod_{m: X^*(T) \setminus \{0\} \rightarrow \mathbb{N}} \prod_{\lambda} B \text{GL}_{m_\lambda},$$

where $m_\lambda = \text{rk}(\mathcal{F}_\lambda)$ and m has finite support.

Example 3.10. As in the previous example, we still assume that k is algebraically closed. Let $\mathcal{C} = \text{QCoh}(BG)$ for G a reductive group. If $V \in \mathcal{R}ep_n(\text{QCoh}(BG))(k)$, then $(V, \varphi) \in \text{QCoh}(BG) = \text{Rep}(G)$ is a pair of a G -representation V equipped with an isomorphism $\varphi : V^G := \text{Hom}_G(k, V) \xrightarrow{\cong} k^n$. There is a decomposition

$$\text{Rep}(G) \cong \prod_{\rho} \text{Vect},$$

where the product is over irreducible representations ρ of G , because k is algebraically closed of characteristic 0 and G is reductive. Thus, the representation V decomposes as

$$V \cong k^n \otimes \rho_{\text{triv}} \oplus \bigoplus_{\rho \neq \rho_{\text{triv}}} M_\rho \otimes \rho,$$

where the first summand is the trivial isotype and the other summands are irreducible representations ρ with multiplicities M_ρ .

Passing to S -points, as in the case of BT , we have families of G -representations $\mathcal{F} \in \mathrm{QCoh}(BG \times S) \cong \mathrm{QCoh}(BG) \otimes \mathrm{QCoh}(S)$ together with isomorphisms $\varphi : p_{S*} \mathcal{F} \xrightarrow{\cong} \mathcal{O}_S^n$. As in the case of k -points, we have an isotypic decomposition

$$\mathcal{F} \cong \bigoplus_{\rho} \mathcal{M}_{\rho} \otimes \rho,$$

where the $\mathcal{M}_{\rho} \in \mathrm{QCoh}(S)$ and the isomorphism φ induces $\mathcal{M}_{\rho_{\mathrm{triv}}} \xrightarrow{\cong} \mathcal{O}_S^n$. If we restrict to coherent sheaves— G -equivariant vector bundles on S , then the classical truncation of the moduli has a presentation as

$$\bigsqcup_m \prod_{\rho \neq \rho_{\mathrm{triv}}} B\mathrm{GL}_{m_{\rho}},$$

where the coproduct is over all assignments with finite support of multiplicities to irreducible representations ρ .

4 Trace Maps II

In Theorem 2.8, we have constructed an S^1 -equivariant trace map

$$\mathrm{HH}_*(A) \rightarrow \mathbb{L}(A)_n,$$

which corresponds by the universal property of S^1 -coinvariants to a trace map

$$\mathrm{HC}_*(A) \rightarrow \mathbb{L}(A)_n,$$

arising from a domain wall between the field theories defined by A -mod and $\mathbb{L}(A)_n$ -mod. We claim that we have such a domain wall for representation moduli defined by the “universal representation.” More precisely, given a quasi-projective classical scheme X , we will show that there is a Fourier-Mukai integral transform from $\mathrm{QCoh}(X)$ to

$$\mathrm{QCoh}(\mathrm{Quot}_{\mathcal{O}_X^n}^{n, \mathrm{fr}}(X)),$$

where $\mathrm{Quot}_{\mathcal{O}_X^n}^{n, \mathrm{fr}}(X)$ is a derived enhancement of the framed locus of the Quot scheme of points on X relative to $\mathcal{O}_X^n$. This scheme will be constructed as a substack of “coherent representations.”

$$\mathcal{R}ep_n^{\mathrm{coh}}(X) := \mathrm{Coh}_{\mathrm{prop}}(X) \times_{\mathrm{Vect}} \mathrm{pt} \subset \mathcal{R}ep_n(X),$$

which is the substack of $\mathcal{R}ep_n(\mathrm{QCoh}(X))$ of coherent sheaves with proper support (as in [PS22]). This integral transform will formally induce a map on Hochschild homologies, which, composed with the constant loop map

$$\mathrm{Quot}_{\mathcal{O}_X^n}^{n, \mathrm{fr}}(X) \rightarrow \mathcal{L} \mathrm{Quot}_{\mathcal{O}_X^n}^{n, \mathrm{fr}}(X),$$

will give a generalization of the trace map of [BKR13] (see Theorem 4.17).

4.1 Coherent Representation Moduli and Derived Quot Schemes

We now recall the definition of derived coherent sheaves relative to a morphism of stacks. The following definitions can be found (with homological, rather than cohomological conventions) at the beginning of [PS22, §2]; we reproduce them here for the reader's convenience. Recall that for $A \in \mathbf{CAlg}$, an A -module M is said to be **almost perfect** if it is eventually connective and all truncations $\tau^{\geq i} M$ are compact in $A\text{-mod}^{\geq i}$. We denote the full subcategory of $A\text{-mod}$ of almost perfect modules by $\mathbf{APerf}(\mathrm{Spec} A)$. For any prestack X , we can define its **category of almost perfect complexes**

$$\mathbf{APerf}(X) := \lim_{S \rightarrow X} \mathbf{APerf}(S).$$

Given a morphism $f : X \rightarrow S$ in $\mathbf{Stk}$, a quasi-coherent sheaf $\mathcal{F} \in \mathbf{QCoh}(X)$ has **tor amplitude $\geq -n$ relative to S** if for every $\mathcal{G} \in \mathbf{QCoh}(S)^\vee$,

$$H^{-i}(\mathcal{F} \otimes f^* \mathcal{G}) = 0$$

for $i \notin [0, n]$. The full subcategory of $\mathbf{APerf}(X)$ spanned by sheaves of tor amplitude $\geq -n$ relative to S is denoted $\mathbf{APerf}_S^{\geq -n}(X)$. Define

$$\mathbf{Coh}_S(X) := \mathbf{APerf}_S^{\geq 0}(X)$$

as the category of **flat families of coherent sheaves on X relative to S** . By Lemma 2.4 of *op. cit.*, the assignment

$$\begin{array}{c} \mathbf{Aff}^{\mathrm{op}} \rightarrow \mathbf{Spc} \\ S \mapsto \overline{\mathbf{Coh}_S(X \times S)} \end{array}$$

defines a stack $\mathbf{Coh}(X)$ called the **derived stack of coherent sheaves on X** . Now, if $X \rightarrow S$ is a map of schemes locally almost of finite type (laft) and $\mathcal{F} \in \mathbf{APerf}(X)$, we say that $\mathcal{F}$ has **proper support relative to S** if there is a closed subscheme $Z \hookrightarrow X$ such that $Z \rightarrow S$ is proper and $\mathcal{F}|_{X \setminus Z} \cong 0$. Denote the category of almost perfect complexes with proper support by $\mathbf{APerf}_{\mathrm{prop}}(X)$. For a quasi-projective classical scheme X , define the **stack of coherent sheaves with proper support** as

$$\mathbf{Coh}_{\mathrm{prop}}(X) := \mathbf{APerf}_{\mathrm{prop}}(X) \times_{\mathbf{APerf}(X)} \mathbf{Coh}(X).$$

Remark 4.1. In [PS22], this stack is defined as

$$\mathbf{Perf}_{\mathrm{prop}}(X) \times_{\mathbf{Perf}(X)} \mathbf{Coh}(X),$$

but the map $\mathbf{Coh}(X) \rightarrow \mathbf{Perf}(X)$ does not exist unless X is smooth (see [PS22, Lemma 2.18 and Corollary 2.19]). The first author of *op. cit.* has informed me in private correspondence that this fiber product was intended to be as in our definition above.

Example 4.2. Let $X = \mathrm{Spec} k[x]/(x^2)$, $k \in \mathbf{QCoh}(X)$, and $\mathrm{pt} = \mathrm{Spec} k$. Then k is almost perfect, as it has bounded, coherent cohomologies. It is also of tor amplitude 0 relative to pt , since if $V \in \mathbf{Vect}^\vee$, then

$$k \otimes_{k[x]/(x^2)} (k[x]/(x^2) \otimes V) = V$$

is tor amplitude 0. But it is not perfect, since it has infinite projective dimension as a $k[x]/(x^2)$ -module. Thus, it is not an object of

$$\mathrm{Perf}_{\mathrm{prop}}(X)(\mathrm{pt}) \times_{\mathrm{APerf}(X)(\mathrm{pt})} \mathrm{Coh}(X)(\mathrm{pt}).$$

Since our eventual goal is a definition of the representation moduli and, for an algebra A , $k \in \mathrm{Rep}_1(A)(k)$, we see that the definition of the stack of coherent sheaves with proper support given above is more suitable for a definition of representation varieties.

Proposition 4.3. Let X be a classical quasi-projective scheme. Then the stack

$$\mathrm{Coh}_{\mathrm{prop}}(X)$$

is a 1-Artin stack locally almost of finite type with affine diagonal.

Proof. The proof of Proposition 2.29 of [PS22] applies verbatim, as it does not use the fact that the moduli parametrizes perfect, rather than almost perfect, complexes. Similarly, the morphism $\mathrm{Coh}(X) \rightarrow \mathrm{APerf}(X)$ is an infinitesimally cohesive and nilcomplete morphism of stacks which are themselves infinitesimally cohesive and nilcomplete. Since these properties are stable under base change (cf. the proof of Corollary 2.14 of *op. cit.*), it follows that $\mathrm{Coh}_{\mathrm{prop}}(X) \rightarrow \mathrm{APerf}_{\mathrm{prop}}(X)$ is infinitesimally cohesive and nilcomplete. It remains to check that $\mathrm{APerf}_{\mathrm{prop}}(X) \rightarrow \mathrm{APerf}(X)$ is infinitesimally cohesive and nilcomplete. Once we have this, we can use [Lur18, Theorem 18.1.0.2] to conclude that $\mathrm{Coh}_{\mathrm{prop}}(X)$ is a 1-Artin stack locally almost of finite type.

We now need the following result, which says that proper support is detected on the classical truncation (in fact, on the underlying reduced, classical scheme).

Lemma 4.4. If S is an affine scheme and $i : X \times S_{\mathrm{red}} \rightarrow X \times S$ is the inclusion, then $\mathcal{F} \in \mathrm{APerf}(X \times S)$ has proper support relative to S if and only if $i^* \mathcal{F}$ has proper support relative to S_{red} .

Proof of Lemma. The forward implication is immediate by base change, while the reverse implication can be seen as follows. Suppose $i^* \mathcal{F}$ is supported on a closed subscheme $Z \hookrightarrow X \times S_{\mathrm{red}}$ such that $Z \rightarrow S_{\mathrm{red}}$ is proper. Then the composition $Z \rightarrow S_{\mathrm{red}} \rightarrow S$ is also proper, so we see that Z is a closed subscheme of $X \times S$ proper over S . Denoting $U := (X \times S) \setminus Z$, we see that $i^*(\mathcal{F}|_U) \cong 0$. It is now sufficient to recall that pullbacks to geometric fibers are jointly conservative on almost perfect complexes to conclude. $\square$

Now, to show that $\mathrm{APerf}_{\mathrm{prop}}(X) \rightarrow \mathrm{APerf}(X)$ is nilcomplete, we would like to show that given an affine S and $S_n = \tau_{\geq n} S$ its truncations,

$$\mathrm{APerf}_{\mathrm{prop}}(X)(S) \rightarrow \mathrm{APerf}(X)(S) \times_{\lim \mathrm{APerf}(X)(S_n)} \lim \mathrm{APerf}_{\mathrm{prop}}(X)(S_n)$$

is an equivalence. If $\mathcal{F} \in \mathrm{APerf}(X \times S)$ and every $\mathcal{F}|_{X \times S_n}$ has proper support, then its restriction to S_{red} has proper support. Thus, $\mathcal{F}$ has proper support by the above Lemma. This proves essential surjectivity, while fully faithfulness follows from the fact that the map $\mathrm{APerf}_{\mathrm{prop}}(X) \rightarrow \mathrm{APerf}(X)$ is an inclusion.

To show infinitesimal cohesiveness, we follow the proof of [PS22, Lemma 2.13]. Let $S = \mathrm{Spec} A$ be an affine scheme, $M \in \mathrm{QCoh}(S)^{\geq 1}$, $S[M] := \mathrm{Spec}(A \oplus M)$, and $d : S[M] \rightarrow S$ be a derivation, and consider the pushout diagram

$$\begin{array}{ccc} S[M] & \xrightarrow{d} & S \\ \downarrow d_0 & & \downarrow f_0 \\ S & \xrightarrow{f} & S_d[M[-1]] \end{array},$$

where d_0 is the 0 derivation. Notice that all of the schemes in the diagram above have the same classical reduction S_{red} . Let $\mathcal{F} \in \mathbf{APerf}(X \times S_d[M[-1]])$, and suppose the restrictions of $\mathcal{F}$ along f , f_0 and $fd \cong f_0 d_0$ have proper support. Since the subsequent restrictions to S_{red} give the same almost perfect complex with proper support, we see that $\mathcal{F}$ has proper support over $S_d[M[-1]]$ by Lemma 4.4. This proves essential surjectivity of the relevant map; fully faithfulness follows because $\mathbf{APerf}_{\text{prop}}(X) \rightarrow \mathbf{APerf}(X)$ is an inclusion.

Finally, the stack $\text{Coh}_{\text{prop}}(X)$ has affine diagonal by [GR17, Chapter 2 Corollary 3.2.8], as its classical truncation is the stack of coherent complexes with proper support, which has affine diagonal by [Aut, Tag 0DLY]. $\square$

Theorem 4.5. Let $\mathcal{C} = \text{QCoh}(X)$ for a classical quasi-projective scheme X . Then $\mathcal{R}ep_n(\mathcal{C})$ has a substack

$$\mathcal{R}ep_n^{\text{coh}}(X) := \text{Coh}_{\text{prop}}(X) \times_{\text{Vect pt}} \text{pt} \subset \mathcal{R}ep_n(\mathcal{C}),$$

called the **coherent representation moduli**, which is a 1-Artin stack locally almost of finite type (laft). Moreover, this stack has affine diagonal.

Proof. We will show that $\mathcal{R}ep_n^{\text{coh}}(X)$ is an open substack of a stack $\mathcal{V}$ which is affine and almost of finite type over $\text{Coh}_{\text{prop}}(X)$. This proves that $\mathcal{R}ep_n^{\text{coh}}(X)$ is quasi-affine and almost of finite type over $\text{Coh}_{\text{prop}}(X)$. Since $\text{Coh}_{\text{prop}}(X)$ is 1-Artin and laft by Proposition 4.3, this proves that $\mathcal{R}ep_n^{\text{coh}}(X)$ is also 1-Artin and laft.

We first show that if $\mathcal{E} \in \text{Coh}_{\text{prop}}(X)(S)$ is an S -point, then $\mathcal{V}_{\mathcal{E}} = p_{S*}\mathcal{E}$, where $p_S : X \times S \rightarrow S$, is perfect and its construction commutes with arbitrary base change. Indeed, to see that we have base change, it is sufficient to observe that p_S is schematic quasi-compact, hence satisfies the conditions of [GR17, Chapter 3 Section 2.2]. By [PS22, Proposition 2.28], $\mathcal{V}_{\mathcal{E}}$ is almost perfect. Let $M \in \text{QCoh}(S)^{\heartsuit}$. Then $\mathcal{V}_{\mathcal{E}} \otimes_{\mathcal{O}_S} M \cong p_{S*}(\mathcal{E} \otimes_{\mathcal{O}_X} p_{S*}M)$ by the projection formula ([GR17, Chapter 3 Lemma 3.2.4]). By definition of $\mathcal{E}$, $\mathcal{E} \otimes_{\mathcal{O}_X} p_{S*}M$ is tor amplitude 0. Since X is quasi-compact, there is a finite Zariski cover; thus, for $\mathcal{F} \in \text{QCoh}(X \times S)^{\heartsuit}$, $p_*\mathcal{F}$ is computed by a finite Čech complex where the terms are t -exact pushforwards of restrictions of $\mathcal{F}$. Thus, $p_*\mathcal{F}$ is of finite tor amplitude; moreover, it is coconnective. It follows that $\mathcal{V}_{\mathcal{E}}$ is perfect coconnective by [Lur17, Proposition 7.2.4.23].

Because of the base change compatibility of the definition of $\mathcal{V}_{\mathcal{E}}$, the assignments $\mathcal{E} \mapsto p_*\mathcal{E}$ glue to give a perfect complex $\mathcal{V} \in \text{Perf}(\text{Coh}_{\text{prop}}(X))$. Then $(\mathcal{V}^n)^{\vee}$ is a connective perfect complex; hence, defines a stack

$$\mathbb{V} := \text{Spec}_{\text{Coh}_{\text{prop}}(X)} \text{Sym}((\mathcal{V}^n)^{\vee}) \xrightarrow{p} \text{Coh}_{\text{prop}}(X)$$

which is affine almost of finite type over $\text{Coh}_{\text{prop}}(X)$. Notice that S -points of $\mathbb{V}$ are precisely given by S -points $f : S \rightarrow \text{Coh}_{\text{prop}}(X)$ together with a map $\mathcal{O}_S^n \rightarrow f^*\mathcal{V} = p_{S*}\mathcal{E}$ by the base change compatibility of $\mathcal{V}$. We thus see that S -points of $\mathbb{V}$ are pairs of an almost perfect complex $\mathcal{E}$ on $X \times S$ and a map $\mathcal{O}_S^n \rightarrow p_{S*}\mathcal{E}$.

It remains to show that $\mathcal{R}ep_n^{\text{coh}}(X)$ is an open subfunctor of this stack. Let $\mathcal{O}_{\mathbb{V}}^n \rightarrow p^*\mathcal{V}$ be the universal map, and observe that its cofiber is a perfect complex. Clearly, the locus where that cofiber vanishes is open; moreover, that locus is precisely the locus where the map

$$\mathcal{O}_{\mathbb{V}}^n \rightarrow p^*\mathcal{V}$$

is an isomorphism.

Now, we will show that $\mathcal{R}ep_n^{\text{coh}}(X)$ has affine diagonal. Note that $\text{Coh}_{\text{prop}}(X)$ has affine diagonal by Proposition 4.3. Since $\mathbb{V} \rightarrow \text{Coh}_{\text{prop}}(X)$ is affine, it is separated, so the diagonal

$$\Delta_{\mathbb{V}} : \mathbb{V} \rightarrow \mathbb{V} \times_{\text{Coh}_{\text{prop}}(X)} \mathbb{V}$$

is also affine. We see that the diagonal of $\mathcal{R}ep_n^{\text{coh}}(X)$ factors as

$$\Delta_{\mathcal{R}ep_n^{\text{coh}}(X)} : \mathcal{R}ep_n^{\text{coh}}(X) \rightarrow \mathcal{R}ep_n^{\text{coh}}(X) \times_{\text{Coh}_{\text{prop}}(X)} \mathcal{R}ep_n^{\text{coh}}(X) \rightarrow \mathcal{R}ep_n^{\text{coh}}(X) \times \mathcal{R}ep_n^{\text{coh}}(X),$$

where the first map is affine because the map $\mathcal{R}ep_n^{\text{coh}}(X) \rightarrow \text{Coh}_{\text{prop}}(X)$ is quasi-affine, hence has affine relative diagonal (it is a base change of $\Delta_{\mathbb{V}}$), and the second is affine because it is the base change of the affine diagonal of $\text{Coh}_{\text{prop}}(X)$. $\square$

For a classical quasi-projective scheme X , we have the substack

$$\text{Coh}_{\text{prop}}^{\leq d}(X) \subset \text{Coh}_{\text{prop}}(X)$$

of sheaves whose proper support is dimension $\leq d$. The stack $\text{Coh}_{\text{prop}}^{\leq 0}(X)$ has a decomposition

$$\text{Coh}_{\text{prop}}^{\leq 0}(X) = \coprod_{m \in \mathbb{N}_0} \text{Coh}_0^m(X)$$

by substacks of sheaves with Hilbert polynomial identically equal to m , which follows from the decomposition of the underlying classical stack. In particular, we have a stack

$$\text{Coh}_0^n(X)$$

of length n sheaves on X .

Corollary 4.6. Let X be a quasi-projective classical scheme. Then the fiber

$$\text{Quot}_{\mathcal{O}_X^n}^{n, \text{fr}}(X) := \text{Coh}_0^n(X) \times_{\text{Vect}} \text{pt} \subset \mathcal{R}ep_n(X)$$

is a quasi-compact scheme almost of finite type with affine diagonal.

Proof. Since

$$Y = \text{Quot}_{\mathcal{O}_X^n}^{n, \text{fr}}(X) \subset \mathcal{R}ep_n^{\text{coh}}(X)$$

has classical truncation a scheme of finite type by the proof of [KV22, Proposition 4.4.1], it is a quasi-compact scheme almost of finite type. This immediately implies that Y is Deligne-Mumford since its classical truncation is [Kha26, Definition 8.5.7(iii)]. One can then use [Lur18, Corollary 1.6.7.4] to conclude that Y is in fact a scheme.

Further, the scheme Y has affine diagonal because it is a substack of $\mathcal{R}ep_n^{\text{coh}}(X)$, which has affine diagonal by Theorem 4.5. $\square$

Remark 4.7. An S -point of $\mathrm{Quot}_{\mathcal{O}_X^n}^{n,\mathrm{fr}}(X)$ is thus a family $\mathcal{E} \in \mathrm{Coh}_0^n(X)(S)$ together with an isomorphism

$$\mathcal{O}_S^n \xrightarrow{\cong} p_{S*}\mathcal{E}.$$

By adjunction, this isomorphism determines a quotient map $q : \mathcal{O}_{X \times S}^n \rightarrow \mathcal{E}$ which is surjective on H^0 . Hence, the classical truncation of this derived scheme is the open locus of the classical Quot scheme of length n quotients whose adjoint $\mathcal{O}_S^n \rightarrow p_{S*}\mathcal{E}$ is an isomorphism. We call this the framed locus.

Even for affine $X = \mathrm{Spec} A$, we don't necessarily have that coherent representations match all representations. Indeed, suppose A is the free commutative algebra on infinitely many generators. Then k is not finitely presented as an A -module. By [Lur17, Proposition 7.2.4.17], this implies that it is not almost perfect as an A -module, hence not coherent in the above sense. However, we do have the following.

Proposition 4.8. Let A be a classical finitely presented commutative algebra. Then

$$\mathrm{Rep}_n(A) \cong \mathrm{Quot}_{A^n}^{n,\mathrm{fr}}(\mathrm{Spec} A).$$

Proof. Note that by Theorem 3.7, we have an inclusion

$$\mathrm{Quot}_{A^n}^{n,\mathrm{fr}}(\mathrm{Spec} A) \hookrightarrow \mathrm{Rep}_n(A),$$

and it is sufficient to show that this map is surjective. Let $\mathrm{Spec} B$ be an affine scheme. Any almost perfect $(A \otimes B)$ -module M such that $M \cong B^n$ as B -modules has support which is finite over $\mathrm{Spec} B$; hence, the coherent sheaf on $\mathrm{Spec}(A \otimes B)$ which it defines has proper support relative to $\mathrm{Spec} B$. Since the pushforward of the sheaf corresponding to M to $\mathrm{Spec} B$ is locally free of rank n and its support is finite over $\mathrm{Spec} B$, it is a flat family of length n sheaves, hence, it defines a point of $\mathrm{Coh}_0^n(\mathrm{Spec} A)$. Clearly, M is tor amplitude 0 over B , as, for $N \in B\text{-mod}^\heartsuit$,

$$M \otimes_{A \otimes B} (A \otimes B \otimes_B N) \cong M \otimes_B N \cong B^n \otimes_B N \cong N^n.$$

Thus, it is sufficient to show that any $(A \otimes B)$ -module M with an isomorphism $M \cong B^n$ is almost perfect.

We will first prove the following.

Lemma 4.9. Let $R \rightarrow S$ be a map of connective algebras such that S is almost perfect as an $(S \otimes_R S)$ -module. If M is an S -module which is almost perfect over R , then it is almost perfect over S .

Proof of Lemma. Without loss of generality, we can assume that M is connective; indeed, since M is almost perfect over R , it is bounded above. Let $T = S \otimes_R S$, and consider the functor

$$- \otimes_S M : T\text{-mod} \rightarrow S\text{-mod},$$

where $N \otimes_S M$ has an S -module structure via the left S -action on the first factor. Since S is almost perfect over T , we can, by [Lur17, Proposition 7.4.2.11], choose for each m a perfect T -module P_m equipped with a map $P_m \rightarrow S$ that has m -connective fiber. Applying $- \otimes_S M$, the map $P_m \otimes_S M \rightarrow M$ has m -connective fiber. Since $P_m \in \mathrm{Perf}(T)$, $P_m \otimes_S M$ is in the thick subcategory of $S\text{-mod}$ generated by $S \otimes_R M$. Since $S \otimes_R M$ is almost perfect over S by base change of the almost perfect R -module M and almost perfect modules are closed under finite colimits and retracts by *loc. cit.* $P_m \otimes_S M$ is almost perfect over S . Now, if M is bounded above and for every m there is an almost perfect $Q_m \rightarrow M$ whose fiber is m -connective, then M is almost perfect by *loc. cit.* $\square$

We would like to apply the Lemma to our situation, where $R = B$, $S = A \otimes B$, which will conclude the proof that M is almost perfect over $A \otimes B$. Note that R and S are connective because $\operatorname{Spec} B$ and $\operatorname{Spec}(A \otimes B)$ are affine schemes, and M is almost perfect over B . Since A is finitely presented over k , it is almost perfect as a module over $A \otimes A$. Indeed, $A \otimes A$ is left coherent (because it's noetherian); hence, by [Lur17, Proposition 7.2.4.17], A is almost perfect over $A \otimes A$. Indeed, it is connective, and the multiplication

$$A \otimes A \rightarrow A$$

has finitely generated image. Thus, base change along a connective algebra B gives

$$A \otimes B \cong A \otimes_{A \otimes A} (A \otimes A \otimes B),$$

so $A \otimes B$ is almost perfect over $A \otimes A \otimes B = S \otimes_R S$. □

Corollary 4.10. If $X = \operatorname{Spec} A$ is affine of finite presentation, then

$$\mathcal{R}ep_n^{\operatorname{coh}}(X) \cong \operatorname{Rep}_n(A).$$

Remark 4.11. We have thus shown that in the case of $X = \operatorname{Spec} A$ of finite presentation, coherent moduli, representation moduli, derived representation varieties, and the derived framed locus $\operatorname{Quot}_{\mathcal{O}_X^n}^{n, \operatorname{fr}}(X)$ all agree. This completes the proof of Theorem 1.2 from the introduction.

4.2 The Universal Representation

From now on, let X be a classical quasi-projective scheme. Associated to the identity map

$$\operatorname{Coh}_0^n(X) \rightarrow \operatorname{Coh}_0^n(X),$$

there is a universal sheaf $\mathcal{U}$ on

$$\operatorname{Coh}_0^n(X) \times X$$

defined as follows. We can extend the moduli functor $\operatorname{Coh}_0^n(X)$ from affine schemes to stacks by right Kan extension along the inclusion $\operatorname{Aff} \rightarrow \operatorname{Stk}$. Explicitly, if Z is a stack, then we define

$$\operatorname{Coh}_0^n(X)(Z) := \lim_{S \rightarrow Z} \operatorname{Coh}_0^n(X)(S),$$

where the limit is taken over all affine schemes S mapping to Z . Thus, there are equivalences of spaces

$$\operatorname{Map}(\operatorname{Coh}_0^n(X), \operatorname{Coh}_0^n(X)) \cong \lim_{S \rightarrow \operatorname{Coh}_0^n(X)} \operatorname{Coh}_0^n(X)(S).$$

In particular, we see that there is an object in

$$\lim_{S \rightarrow \operatorname{Coh}_0^n(X)} \operatorname{Coh}_0^n(X)(S)$$

which corresponds to the identify morphism $\mathrm{Coh}_0^n(X) \rightarrow \mathrm{Coh}_0^n(X)$. Since $\mathrm{Coh}_0^n(X)(S) \subset \mathrm{Coh}_{\mathrm{prop}}(X)(S) \subset \mathrm{Coh}(X)(S) \subset \mathrm{APerf}(X \times S)$, descent for almost perfect complexes implies that

$$\mathrm{APerf}(\mathrm{Coh}_0^n(X) \times X) \cong \lim_{S \rightarrow \mathrm{Coh}_0^n(X)} \mathrm{APerf}(S \times X) \supset \lim_{S \rightarrow \mathrm{Coh}_0^n(X)} \mathrm{Coh}_0^n(X)(S).$$

This defines an almost perfect complex $\mathcal{U} \in \mathrm{APerf}(\mathrm{Coh}_0^n(X) \times X)$. Moreover, if $f : S \rightarrow \mathrm{Coh}_0^n(X)$ is an S -point, then the pullback of $\mathcal{U}$ along $f \times \mathrm{id}_X : S \times X \rightarrow \mathrm{Coh}_0^n(X) \times X$ gives a length n torsion sheaf on $S \times X$ with proper support relative to S .

Informally, this sheaf can be understood as follows. Given an S -point

$$(f, x) : S \rightarrow \mathrm{Coh}_0^n(X) \times X$$

we obtain a sheaf on $S \times X$ via the map f together with a map $x : S \rightarrow X$. By pulling back this sheaf along the graph of x in $S \times X$, we obtain a sheaf on S , and this is precisely the pullback of the sheaf $\mathcal{U}$ to S along (f, x) .

Denote $Y = \mathrm{Quot}_{\mathcal{O}_X^n}^{n, \mathrm{fr}}(X)$. Pulling back $\mathcal{U}$ along

$$\pi \times \mathrm{id}_X : Y \times X \rightarrow \mathrm{Coh}_0^n(X) \times X,$$

where $\pi : Y \rightarrow \mathrm{Coh}_0^n(X)$ is the projection, gives a universal sheaf on $Y \times X$,

$$\mathcal{R} := (\pi \times \mathrm{id}_X)^* \mathcal{U}.$$

Note that $\mathcal{R}$ is an almost perfect complex.

Lemma 4.12. The sheaf $\mathcal{R}$ has proper support relative to $Y = \mathrm{Quot}_{\mathcal{O}_X^n}^{n, \mathrm{fr}}(X)$.

Proof. Let $Z \subset (Y \times X)^{\mathrm{cl}} \subset Y \times X$ be the subscheme of $Y \times X$ defined by equipping the classical support of $\mathcal{R}|_{(Y \times X)^{\mathrm{cl}}}$ with the reduced induced subscheme structure. Note that Z is closed because $\mathcal{R}$ is almost perfect and flat relative to Y ; hence, its restriction to the classical truncation is a coherent sheaf in degree 0 and therefore has closed support. Since an almost perfect complex is 0 if it is 0 on the classical truncation,¹ we see that

$$\mathcal{R}|_{(Y \times X) \setminus Z} \cong 0.$$

It remains to check that $Z \rightarrow Y$ is proper, which is equivalent to checking that $Z \rightarrow Y^{\mathrm{cl}}$ is proper. Since properness is affine-local on the target, it is sufficient to check that for S an affine open of Y^{cl} , $\mathcal{R}|_{S \times X}$ has proper support relative to S . We observe that $Z_S = S \times_{Y^{\mathrm{cl}}} Z$ is the classical support of $\mathcal{R}|_{S \times X}$ with its reduced induced subscheme structure (by [Aut, Tag 056J] applied to the cohomology sheaves of $\mathcal{R}|_{S \times X}$). Since $\mathcal{R}|_{S \times X}$ has proper support relative to S via the composition $S \rightarrow Y \rightarrow \mathrm{Coh}_0^n(X) \subset \mathrm{Coh}_{\mathrm{prop}}(X)$, there is a closed subscheme $Z' \subset S \times X$ proper over S with $\mathcal{R}|_{(S \times X) \setminus Z'} \cong 0$; moreover, Z_S is a closed subscheme of Z' by definition. Thus, we see that $Z \rightarrow S$ is proper, which proves the claim. $\square$

Corollary 4.13. There is a closed subscheme $Z \subset Y \times X$ such that $\mathcal{R}|_{(Y \times X) \setminus Z} \cong 0$ and the composition $Z \rightarrow Y \times X \rightarrow Y$ is finite.

¹If B is a connective commutative algebra and M is a bounded above B -module, then $H^m(H^0(B) \otimes_B M) \cong H^m(M)$, and this is nonzero if and only if $H^m(M)$ is. Thus, we see that restriction to the classical truncation is indeed conservative on bounded above complexes.

Proof. The scheme Z in the proof of Lemma 4.12 has finite fibers over Y ; hence, $Z \rightarrow Y$ is quasi-finite. Using Lemma 4.12, we see that $Z \rightarrow Y$ is proper and quasi-finite, hence finite. $\square$

Proposition 4.14. Denote $Y = \text{Quot}_{\mathcal{O}_X^n}^{\text{fr}}(X)$, and let $p_Y : Y \times X \rightarrow Y$ be the projection. We then have

$$p_{Y*}\mathcal{R} \cong \mathcal{O}_Y^n.$$

Remark 4.15. This proposition shows that $\mathcal{R}$ is the analog in the non-affine setting of the $(A, \mathbb{L}(A)_n)$ -bimodule $(\mathbb{L}(A)_n)^n$ of Lemma 2.4.

Proof. By considering the Kan-extended functor of points of Y , which takes values on stacks, we see that the map $Y \rightarrow Y$ is a tautological point of $Y(Y)$; hence, there is a universal equivalence $\mathcal{O}_Y^n \cong p_{Y*}\mathcal{R}$. Indeed, given $y : S \rightarrow Y$, the composition

$$S \xrightarrow{y} Y \xrightarrow{\pi} \text{Coh}_0^n(X)$$

thus classifies a sheaf on $S \times X$, and this sheaf corresponds to the sheaf on $S \times X$

$$\mathcal{R}|_{S \times X} = (y \times \text{id}_X)^*\mathcal{R} = ((\pi \circ y) \times \text{id}_X)^*\mathcal{U}$$

by universality of $\mathcal{U}$. Further, under pushforward along $p_S : S \times X \rightarrow S$, we have an isomorphism

$$p_{S*}\mathcal{R}|_{S \times X} \cong \mathcal{O}_S^n$$

by construction of Y . Now, consider the following cartesian square

$$\begin{array}{ccc} S \times X & \xrightarrow{y \times \text{id}_X} & Y \times X \\ \downarrow p_S & & \downarrow p_Y \\ S & \xrightarrow{y} & Y \end{array} \quad .$$

Base change along this square gives an isomorphism

$$p_{S*}(y \times \text{id}_X)^* \cong y^* p_{Y*},$$

so that

$$p_{S*}\mathcal{R}|_{S \times X} \cong (p_{Y*}\mathcal{R})|_S \cong \mathcal{O}_S^n.$$

This thus identifies

$$p_{Y*}\mathcal{R} \cong \mathcal{O}_Y^n.$$

$\square$

4.3 The Universal Representation Defines a Domain Wall

By Corollary 4.6, $Y = \text{Quot}_{\mathcal{O}_X^n}^{\text{fr}}(X)$ is a quasi-compact scheme almost of finite type with affine diagonal; hence, it is perfect in the sense of [BFN10]. Since X and Y are perfect, we have equivalences

$$\text{QCoh}(Y \times X) \cong \text{QCoh}(Y) \otimes \text{QCoh}(X) \cong \text{Map}_{\text{Pr}^L}(\text{QCoh}(X), \text{QCoh}(Y)),$$

by e.g. [GR17, Proposition 3.3.4.2]. Thus, the sheaf $\mathcal{R}$ defines a left adjoint functor

$$\begin{aligned} \Phi_{\mathcal{R}} : \text{QCoh}(X) &\rightarrow \text{QCoh}(Y) \\ \mathcal{F} &\mapsto p_{Y*}(p_X^* \mathcal{F} \otimes \mathcal{R}), \end{aligned}$$

where p_Y and p_X are the corresponding projections from $Y \times X$. This functor abstractly has a right adjoint, but this right adjoint may not necessarily preserve colimits and hence may not be a right adjoint *in the category* Pr^L .

Proposition 4.16. Let X be a quasi-projective classical scheme. Then the right adjoint of $\Phi_{\mathcal{R}}$ preserves colimits.

Proof. It is sufficient to show that $\Phi_{\mathcal{R}}$ preserves compact objects. Since X and Y are perfect, so is the map $p_Y : Y \times X \rightarrow Y$ by [BFN10, Corollary 3.23]. In particular, p_Y satisfies the projection formula by [BFN10, Proposition 3.10]. Further, the compact objects of $\text{QCoh}(Z)$ for a perfect stack Z with affine diagonal are precisely the perfect complexes $\text{Perf}(Z)$ ([BFN10, Proposition 3.9]). Thus, we need to show that given a perfect complex $\mathcal{P}$ on X , $\Phi_{\mathcal{R}}(\mathcal{P}) = p_{Y*}(p_X^* \mathcal{P} \otimes \mathcal{R})$ is also perfect. Denote $\mathcal{E} := p_X^* \mathcal{P} \otimes \mathcal{R}$. Since by Lemma 4.12 $\mathcal{R}$ has proper support relative to Y , $\mathcal{E}$ has proper support relative to Y and is almost perfect; hence, $p_{Y*} \mathcal{E}$ is almost perfect by [PS22, Proposition 2.28]. It thus suffices to check that

$$p_{Y*} \mathcal{E} \otimes \mathcal{F}$$

has universally bounded cohomology for any $\mathcal{F} \in \text{QCoh}(Y)^{\heartsuit}$. It must be bounded from above. To see it is bounded below, write

$$p_{Y*} \mathcal{E} \otimes \mathcal{F} = p_{Y*}(\mathcal{E} \otimes p_Y^* \mathcal{F})$$

by the projection formula. By assumption, $\mathcal{E} \otimes p_Y^* \mathcal{F}$ has cohomology sheaves in a finite range because $\mathcal{P}$ is perfect. Since $\mathcal{R}$ has finite support $Z \rightarrow Y \times X$ by Corollary 4.13, so does $\mathcal{E} \otimes p_Y^* \mathcal{F}$. Therefore, p_{Y*} is t -exact on $\mathcal{E} \otimes p_Y^* \mathcal{F}$. $\square$

The above proposition shows that $\Phi_{\mathcal{R}}$ is actually a 1-morphism in $(\text{Pr}^L)^{\text{cont}}$, which is the $(\infty, 2)$ -subcategory of Pr^L of dualizable objects and 1-morphisms whose adjoints are also morphisms. Thus, we are in the setting of [BN19, §3.4], so there is a well-defined map

$$\text{HH}_*(X) \rightarrow \text{HH}_*(Y)$$

induced by the 1-morphism $\Phi_{\mathcal{R}}$, where for a stack Z , $\text{HH}_*(Z) := \text{HH}_*(\text{QCoh}(Z))$. In other words, $\Phi_{\mathcal{R}}$ is a “domain wall” in the sense of §2. Constant loops are a map $Y \rightarrow \mathcal{L}Y$, hence they induce a map

$$\text{HH}_*(Y) = \mathcal{O}(\mathcal{L}Y) \rightarrow \mathcal{O}(Y).$$

By composing this with the induced map on Hochschild homologies above, we have thus obtained a map

$$\mathrm{HH}_*(X) \rightarrow \mathcal{O}(Y).$$

Moreover, this composition is equivariant with respect to the natural S^1 -actions (because each of the composed maps is by [HSS17, Section 2.3]), so it descends to a map

$$\mathrm{HC}_*(X) \rightarrow \mathcal{O}(Y).$$

The next theorem shows that in the case of $X = \mathrm{Spec} A$ of finite presentation, this is the trace map of [BKR13].

Theorem 4.17. Let $X = \mathrm{Spec} A$ be an affine scheme of finite presentation. Then $\mathcal{R} = (\mathbb{L}(A)_n)^n$ as an $(A \otimes \mathbb{L}(A)_n)$ -module. Moreover, the pair of functors it defines,

$$\Phi_{\mathcal{R}} : \mathrm{QCoh}(X) \rightleftarrows \mathrm{QCoh}(\mathrm{Quot}_{\mathcal{O}_X^n}^{n, \mathrm{fr}}(X)) : \Phi_{\mathcal{R}}^R,$$

are precisely those defined by the domain wall $(\mathbb{L}(A)_n)^n$ in §2.2.

Proof. Consider $\mathrm{Quot}_{\mathcal{O}_X^n}^{n, \mathrm{fr}}(X) \times X$, which is isomorphic to $\mathrm{Rep}_n(A) \times X$ by Theorem 3.7. By Proposition 4.14, we see immediately that as an $\mathbb{L}(A)_n$ -module, $\mathcal{R}$ is free of rank n . We would now like to show that the A -module structure is precisely the one arising from Lemma 2.4. For $S = \mathrm{Spec} B$, an S -point of $\mathrm{Rep}_n(A)$ is precisely a map from $QA \otimes B \rightarrow \mathrm{End}_B((s \times \mathrm{id}_X)^* \mathcal{R}) = M_n(B)$, where

$$(s \times \mathrm{id}_X)^* \mathcal{R} = A \otimes B \otimes_{\mathbb{L}(A)_n \otimes A} (\mathbb{L}(A)_n)^n = B^n.$$

We claim that the maps

$$QA \rightarrow M_n(B)$$

are natural in B ; hence, they give a map of $\mathbb{L}(A)_n$ -algebras

$$QA \otimes \mathbb{L}(A)_n \rightarrow M_n(\mathbb{L}(A)_n).$$

Indeed, this follows from naturality of the universal sheaf $\mathcal{R}$ with respect to pullback. The A -module structure is then induced under $A\text{-mod} = QA\text{-mod}$. We next claim that this module structure is the “universal representation” defined in the proof of Lemma 2.4. We observe that the QA -action on the restriction of $\mathcal{R}$ to $\mathrm{Spec}(B)$ -points of $\mathrm{Rep}_n(A)$ is the action of QA on $M_n(B)$ by construction, which is by definition the same action as that in Lemma 2.4.

Lastly, we identify the functors defined by $\mathcal{R}$. The functor

$$\mathrm{QCoh}(X) \rightarrow \mathrm{QCoh}(\mathrm{Rep}_n(A))$$

sends an A -module M to

$$\begin{aligned} \Phi_{\mathcal{R}}(M) &= M \otimes_A (A \otimes B) \otimes_{A \otimes B} \mathcal{R} \\ &= M \otimes_A \mathcal{R}. \end{aligned}$$

The functor in the reverse direction (which is the right adjoint) sends an $\mathbb{L}(A)_n$ -module N to

$$N \otimes_{\mathbb{L}(A)_n} \mathcal{R}^R = N \otimes_{\mathbb{L}(A)_n} \mathrm{Hom}_{\mathbb{L}(A)_n}(\mathcal{R}, \mathbb{L}(A)_n) \cong N^n.$$

These are the functors identified in §2. □

5 The Shifted Symplectic Structure

For X a smooth classical quasi-projective scheme, we will show that there is a GL_n action on $Y = \mathrm{Quot}_{\mathcal{O}_X^n}^{n,\mathrm{fr}}(X)$ such that the quotient stack Y/GL_n admits a natural shifted symplectic structure (see [Pan+11] for what this means). In the case of $X = \mathrm{Spec} A$ with $\dim X = 2$, this structure generalizes the 0-shifted symplectic structure constructed on the GL_n character variety of a genus $g \leq 1$ surface by Goldman in [Gol84].

Proposition 5.1. The derived framed locus of the Quot scheme of points has a presentation

$$\mathrm{Quot}_{\mathcal{O}_X^n}^{n,\mathrm{fr}}(X) \cong \mathrm{Coh}_0^n(X) \times_{B\mathrm{GL}_n} \mathrm{pt},$$

where $\mathrm{Coh}_0^n(X) \rightarrow B\mathrm{GL}_n$ classifies the pushforward of the universal family $\mathcal{U}$ to $\mathrm{Coh}_0^n(X)$. In particular, the projection

$$\mathrm{Quot}_{\mathcal{O}_X^n}^{n,\mathrm{fr}}(X) \rightarrow \mathrm{Coh}_0^n(X)$$

is a GL_n -torsor.

Proof. Let $p : \mathrm{Coh}_0^n(X) \times X \rightarrow \mathrm{Coh}_0^n(X)$ be the projection, and denote $\mathcal{V} := p_*\mathcal{U}$, where $\mathcal{U}$ is the universal sheaf of Section 4.2. We claim that $\mathcal{V}$ is locally free of rank n . It suffices to show that given an S -point $f : S \rightarrow \mathrm{Coh}_0^n(X)$, $f^*\mathcal{V}$ is locally free of rank n . The map p is schematic quasi-compact; indeed, it is the base change of such a map. Thus, by [GR17, Chapter 3 Proposition 2.2.2], base change holds for the diagram

$$\begin{array}{ccc} S \times X & \xrightarrow{f \times \mathrm{id}_X} & \mathrm{Coh}_0^n(X) \times X \\ \downarrow p_S & & \downarrow p \\ S & \xrightarrow{f} & \mathrm{Coh}_0^n(X) \end{array} .$$

This allows us to write

$$f^*\mathcal{V} \cong p_{S*}(f \times \mathrm{id}_X)^*\mathcal{U}.$$

By construction of the universal family, $f^*\mathcal{V}$ is precisely the pushforward along p_{S*} of the sheaf on $S \times X$ classified by f , which we denote by $\mathcal{E}$. At any geometric point s of S , $f^*\mathcal{V} \otimes k(s) \cong \Gamma(X_s, \mathcal{E}_s) \cong k(s)^n$ in degree 0, since $\mathcal{E}_s$ has length n . Thus, the bundle $\mathcal{V}$ defines a map

$$\mathrm{Coh}_0^n(X) \rightarrow B\mathrm{GL}_n .$$

Since $B\mathrm{GL}_n \hookrightarrow \mathrm{Vect}$ is a monomorphism in Stk and $\mathcal{V}$ factors through it, we immediately have an identification

$$\mathrm{Coh}_0^n(X) \times_{B\mathrm{GL}_n} \mathrm{pt} \cong \mathrm{Coh}_0^n(X) \times_{\mathrm{Vect}} \mathrm{pt} =: \mathrm{Quot}_{\mathcal{O}_X^n}^{n,\mathrm{fr}}(X)$$

as desired. □

The stack $\mathrm{Coh}_0^n(X)$ should be thought of as the analog of the (GL_n) character stack when X is not affine. Because

$$\mathrm{Coh}_0^n(X) \subset \mathrm{Coh}_{\mathrm{prop}}(X)$$

is an open and closed substack, its inclusion into $\mathrm{Coh}_{\mathrm{prop}}(X)$ is formally étale. If X is furthermore smooth, there is a map

$$\mathrm{Coh}_{\mathrm{prop}}(X) \rightarrow \mathrm{Perf}_{\mathrm{prop}}(X) \cong \mathrm{APerf}_{\mathrm{prop}}(X) \times_{\mathrm{APerf}(X)} \mathrm{Perf}(X),$$

which exists by [PS22, Corollary 2.19], and this map is formally étale because it is the base change of a formally étale map by *loc. cit.* Thus, the composition

$$\mathrm{Coh}_0^n(X) \rightarrow \mathrm{Perf}_{\mathrm{prop}}(X)$$

is formally étale. By [BD18], when X is Gorenstein and Calabi-Yau of dimension d , the moduli stack of perfect complexes with proper support has a natural $(2 - d)$ -shifted symplectic structure. We have thus nearly obtained the following.

Theorem 5.2. Let X be a classical, smooth, quasi-projective scheme of dimension d equipped with a trivialization of its canonical bundle $\mathcal{O}_X \cong K_X$. Then the stack $\mathrm{Coh}_0^n(X) \cong \mathrm{Quot}_{\mathcal{O}_X^n}^{n, \mathrm{fr}}(X)/\mathrm{GL}_n$ has a natural $(2 - d)$ -shifted symplectic structure.

Proof. Denote the symplectic structure on $\mathrm{Perf}_{\mathrm{prop}}(X)$ by ω , and let

$$f : \mathrm{Coh}_0^n(X) \rightarrow \mathrm{Perf}_{\mathrm{prop}}(X)$$

be the map constructed above. Note that $f^*\omega$ is a closed 2-form of degree $2 - d$ on $\mathrm{Coh}_0^n(X)$, and we need only show that it is nondegenerate. By nondegeneracy of ω , the induced map

$$T_{\mathrm{Perf}_{\mathrm{prop}}(X)} \rightarrow \mathbb{L}_{\mathrm{Perf}_{\mathrm{prop}}(X)}[2 - d]$$

is an isomorphism. By the above, f is formally étale. Since the relative cotangent complex of a formally étale map vanishes, we find that

$$T_{\mathrm{Coh}_0^n(X)} = f^*T_{\mathrm{Perf}_{\mathrm{prop}}(X)}$$

and

$$\mathbb{L}_{\mathrm{Coh}_0^n(X)} = f^*\mathbb{L}_{\mathrm{Perf}_{\mathrm{prop}}(X)}.$$

Thus, the map

$$T_{\mathrm{Coh}_0^n(X)} \rightarrow \mathbb{L}_{\mathrm{Coh}_0^n(X)}[2 - d]$$

induced by $f^*\omega$ is an isomorphism as well. □

A Comparison with Existing Constructions

A.1 Quot Schemes

Recall the construction of [Adh22, Definition 3.3], denoted by $\mathrm{Quot}_{\mathcal{O}_X^n}^{\mathrm{Adh}}(X)$, where X is a nonsingular quasi-projective scheme. For such X , define an open subfunctor of this scheme, $\mathrm{Quot}_{\mathcal{O}_X^n}^{\mathrm{Adh},n,\mathrm{fr}}(X) \subset \mathrm{Quot}_{\mathcal{O}_X^n}^{\mathrm{Adh}}(X)$, whose S -points are length n quotients

$$\mathcal{O}_{X \times S}^n \rightarrow \mathcal{E}$$

for which the adjoint map, under the adjunction

$$\mathrm{Hom}_{X \times S}(\mathcal{O}_{X \times S}^n, \mathcal{E}) \cong \mathrm{Hom}_S(\mathcal{O}_S^n, p_{S*}\mathcal{E}),$$

where $p_S : X \times S \rightarrow S$ is the projection, is an isomorphism.

Proposition A.1. Let X be a smooth quasi-projective scheme. Then

$$\mathrm{Quot}_{\mathcal{O}_X^n}^{n,\mathrm{fr}}(X) \cong \mathrm{Quot}_{\mathcal{O}_X^n}^{\mathrm{Adh},n,\mathrm{fr}}(X).$$

Proof. Let $\mathcal{E} \in \mathrm{Coh}_0^n(X)(S)$, for S a test affine. By definition, an S -point of Adhikari's Quot scheme consists of a morphism

$$q : \mathcal{O}_{X \times S}^n \rightarrow \mathcal{E},$$

where $H^0(q)$ is surjective and $\mathcal{E}$ is perfect, is flat over S , has proper support over S , and has geometric fibers of length n . We will construct a map

$$Y := \mathrm{Quot}_{\mathcal{O}_X^n}^{n,\mathrm{fr}}(X) \rightarrow \mathrm{Quot}_{\mathcal{O}_X^n}^{\mathrm{Adh},n,\mathrm{fr}}(X).$$

Recall the universal representation $\mathcal{R}$ of Section 4.2, which is an almost perfect complex with proper support on $X \times Y$ (Lemma 4.12). It is equipped with a universal framing

$$\alpha_{\mathcal{R}} : \mathcal{O}_Y^n \xrightarrow{\cong} p_{Y*}\mathcal{R},$$

where $p_Y : X \times Y \rightarrow Y$. Consider the sheaf morphism $q_{\mathcal{R}}$ on $X \times Y$ given by the composition

$$q_{\mathcal{R}} : \mathcal{O}_{X \times Y}^n \xrightarrow{p_Y^* \alpha_{\mathcal{R}}} p_Y^* p_{Y*} \mathcal{R} \xrightarrow{\varepsilon_{\mathcal{R}}} \mathcal{R},$$

where ε is the counit of the adjunction (p_Y^*, p_{Y*}) . We next claim that $H^0(q_{\mathcal{R}})$ is surjective. Denoting $C = \mathrm{coker}(H^0(q_{\mathcal{R}}))$, it is sufficient to show that the restriction of C to every geometric point is 0, by Nakayama and the fact that C is of finite type. Thus, let (x, y) be a geometric point of $X \times Y$, and set $K = k(y)$ and $\Omega = k(x, y)$. Let $\mathcal{R}_y$ denote the restriction of $\mathcal{R}$ to X_y . The scheme theoretic support of $\mathcal{R}_y$ is a finite scheme $\mathrm{Spec} B$ over $\mathrm{Spec} K$. Thus, we see that

$$\mathcal{R}_y \cong i_* \widetilde{M},$$

where $i : \text{Spec } B \rightarrow X_y$ and $M = \Gamma(Z, \widetilde{M}) \cong \Gamma(X_y, \mathcal{R}_y)$. Supposing that (x, y) is in the support (otherwise the fiber is 0), we obtain a map $B \rightarrow \Omega$; hence,

$$H^0(\mathcal{R}_{(x,y)}) \cong \Omega \otimes_B M.$$

In particular, the restriction of $H^0(q_{\mathcal{R}})$ to (x, y) is a map

$$\Omega^n \xrightarrow{\Omega \otimes_K \alpha_y} \Omega \otimes_K M \rightarrow \Omega \otimes_B M,$$

where

$$\alpha_y : K^n \xrightarrow{\cong} \Gamma(X_y, E_y)$$

The first arrow in the composition is an isomorphism, and the second arrow is surjective. Indeed, the base change over B is obtained from the base change over K by imposing additional relations. Thus, we see that by the representing property of $\text{Quot}_{\mathcal{O}_X^n}^{\text{Adh}}(X)$, we have defined a map

$$f : Y \rightarrow \text{Quot}_{\mathcal{O}_X^n}^{\text{Adh}}(X).$$

Since the adjoint of the map $q_{\mathcal{R}}$ is an equivalence

$$\mathcal{O}_Y^n \xrightarrow{\cong} p_{Y*} \mathcal{R},$$

we see that f factors through the open subfunctor $U := \text{Quot}_{\mathcal{O}_X^n}^{\text{Adh}, n, \text{fr}}(X) \subset \text{Quot}_{\mathcal{O}_X^n}^{\text{Adh}}(X)$. Going through the same procedure, one can define an inverse by using the universal quotient

$$q_U : \mathcal{O}_{X \times U}^n \rightarrow \mathcal{E},$$

where $\mathcal{E}$ is the universal family on $X \times U$, defined in the same procedure as was used to define $\mathcal{R}$. The adjoint to q_U is

$$\mathcal{O}_U^n \xrightarrow{\eta} p_{U*} p_U^* \mathcal{O}_U^n \xrightarrow{p_U^* q_U} p_{U*} \mathcal{E};$$

by definition of q_U , this is an equivalence. Hence, we have a well-defined map

$$g : U \rightarrow Y.$$

It is easy to see that f and g are mutually inverse. □

References

- [Adh22] Nachiketa Adhikari. “Derived quot schemes”. In: arXiv:2211.13845 (Nov. 2022). arXiv:2211.13845 [math]. DOI: 10.48550/arXiv.2211.13845. URL: <http://arxiv.org/abs/2211.13845>.
- [AMR96] Jørgen Ellegaard Andersen, Josef Mattes, and Nicolai Reshetikhin. “The poisson structure on the moduli space of flat connections and chord diagrams”. In: *Topology* 35.4 (Oct. 1996), pp. 1069–1083. ISSN: 0040-9383. DOI: 10.1016/0040-9383(95)00059-3.
- [AMR98] Jørgen Ellegaard Andersen, Josef Mattes, and Nicolai Reshetikhin. “Quantization of the Algebra of Chord Diagrams”. In: *Mathematical Proceedings of the Cambridge Philosophical Society* 124.3 (Nov. 1998). arXiv:q-alg/9701018, pp. 451–467. ISSN: 0305-0041, 1469-8064. DOI: 10.1017/S0305004198002813.
- [Aut] the Stacks Project Authors. *Stacks Project*. URL: <https://stacks.math.columbia.edu/>.
- [BBJ18] David Ben-Zvi, Adrien Brochier, and David Jordan. “Quantum character varieties and braided module categories”. en. In: *Selecta Mathematica* 24.5 (Nov. 2018). Company: Springer Distributor: Springer Institution: Springer Label: Springer publisher: Springer International Publishing, pp. 4711–4748. ISSN: 1420-9020. DOI: 10.1007/s00029-018-0426-y.
- [BCS24] Tristan Bozec, Damien Calaque, and Sarah Scherotzke. “Topological field theories associated with Calabi-Yau categories”. In: arXiv:2410.17726 (Oct. 2024). arXiv:2410.17726 [math.AT]. DOI: 10.48550/arXiv.2410.17726. URL: <http://arxiv.org/abs/2410.17726>.
- [BD18] Christopher Brav and Tobias Dyckerhoff. “Relative Calabi-Yau structures II: Shifted Lagrangians in the moduli of objects”. en. In: *arXiv.org* (Dec. 2018). URL: <https://arxiv.org/abs/1812.11913v1>.
- [BFN10] David Ben-Zvi, John Francis, and David Nadler. “Integral Transforms and Drinfeld Centers in Derived Algebraic Geometry”. en. In: arXiv:0805.0157 (Mar. 2010). arXiv:0805.0157 [math]. URL: <http://arxiv.org/abs/0805.0157>.
- [BKR13] Yuri Berest, George Khachatryan, and Ajay Ramadoss. “Derived representation schemes and cyclic homology”. In: *Advances in Mathematics* 245 (2013), pp. 625–689. ISSN: 0001-8708. DOI: <https://doi.org/10.1016/j.aim.2013.06.020>. URL: <https://www.sciencedirect.com/science/article/pii/S0001870813002296>.
- [BKS19] Dennis Borisov, Ludmil Katzarkov, and Artan Sheshmani. “Shifted symplectic structures on derived Quot-stacks”. en. In: *arXiv.org* (Aug. 2019). URL: <https://arxiv.org/abs/1908.03021v2>.
- [BN19] David Ben-Zvi and David Nadler. “Nonlinear Traces”. In: arXiv:1305.7175 (June 2019). arXiv:1305.7175 [math]. DOI: 10.48550/arXiv.1305.7175. URL: <http://arxiv.org/abs/1305.7175>.
- [BR22] Yuri Berest and Ajay C. Ramadoss. “Symmetric Homology is Representation Homology”. In: (2022). arXiv: 2210.10131 [math.AT].

- [CK99] I. Ciocan-Fontanine and M. Kapranov. “Derived Quot schemes”. en. In: *arXiv.org* (May 1999). URL: <https://arxiv.org/abs/math/9905174v2>.
- [Gol09] William M. Goldman. “Trace Coordinates on Fricke spaces of some simple hyperbolic surfaces”. en. In: *arXiv.org* (Jan. 2009). URL: <https://arxiv.org/abs/0901.1404v1>.
- [Gol84] William M Goldman. “The symplectic nature of fundamental groups of surfaces”. In: *Advances in Mathematics* 54.2 (Nov. 1984), pp. 200–225. ISSN: 0001-8708. DOI: 10.1016/0001-8708(84)90040-9.
- [GR17] Dennis Gaitsgory and Nick Rozenblyum. *A Study in Derived Algebraic Geometry: Volume I: Correspondences and Duality*. en. Vol. 221. Mathematical Surveys and Monographs. American Mathematical Society, 2017. URL: <https://bookstore.ams.org/SURV/221.1>.
- [Gro61] Alexander Grothendieck. “Techniques de construction et théorèmes d’existence en géométrie algébrique IV : les schémas de Hilbert”. fr. In: *Séminaire Bourbaki : années 1960/61, exposés 205-222*. Séminaire Bourbaki 6. talk:221. Société mathématique de France, 1961, pp. 249–276. URL: https://www.numdam.org/item/SB_1960-1961__6__249_0/.
- [GS18] Owen Gwilliam and Claudia Scheimbauer. “Duals and adjoints in higher Morita categories”. In: arXiv:1804.10924 (June 2018). arXiv:1804.10924 [math.CT]. DOI: 10.48550/arXiv.1804.10924. URL: <http://arxiv.org/abs/1804.10924>.
- [Hau17] Rune Haugseng. “The higher Morita category of E_n -algebras”. In: *Geometry & Topology* 21.3 (May 2017). arXiv:1412.8459 [math], pp. 1631–1730. ISSN: 1364-0380, 1465-3060. DOI: 10.2140/gt.2017.21.1631.
- [Hin97] Vladimir Hinich. “Homological algebra of homotopy algebras”. In: arXiv:q-alg/9702015 (Feb. 1997). arXiv:q-alg/9702015. DOI: 10.48550/arXiv.q-alg/9702015. URL: <http://arxiv.org/abs/q-alg/9702015>.
- [HL10] Daniel Huybrechts and Manfred Lehn. *The Geometry of Moduli Spaces of Sheaves*. 2nd ed. Cambridge Mathematical Library. Cambridge: Cambridge University Press, 2010. ISBN: 978-0-521-13420-0. DOI: 10.1017/CBO9780511711985. URL: <https://www.cambridge.org/core/books/geometry-of-moduli-spaces-of-sheaves/E69325DA1892E9BA76>.
- [HSS17] Marc Hoyois, Sarah Scherotzke, and Nicolò Sibilla. “Higher traces, noncommutative motives, and the categorified Chern character”. In: arXiv:1511.03589 (Jan. 2017). arXiv:1511.03589 [math]. DOI: 10.48550/arXiv.1511.03589. URL: <http://arxiv.org/abs/1511.03589>.
- [Jor23] David Jordan. “Quantum character varieties”. In: arXiv:2309.06543 (Sept. 2023). arXiv:2309.06543 [math.QA]. DOI: 10.48550/arXiv.2309.06543. URL: <http://arxiv.org/abs/2309.06543>.
- [Kha26] Adeel A. Khan. “Lectures on algebraic stacks”. In: arXiv:2310.12456 (Apr. 2026). arXiv:2310.12456 [math.AG]. DOI: 10.48550/arXiv.2310.12456. URL: <http://arxiv.org/abs/2310.12456>.
- [KV22] Mikhail Kapranov and Eric Vasserot. “The cohomological Hall algebra of a surface and factorization cohomology”. In: arXiv:1901.07641 (Mar. 2022). arXiv:1901.07641 [math]. DOI: 10.48550/arXiv.1901.07641. URL: <http://arxiv.org/abs/1901.07641>.

- [LM85] Alexander Lubotzky and Andy R. Magid. “Varieties of representations of finitely generated groups”. en. In: *Memoirs of the American Mathematical Society* 58.336 (1985), 0–0. ISSN: 0065-9266, 1947-6221. DOI: 10.1090/memo/0336.
- [Lur09a] Jacob Lurie. *Higher topos theory*. en. Annals of mathematics studies. Princeton, N.J: Princeton University Press, 2009. ISBN: 978-0-691-14048-3.
- [Lur09b] Jacob Lurie. “On the Classification of Topological Field Theories”. en. In: arXiv:0905.0465 (May 2009). arXiv:0905.0465 [math]. URL: <http://arxiv.org/abs/0905.0465>.
- [Lur17] Jacob Lurie. *Higher Algebra*. Sept. 2017. URL: <https://www.math.ias.edu/~lurie/papers/HA.pdf>.
- [Lur18] Jacob Lurie. “Spectral Algebraic Geometry”. In: (Feb. 2018). URL: <https://www.math.ias.edu/~lurie/papers/SAG-rootfile.pdf>.
- [Nit05] Nitin Nitsure. “Construction of Hilbert and Quot Schemes”. In: arXiv:math/0504590 (Apr. 2005). arXiv:math/0504590. DOI: 10.48550/arXiv.math/0504590. URL: <http://arxiv.org/abs/math/0504590>.
- [NS18] Thomas Nikolaus and Peter Scholze. “On topological cyclic homology”. In: arXiv:1707.01799 (Sept. 2018). arXiv:1707.01799 [math.AT]. DOI: 10.48550/arXiv.1707.01799. URL: <http://arxiv.org/abs/1707.01799>.
- [Pan+11] T. Pantev et al. “Shifted Symplectic Structures”. en. In: *arXiv.org* (Nov. 2011). URL: <https://arxiv.org/abs/1111.3209v4>.
- [PS22] Mauro Porta and Francesco Sala. “Two-dimensional categorified Hall algebras”. In: *Journal of the European Mathematical Society* 25.11 (Nov. 2022). arXiv:1903.07253 [math], pp. 4221–4289. ISSN: 1435-9855, 1435-9863. DOI: 10.4171/JEMS/1264.
- [Saf16] Pavel Safronov. “Quasi-Hamiltonian reduction via classical Chern-Simons theory”. In: *Advances in Mathematics* 287 (Jan. 2016). arXiv:1311.6429 [math], pp. 733–773. ISSN: 00018708. DOI: 10.1016/j.aim.2015.09.031.
- [Toe06] B. Toen. “The homotopy theory of dg-categories and derived Morita theory”. In: arXiv:math/0408337 (Oct. 2006). arXiv:math/0408337. DOI: 10.48550/arXiv.math/0408337. URL: <http://arxiv.org/abs/math/0408337>.
- [Tur91] Vladimir G. Turaev. “Skein quantization of Poisson algebras of loops on surfaces”. en. In: *Annales scientifiques de l’École Normale Supérieure* 24.6 (1991), pp. 635–704. ISSN: 1873-2151. DOI: 10.24033/asens.1639.
- [TV07] B. Toen and M. Vaquie. “Moduli of objects in dg-categories”. In: arXiv:math/0503269 (May 2007). arXiv:math/0503269. DOI: 10.48550/arXiv.math/0503269. URL: <http://arxiv.org/abs/math/0503269>.
- [Yeu18] Wai-Kit Yeung. “Pre-Calabi-Yau structures and moduli of representations”. en. In: *arXiv.org* (Feb. 2018). URL: <https://arxiv.org/abs/1802.05398v4>.