

MATH512 - Project 4

Wasif Ahmed, Haoxiang Deng, Jacob Fein-Ashley, Kanav Malhotra, Longzhe Yang

April 2024

Question 1

Refer to figure ?? and ??.




Figure 1: $|\delta W_i|$




Figure 2: δW_i^2

Notice that as the L parameter increases, the $|\delta W_i|$ term is unbounded while δW_i^2 converges to 2 in probability.

Question 2




Figure 3: 2a

1.




Figure 4: 2b

2.




Figure 5: 2c

3.




Figure 6: 2d

4.




Figure 7: 2e

5.




Figure 8: 2f

6.

Question 3

Objective: Analyze the weak and strong order of convergence of the EM method for solving the given SDE:

$$dX(t) = \mu X(t)dt + \sigma X(t)dW(t), \quad X(0) = 3, \quad \mu = 2, \quad \sigma = 0.10$$

- **Weak order of convergence equal to 1:** to show that

$$|E[X_1] - E[X(1)]| = C\Delta t$$

- **Strong order of convergence equal to 0.5:** to show that

$$E|X_1 - X(1)| = C\Delta t^{0.5}$$

where X_1 stands for the exact solution and $X(1)$ stands for the estimated solution.

Weak Order of Convergence focuses on the **expected values** of the numerical solution compared to the exact solution

$$|E[X_1] - E[X(1)]| = C\Delta t$$

- A plot of the error versus Δt for the weak order of convergence of 1.

imgs/fig1.png

Strong Order of Convergence to 0.5 is concerned with the **pathwise accuracy**, evaluating how closely the numerical solution follows individual realizations of the exact solution.

$$E|X_1 - X(1)| = C\Delta t^{0.5}$$

- A plot of the error versus $\Delta t^{0.5}$ for the strong order of convergence of 0.5.

imgs/fig2.png

Question 4




Figure 9: Implicit Euler




Figure 10: Explicit Euler




Figure 11: Mean Square Stable Values

Question 5

An algorithm to simulate exit times for the SDE:

- 1: Choose a step size Δt
- 2: Choose some paths, M

```

3: for  $s = 1$  to  $M$  do
4:   Set  $t_n = 0$  and  $X_n = X_0$ 
5:   while  $X_n > a$  and  $X_n < b$  do
6:     Compute a  $N(0, 1)$  sample  $\xi_n$ 
7:     Replace  $X_n$  by  $X_n + \mu X_n \Delta t + \sigma X_n \xi_n$ 
8:     Replace  $t_n$  by  $t_n + \Delta t$ 
9:   end while
10:  Set  $T_s^{exit} = t_n - 1/2\Delta t$ 
11: end for
12: Set  $a_M = \frac{1}{M} \sum_{s=1}^M T_s^{exit}$ 
13: Set  $b_M^2 = \frac{1}{M-1} \sum_{s=1}^M (T_s^{exit} - a_M)^2$ 

```




Figure 12: 5




Figure 13: 5