Jacob Hauck

Math 6010

9-20-2023

# Objectives

▶ Define exponential functions  $(f(x) = C \cdot a^x)$ 

### Objectives

▶ Define exponential functions  $(f(x) = C \cdot a^x)$ 

▶ Understand basic properties of exponential functions

# Objectives

▶ Define exponential functions  $(f(x) = C \cdot a^x)$ 

▶ Understand basic properties of exponential functions

► Graph exponential functions

Want a function like

$$f(x) = C \cdot a^x$$

Want a function like

$$f(x) = C \cdot a^x$$

Recall: if 
$$x = \frac{m}{n}$$
, then  $a^x = a^{\frac{m}{n}} = \sqrt[n]{a^m}$ 

Want a function like

$$f(x) = C \cdot a^x$$

Recall: if 
$$x = \frac{m}{n}$$
, then  $a^x = a^{\frac{m}{n}} = \sqrt[n]{a^m}$ 

But x might not be  $\frac{m}{n}$ , for integers m, n...

Want a function like

$$f(x) = C \cdot a^x$$

Recall: if  $x = \frac{m}{n}$ , then  $a^x = a^{\frac{m}{n}} = \sqrt[n]{a^m}$ 

But x might not be  $\frac{m}{n}$ , for integers m, n...

 $ightharpoonup x = \pi, \sqrt{2}$ , and so on

Want a function like

$$f(x) = C \cdot a^x$$

Recall: if 
$$x = \frac{m}{n}$$
, then  $a^x = a^{\frac{m}{n}} = \sqrt[n]{a^m}$ 

But x might not be  $\frac{m}{n}$ , for integers m, n...

$$ightharpoonup x = \pi, \sqrt{2}$$
, and so on

...but every number has a decimal expansion

$$3 = 3,$$
  $\frac{2}{5} = 0.4,$   $\pi = 3.14159265...,$   $\sqrt{2} = 1.41421356...$ 

Want a function like

$$f(x) = C \cdot a^x$$

Recall: if 
$$x = \frac{m}{n}$$
, then  $a^x = a^{\frac{m}{n}} = \sqrt[n]{a^m}$ 

But x might not be  $\frac{m}{n}$ , for integers m, n...

$$ightharpoonup x = \pi, \sqrt{2}$$
, and so on

...but every number has a decimal expansion

$$3 = 3,$$
  $\frac{2}{5} = 0.4,$   $\pi = 3.14159265...,$   $\sqrt{2} = 1.41421356...$ 

Approximate any number x by finitely many of its decimal digits.

$$a^{\pi} \approx a^{3.14} = a^{\frac{314}{100}} = \sqrt[100]{a^{314}}$$

The more digits we use, the closer we get to the true value of  $a^{\pi}$ 

$2^3$	$2^{3.1}$	$2^{3.14}$	$2^{3.141}$	$2^{3.1415}$	 $2^{\pi}$
8.0000	8.5742	8.8152	8.8213	8.8244	 8.8250

The more digits we use, the closer we get to the true value of  $a^{\pi}$ 

$2^3$	$2^{3.1}$	$2^{3.14}$	$2^{3.141}$	$2^{3.1415}$	• • •	$2^{\pi}$
8.0000	8.5742	8.8152	8.8213	8.8244		8.8250

So we can use any number x in the exponent. Hooray!

The more digits we use, the closer we get to the true value of  $a^{\pi}$ 

$2^3$	$2^{3.1}$	$2^{3.14}$	$2^{3.141}$	$2^{3.1415}$	 $2^{\pi}$
8.0000	8.5742	8.8152	8.8213	8.8244	 8.8250

So we can use any number x in the exponent. Hooray!

Better still, we get to keep all the properties of exponents that we already know:

#### Exponent Properties

If x, y are real numbers, and a, b > 0, then

$$\bullet a^{x} \cdot a^{y} = a^{x+y} \quad \bullet (a^{x})^{y} = a^{xy} \quad \bullet (ab)^{x} = a^{x} \cdot b^{x}$$

$$\bullet 1^{x} = 1 \quad \bullet a^{-x} = \frac{1}{a^{x}} = \left(\frac{1}{a}\right)^{x} \quad \bullet a^{0} = 1$$

$$\bullet \ 1^x = 1 \qquad \bullet \ a^{-x} = \frac{1}{a^x} = \left(\frac{1}{a}\right)^x \quad \bullet \ a^0 = 1$$

Now that we know what  $a^x$  means for any x, we can define an exponential function.

#### Exponential Function

An **exponential function** is a function f such that

$$f(x) = C \cdot a^x,$$

where

- $ightharpoonup C \neq 0$  is called the **initial value**, and
- ▶ a > 0,  $a \neq 1$  is called the **growth factor**.

Consider the exponential function  $f(x) = 5 \cdot 2^x$ . Let's make a table of values

$$\begin{array}{ccc} x & f(x) \\ -2 & 1.25 \\ -1 & 2.5 \\ 0 & 5 \\ 1 & 10 \\ 2 & 20 \\ \end{array}$$

Consider the exponential function  $f(x) = 5 \cdot 2^x$ . Let's make a table of values

$$\begin{array}{ccc} x & f(x) \\ -2 & 1.25 \\ -1 & 2.5 \\ 0 & 5 \\ 1 & 10 \\ 2 & 20 \end{array}$$

ightharpoonup f(x) goes up by  $\times 2 = a$  each time x increases

Consider the exponential function  $f(x) = 5 \cdot 2^x$ . Let's make a table of values

$$\begin{array}{c|cc} x & f(x) \\ \hline -2 & 1.25 \\ -1 & 2.5 \\ 0 & 5 \\ 1 & 10 \\ 2 & 20 \\ \end{array}$$

- ightharpoonup f(x) goes up by  $\times 2 = a$  each time x increases
- f(0) = 5 = C.

$$f(x) = C \cdot a^x$$

•  $f(0) = C \cdot a^0 = C$ , the initial value

$$f(x) = C \cdot a^x$$

- $ightharpoonup f(0) = C \cdot a^0 = C$ , the initial value
- ► The growth factor: how much the function grows (or shrinks) every time x goes up by 1

$$f(x+1) = a \cdot f(x)$$

$$f(x) = C \cdot a^x$$

- $ightharpoonup f(0) = C \cdot a^0 = C$ , the initial value
- ► The growth factor: how much the function grows (or shrinks) every time x goes up by 1

$$f(x+1) = a \cdot f(x)$$

because

$$\frac{f(x+1)}{f(x)} = \frac{C \cdot a^{x+1}}{C \cdot a^x}$$

$$f(x) = C \cdot a^x$$

- $ightharpoonup f(0) = C \cdot a^0 = C$ , the initial value
- The growth factor: how much the function grows (or shrinks) every time x goes up by 1

$$f(x+1) = a \cdot f(x)$$

because

$$\frac{f(x+1)}{f(x)} = \frac{C \cdot a^{x+1}}{C \cdot a^x} = \frac{a^{x+1}}{a^x}$$

$$f(x) = C \cdot a^x$$

- $ightharpoonup f(0) = C \cdot a^0 = C$ , the initial value
- ► The growth factor: how much the function grows (or shrinks) every time x goes up by 1

$$f(x+1) = a \cdot f(x)$$

because

$$\frac{f(x+1)}{f(x)} = \frac{C \cdot a^{x+1}}{C \cdot a^x} = \frac{a^{x+1}}{a^x} = a^{x+1-x} = a$$

Let's try to graph the function  $f(x) = 5 \cdot 2^x$  from before. First, extend the table of values

Let's try to graph the function  $f(x) = 5 \cdot 2^x$  from before. First, extend the table of values

j

Let's try to graph the function  $f(x) = 5 \cdot 2^x$  from before. First, extend the table of values

x	f(x)
-5	0.16125
-4	0.3125
-3	0.625
-2	1.25
-1	2.5
0	5
1	10
2	20
3	40
4	80
5	160

It seems that  $f(x) \to 0$  as  $x \to -\infty$ , and  $f(x) \to \infty$  as  $x \to \infty$ 







