

The Fréchet Derivative

Jacob Hauck

Math 6418

Motivation

Let X, Y be normed linear spaces. We know a lot about linear maps $A : X \rightarrow Y$ – what can we do about nonlinear maps?

Motivation

Let X, Y be normed linear spaces. We know a lot about linear maps $A : X \rightarrow Y$ – what can we do about nonlinear maps?

Approximate a nonlinear map $f : U \rightarrow Y$ by a linear operator $A \in B(X, Y)$.

Motivation

Say $X = Y = \mathbf{R}$, then A_{x_0} is given by multiplication by a number $a_{x_0} \in \mathbf{R}$, and a natural choice for a_{x_0} is $f'(x_0)$, as

$$f(x) - f(x_0) \approx f'(x_0)(x - x_0).$$

Motivation

Say $X = Y = \mathbf{R}$, then A_{x_0} is given by multiplication by a number $a_{x_0} \in \mathbf{R}$, and a natural choice for a_{x_0} is $f'(x_0)$, as

$$f(x) - f(x_0) \approx f'(x_0)(x - x_0).$$

By definition,

$$\left| \frac{f(x_0 + h) - f(x_0)}{h} - f'(x_0) \right| \rightarrow 0 \quad \text{as} \quad h \rightarrow 0.$$

Motivation

Say $X = Y = \mathbf{R}$, then A_{x_0} is given by multiplication by a number $a_{x_0} \in \mathbf{R}$, and a natural choice for a_{x_0} is $f'(x_0)$, as

$$f(x) - f(x_0) \approx f'(x_0)(x - x_0).$$

By definition,

$$\left| \frac{f(x_0 + h) - f(x_0)}{h} - f'(x_0) \right| \rightarrow 0 \quad \text{as } h \rightarrow 0.$$

How can we generalize to the case that $X, Y \neq \mathbf{R}$? Can't divide by h , so rewrite as

$$\frac{|f(x_0 + h) - f(x_0) - f'(x_0)h|}{|h|} \rightarrow 0 \quad \text{as } h \rightarrow 0.$$

Definition of the Fréchet Derivative

Definition (Fréchet Derivative)

A function $f : U \rightarrow Y$ is called **Fréchet differentiable** at $x \in U$ if there exists $A \in B(X, Y)$ such that

$$\frac{\|f(x+h) - f(x) - Ah\|_Y}{\|h\|_X} \rightarrow 0 \quad \text{as} \quad \|h\|_X \rightarrow 0,$$

in which case A is called the **Fréchet derivative** of f at x , also denoted by

$$A = f'(x) = Df(x)$$

Definition of the Fréchet Derivative

Definition (Fréchet Derivative)

A function $f : U \rightarrow Y$ is called **Fréchet differentiable** at $x \in U$ if there exists $A \in B(X, Y)$ such that

$$\frac{\|f(x+h) - f(x) - Ah\|_Y}{\|h\|_X} \rightarrow 0 \quad \text{as} \quad \|h\|_X \rightarrow 0,$$

in which case A is called the **Fréchet derivative** of f at x , also denoted by

$$A = f'(x) = Df(x)$$

The Fréchet derivative is the same as the usual derivative if $f \in C^1(\mathbf{R})$.

Basic Properties

1. Uniqueness (justifying the phrase “*the* Fréchet derivative”)

Basic Properties

1. Uniqueness (justifying the phrase “*the* Fréchet derivative”)
2. Linearity

$$D(\alpha f + \beta g)(x) = \alpha Df(x) + \beta Dg(x)$$

Basic Properties

1. Uniqueness (justifying the phrase “*the* Fréchet derivative”)
2. Linearity

$$D(\alpha f + \beta g)(x) = \alpha Df(x) + \beta Dg(x)$$

3. Fréchet differentiable \implies continuous

Basic Properties

1. Uniqueness (justifying the phrase “*the* Fréchet derivative”)
2. Linearity

$$D(\alpha f + \beta g)(x) = \alpha Df(x) + \beta Dg(x)$$

3. Fréchet differentiable \implies continuous
4. Fréchet differentiable \implies locally Lipschitz
 $f : X \rightarrow Y$ is **locally Lipschitz** at $x \in X$ if there exists $\delta > 0$ and $L > 0$ such that

$$\|f(x) - f(y)\|_Y \leq L\|x - y\|_X \quad \text{if} \quad \|x - y\| < \delta.$$

Basic Properties

1. Uniqueness (justifying the phrase “*the* Fréchet derivative”)
2. Linearity

$$D(\alpha f + \beta g)(x) = \alpha Df(x) + \beta Dg(x)$$

3. Fréchet differentiable \implies continuous
4. Fréchet differentiable \implies locally Lipschitz
 $f : X \rightarrow Y$ is **locally Lipschitz** at $x \in X$ if there exists $\delta > 0$ and $L > 0$ such that

$$\|f(x) - f(y)\|_Y \leq L\|x - y\|_X \quad \text{if} \quad \|x - y\| < \delta.$$

For any $\varepsilon > 0$, can choose δ small enough so that $L = \|Df(x)\|_{B(X,Y)} + \varepsilon$ works.

Examples – Linear Operators

Example 1. Suppose that $f : X \rightarrow Y$ is actually linear: $f(x) = Ax$, where $A \in B(X, Y)$. Then, as expected,

$$Df(x) = A \quad \text{for all } x \in X.$$

Examples – Linear Operators

Example 1. Suppose that $f : X \rightarrow Y$ is actually linear: $f(x) = Ax$, where $A \in B(X, Y)$. Then, as expected,

$$Df(x) = A \quad \text{for all } x \in X.$$

The proof is trivial due to the linearity of A :

$$\frac{\|f(x+h) - f(x) - Ah\|_Y}{\|h\|_X}$$

Examples – Linear Operators

Example 1. Suppose that $f : X \rightarrow Y$ is actually linear: $f(x) = Ax$, where $A \in B(X, Y)$. Then, as expected,

$$Df(x) = A \quad \text{for all } x \in X.$$

The proof is trivial due to the linearity of A :

$$\frac{\|f(x+h) - f(x) - Ah\|_Y}{\|h\|_X} = \frac{\|A(x+h) - Ax - Ah\|_Y}{\|h\|_X} = 0.$$

Examples – A “Quadratic” Operator

Example 2. Suppose that $X = H$ is a Hilbert space over \mathbf{R} , and $f(x) = (x, Ax)$, where $A \in B(H, H)$.

Examples – A “Quadratic” Operator

Example 2. Suppose that $X = H$ is a Hilbert space over \mathbf{R} , and $f(x) = (x, Ax)$, where $A \in B(H, H)$. Then $f : H \rightarrow \mathbf{R}$, and

$$\begin{aligned} f(x + h) - f(x) &= (x + h, A(x + h)) - (x, Ax) \\ &= (x, Ah) + (h, Ax) + (h, Ah) \end{aligned}$$

Examples – A “Quadratic” Operator

Example 2. Suppose that $X = H$ is a Hilbert space over \mathbf{R} , and $f(x) = (x, Ax)$, where $A \in B(H, H)$. Then $f : H \rightarrow \mathbf{R}$, and

$$\begin{aligned} f(x + h) - f(x) &= (x + h, A(x + h)) - (x, Ax) \\ &= (x, Ah) + (h, Ax) + (h, Ah) \\ &= (A^*x + Ax, h) + (h, Ah). \end{aligned}$$

Examples – A “Quadratic” Operator

Example 2. Suppose that $X = H$ is a Hilbert space over \mathbf{R} , and $f(x) = (x, Ax)$, where $A \in B(H, H)$. Then $f : H \rightarrow \mathbf{R}$, and

$$\begin{aligned} f(x+h) - f(x) &= (x+h, A(x+h)) - (x, Ax) \\ &= (x, Ah) + (h, Ax) + (h, Ah) \\ &= (A^*x + Ax, h) + (h, Ah). \end{aligned}$$

Then, viewing $(A^* + A)x$ as an element of $H^* = B(H, \mathbf{R})$,

$$Df(x) = (A^* + A)x$$

Examples – A “Quadratic” Operator

Example 2. Suppose that $X = H$ is a Hilbert space over \mathbf{R} , and $f(x) = (x, Ax)$, where $A \in B(H, H)$. Then $f : H \rightarrow \mathbf{R}$, and

$$\begin{aligned} f(x+h) - f(x) &= (x+h, A(x+h)) - (x, Ax) \\ &= (x, Ah) + (h, Ax) + (h, Ah) \\ &= (A^*x + Ax, h) + (h, Ah). \end{aligned}$$

Then, viewing $(A^* + A)x$ as an element of $H^* = B(H, \mathbf{R})$,

$$Df(x) = (A^* + A)x$$

because, by the Cauchy-Schwarz inequality,

$$\frac{|f(x+h) - f(x) - ((A^* + A)x, h)|}{\|h\|_X} = \frac{|(h, Ah)|}{\|h\|_X}$$

Examples – A “Quadratic” Operator

Example 2. Suppose that $X = H$ is a Hilbert space over \mathbf{R} , and $f(x) = (x, Ax)$, where $A \in B(H, H)$. Then $f : H \rightarrow \mathbf{R}$, and

$$\begin{aligned} f(x+h) - f(x) &= (x+h, A(x+h)) - (x, Ax) \\ &= (x, Ah) + (h, Ax) + (h, Ah) \\ &= (A^*x + Ax, h) + (h, Ah). \end{aligned}$$

Then, viewing $(A^* + A)x$ as an element of $H^* = B(H, \mathbf{R})$,

$$Df(x) = (A^* + A)x$$

because, by the Cauchy-Schwarz inequality,

$$\begin{aligned} \frac{|f(x+h) - f(x) - ((A^* + A)x, h)|}{\|h\|_X} &= \frac{|(h, Ah)|}{\|h\|_X} \\ &\leq \frac{\|h\|_X \|Ah\|_X}{\|h\|_X} = \|Ah\|_X, \end{aligned}$$

Examples – A “Quadratic” Operator

Example 2. Suppose that $X = H$ is a Hilbert space over \mathbf{R} , and $f(x) = (x, Ax)$, where $A \in B(H, H)$. Then $f : H \rightarrow \mathbf{R}$, and

$$\begin{aligned} f(x+h) - f(x) &= (x+h, A(x+h)) - (x, Ax) \\ &= (x, Ah) + (h, Ax) + (h, Ah) \\ &= (A^*x + Ax, h) + (h, Ah). \end{aligned}$$

Then, viewing $(A^* + A)x$ as an element of $H^* = B(H, \mathbf{R})$,

$$Df(x) = (A^* + A)x$$

because, by the Cauchy-Schwarz inequality,

$$\begin{aligned} \frac{|f(x+h) - f(x) - ((A^* + A)x, h)|}{\|h\|_X} &= \frac{|(h, Ah)|}{\|h\|_X} \\ &\leq \frac{\|h\|_X \|Ah\|_X}{\|h\|_X} = \|Ah\|_X, \end{aligned}$$

and $\|Ah\|_X \rightarrow 0$ as $h \rightarrow 0$.

Examples – Gradient

Example 3. Let $f \in C^1(\mathbf{R}^n)$ (first-order partial derivatives exist and are continuous)

Examples – Gradient

Example 3. Let $f \in C^1(\mathbf{R}^n)$ (first-order partial derivatives exist and are continuous)

Guess: $Df(x) = \nabla f(x)$, viewing $\nabla f(x)$ as a linear operator defined by $[\nabla f(x)](h) = [\nabla f(x)]^T h$, for $h \in \mathbf{R}^n$.

Examples – Gradient

Example 3. Let $f \in C^1(\mathbf{R}^n)$ (first-order partial derivatives exist and are continuous)

Guess: $Df(x) = \nabla f(x)$, viewing $\nabla f(x)$ as a linear operator defined by $[\nabla f(x)](h) = [\nabla f(x)]^T h$, for $h \in \mathbf{R}^n$.

Let $n = 2$, and define $\tilde{h} = h_1 e_1$. Then

$$|f(x+h) - f(x) - \nabla f(x)^T h| = |f(x+h) - f(x) - \partial_1 f(x) h_1 - \partial_2 f(x) h_2|$$

Examples – Gradient

Example 3. Let $f \in C^1(\mathbf{R}^n)$ (first-order partial derivatives exist and are continuous)

Guess: $Df(x) = \nabla f(x)$, viewing $\nabla f(x)$ as a linear operator defined by $[\nabla f(x)](h) = [\nabla f(x)]^T h$, for $h \in \mathbf{R}^n$.

Let $n = 2$, and define $\tilde{h} = h_1 e_1$. Then

$$\begin{aligned} |f(x+h) - f(x) - \nabla f(x)^T h| &= |f(x+h) - f(x) - \partial_1 f(x) h_1 - \partial_2 f(x) h_2| \\ &= |f(x+h) - f(x+\tilde{h}) - \partial_2 f(x+\tilde{h}) h_2 \\ &\quad + f(x+\tilde{h}) - f(x) - \partial_1 f(x) h_1 \\ &\quad + (\partial_2 f(x+\tilde{h}) - \partial_2 f(x)) h_2| \end{aligned}$$

Examples – Gradient

Example 3. Let $f \in C^1(\mathbf{R}^n)$ (first-order partial derivatives exist and are continuous)

Guess: $Df(x) = \nabla f(x)$, viewing $\nabla f(x)$ as a linear operator defined by $[\nabla f(x)](h) = [\nabla f(x)]^T h$, for $h \in \mathbf{R}^n$.

Let $n = 2$, and define $\tilde{h} = h_1 e_1$. Then

$$\begin{aligned} |f(x+h) - f(x) - \nabla f(x)^T h| &= |f(x+h) - f(x) - \partial_1 f(x) h_1 - \partial_2 f(x) h_2| \\ &\leq |f(x+h) - f(x+\tilde{h}) - \partial_2 f(x+\tilde{h}) h_2| \\ &\quad + |f(x+\tilde{h}) - f(x) - \partial_1 f(x) h_1| \\ &\quad + |\partial_2 f(x+\tilde{h}) - \partial_2 f(x)| \cdot |h_2| \end{aligned}$$

Examples – Gradient

Example 3. Let $f \in C^1(\mathbf{R}^n)$ (first-order partial derivatives exist and are continuous)

Guess: $Df(x) = \nabla f(x)$, viewing $\nabla f(x)$ as a linear operator defined by $[\nabla f(x)](h) = [\nabla f(x)]^T h$, for $h \in \mathbf{R}^n$.

Let $n = 2$, and define $\tilde{h} = h_1 e_1$. Then

$$\begin{aligned} |f(x+h) - f(x) - \nabla f(x)^T h| &= |f(x+h) - f(x) - \partial_1 f(x) h_1 - \partial_2 f(x) h_2| \\ &\leq |f(x+h) - f(x+\tilde{h}) - \partial_2 f(x+\tilde{h}) h_2| \\ &\quad + |f(x+\tilde{h}) - f(x) - \partial_1 f(x) h_1| \\ &\quad + |\partial_2 f(x+\tilde{h}) - \partial_2 f(x)| \cdot |h_2| \end{aligned}$$

Dividing both sides by $\|h\|$ and applying definition of partial derivative and continuity...

Examples – Gradient

First term:

$$\frac{|f(x+h) - f(x+\tilde{h}) - \partial_2 f(x+\tilde{h})h_2|}{\|h\|}$$

Examples – Gradient

First term:

$$\frac{|f(x+h) - f(x+\tilde{h}) - \partial_2 f(x+\tilde{h})h_2|}{\|h\|} = \underbrace{\frac{h_2}{\|h\|}}_{|\cdot| \leq 1} \cdot \underbrace{\frac{|\cdots|}{h_2}}_{\rightarrow 0 \text{ by definition of } \partial_2} \rightarrow 0$$

as $h \rightarrow 0$.

Examples – Gradient

First term:

$$\frac{|f(x+h) - f(x+\tilde{h}) - \partial_2 f(x+\tilde{h})h_2|}{\|h\|} = \underbrace{\frac{h_2}{\|h\|}}_{|\cdot| \leq 1} \cdot \underbrace{\frac{|\cdots|}{h_2}}_{\rightarrow 0 \text{ by definition of } \partial_2} \rightarrow 0$$

as $h \rightarrow 0$.

Second term:

$$\frac{|f(x+\tilde{h}) - f(x) - \partial_1 f(x)h_1|}{\|h\|} = \frac{h_1}{\|h\|} \cdot \frac{|\cdots|}{h_1} \rightarrow 0 \quad \text{as } h \rightarrow 0.$$

Examples – Gradient

First term:

$$\frac{|f(x+h) - f(x+\tilde{h}) - \partial_2 f(x+\tilde{h})h_2|}{\|h\|} = \underbrace{\frac{h_2}{\|h\|}}_{|\cdot| \leq 1} \cdot \underbrace{\frac{|\cdots|}{h_2}}_{\rightarrow 0 \text{ by definition of } \partial_2} \rightarrow 0$$

as $h \rightarrow 0$.

Second term:

$$\frac{|f(x+\tilde{h}) - f(x) - \partial_1 f(x)h_1|}{\|h\|} = \frac{h_1}{\|h\|} \cdot \frac{|\cdots|}{h_1} \rightarrow 0 \quad \text{as } h \rightarrow 0.$$

Third term:

$$|\partial_2 f(x+\tilde{h}) - \partial_2 f(x)| \cdot \frac{|h_2|}{\|h\|} \rightarrow 0 \quad \text{as } h \rightarrow 0$$

by the continuity of $\partial_2 f$.

Examples – Jacobian

Example 4. Let $f : \mathbf{R}^n \rightarrow \mathbf{R}^m$ be C^1 .

Examples – Jacobian

Example 4. Let $f : \mathbf{R}^n \rightarrow \mathbf{R}^m$ be C^1 .

Then, if $J(x) \in \mathbf{R}^{m \times n}$ is the Jacobian matrix of f ,

$$f(x) = \begin{bmatrix} f_1(x) \\ \vdots \\ f_m(x) \end{bmatrix} \implies J(x) = \begin{bmatrix} \nabla f_1(x)^T \\ \vdots \\ \nabla f_m(x)^T \end{bmatrix},$$

Examples – Jacobian

Example 4. Let $f : \mathbf{R}^n \rightarrow \mathbf{R}^m$ be C^1 .

Then, if $J(x) \in \mathbf{R}^{m \times n}$ is the Jacobian matrix of f ,

$$f(x) = \begin{bmatrix} f_1(x) \\ \vdots \\ f_m(x) \end{bmatrix} \implies J(x) = \begin{bmatrix} \nabla f_1(x)^T \\ \vdots \\ \nabla f_m(x)^T \end{bmatrix},$$

and

$$\frac{\|f(x+h) - f(x) - J(x)h\|}{\|h\|} = \frac{\left\| \begin{bmatrix} f_1(x+h) - f_1(x) - \nabla f_1(x)^T h \\ \vdots \\ f_m(x+h) - f_m(x) - \nabla f_m(x)^T h \end{bmatrix} \right\|}{\|h\|}.$$

Examples – Jacobian

Example 4. Let $f : \mathbf{R}^n \rightarrow \mathbf{R}^m$ be C^1 .

Then, if $J(x) \in \mathbf{R}^{m \times n}$ is the Jacobian matrix of f ,

$$f(x) = \begin{bmatrix} f_1(x) \\ \vdots \\ f_m(x) \end{bmatrix} \implies J(x) = \begin{bmatrix} \nabla f_1(x)^T \\ \vdots \\ \nabla f_m(x)^T \end{bmatrix},$$

and

$$\frac{\|f(x+h) - f(x) - J(x)h\|}{\|h\|} = \frac{\left\| \begin{bmatrix} f_1(x+h) - f_1(x) - \nabla f_1(x)^T h \\ \vdots \\ f_m(x+h) - f_m(x) - \nabla f_m(x)^T h \end{bmatrix} \right\|}{\|h\|}.$$

Each component $\rightarrow 0$ as $h \rightarrow 0$ because ∇f_1 is the Fréchet derivative of f_1 ; therefore $Df(x) = J(x)$, interpreting the matrix as an operator.

Examples – Integral of a Power

Example 5. Let $p > 1$ be an integer, and let $f : L^p(\mathbf{R}) \rightarrow \mathbf{R}$ be defined by

$$f(\varphi) = \int \varphi^p.$$

Examples – Integral of a Power

Example 5. Let $p > 1$ be an integer, and let $f : L^p(\mathbf{R}) \rightarrow \mathbf{R}$ be defined by

$$f(\varphi) = \int \varphi^p.$$

Let $h \in L^p(\mathbf{R})$; then

$$\begin{aligned} f(\varphi + h) - f(\varphi) &= \int [(\varphi + h)^p - \varphi^p] \\ &= \int \left[p\varphi^{p-1}h + \binom{p}{2}\varphi^{p-2}h^2 + \cdots + h^p \right] \end{aligned}$$

Examples – Integral of a Power

Example 5. Let $p > 1$ be an integer, and let $f : L^p(\mathbf{R}) \rightarrow \mathbf{R}$ be defined by

$$f(\varphi) = \int \varphi^p.$$

Let $h \in L^p(\mathbf{R})$; then

$$\begin{aligned} f(\varphi + h) - f(\varphi) &= \int [(\varphi + h)^p - \varphi^p] \\ &= \int \left[p\varphi^{p-1}h + \binom{p}{2}\varphi^{p-2}h^2 + \cdots + h^p \right] \end{aligned}$$

Note that $\varphi^{p-k}h^k$ is integrable for $k = 0, 1, \dots, p$:

$$\int |\varphi|^{p-k}|h|^k \leq \left(\int |\varphi|^p \right)^{\frac{1}{u}} \left(\int |h|^p \right)^{\frac{1}{v}} = \|\varphi\|_{L^p(\mathbf{R})}^{p-k} \|h\|_{L^p(\mathbf{R})}^k$$

if $u = \frac{p}{p-k}$ and $v = \frac{p}{k}$.

Examples – Integral of a Power

Therefore,

$$\begin{aligned} \frac{1}{\|h\|_{L^p(\mathbf{R})}} \left| f(\varphi + h) - f(\varphi) - \int p\varphi^{p-1}h \right| \\ = \frac{1}{\|h\|} \left| \int \left[\binom{p}{2} \varphi^{p-2}h^2 + \dots + h^p \right] \right| \end{aligned}$$

Examples – Integral of a Power

Therefore,

$$\begin{aligned} \frac{1}{\|h\|_{L^p(\mathbf{R})}} & \left| f(\varphi + h) - f(\varphi) - \int p\varphi^{p-1}h \right| \\ &= \frac{1}{\|h\|} \left| \int \left[\binom{p}{2} \varphi^{p-2} h^2 + \dots + h^p \right] \right| \\ &\leq \binom{p}{2} \int |\varphi|^{p-2} |h^2| + \dots + \int |h|^p \end{aligned}$$

Examples – Integral of a Power

Therefore,

$$\begin{aligned} \frac{1}{\|h\|_{L^p(\mathbf{R})}} & \left| f(\varphi + h) - f(\varphi) - \int p\varphi^{p-1}h \right| \\ &= \frac{1}{\|h\|} \left| \int \left[\binom{p}{2} \varphi^{p-2} h^2 + \dots + h^p \right] \right| \\ &\leq \binom{p}{2} \int |\varphi|^{p-2} |h^2| + \dots + \int |h|^p \\ &\leq \frac{1}{\|h\|} \left[\binom{p}{2} \|\varphi\|_{L^p(\mathbf{R})}^{p-2} \|h\|_{L^p(\mathbf{R})}^2 + \dots + \|h\|_{L^p(\mathbf{R})}^p \right] \end{aligned}$$

Examples – Integral of a Power

Therefore,

$$\begin{aligned} \frac{1}{\|h\|_{L^p(\mathbf{R})}} & \left| f(\varphi + h) - f(\varphi) - \int p\varphi^{p-1}h \right| \\ &= \frac{1}{\|h\|} \left| \int \left[\binom{p}{2} \varphi^{p-2} h^2 + \dots + h^p \right] \right| \\ &\leq \binom{p}{2} \int |\varphi|^{p-2} |h^2| + \dots + \int |h|^p \\ &\leq \frac{1}{\|h\|} \left[\binom{p}{2} \|\varphi\|_{L^p(\mathbf{R})}^{p-2} \|h\|_{L^p(\mathbf{R})}^2 + \dots + \|h\|_{L^p(\mathbf{R})}^p \right] \\ &\leq C(\varphi, p) (\|h\| + \|h\|^2 + \dots + \|h\|^{p-2}) \end{aligned}$$

which $\rightarrow 0$ as $\|h\| \rightarrow 0$.

Examples – Integral of a Power

Therefore,

$$\begin{aligned} \frac{1}{\|h\|_{L^p(\mathbf{R})}} & \left| f(\varphi + h) - f(\varphi) - \int p\varphi^{p-1}h \right| \\ &= \frac{1}{\|h\|} \left| \int \left[\binom{p}{2} \varphi^{p-2} h^2 + \dots + h^p \right] \right| \\ &\leq \binom{p}{2} \int |\varphi|^{p-2} |h^2| + \dots + \int |h|^p \\ &\leq \frac{1}{\|h\|} \left[\binom{p}{2} \|\varphi\|_{L^p(\mathbf{R})}^{p-2} \|h\|_{L^p(\mathbf{R})}^2 + \dots + \|h\|_{L^p(\mathbf{R})}^p \right] \\ &\leq C(\varphi, p) (\|h\| + \|h\|^2 + \dots + \|h\|^{p-2}) \end{aligned}$$

which $\rightarrow 0$ as $\|h\| \rightarrow 0$.

Hence,

$$Df(\varphi)h = \int p\varphi^{p-1}h$$

Directional Derivatives

We can also define a directional derivative of an operator $f : X \rightarrow Y$.

Gateaux Derivative

Let $h \in X$ be a unit vector. Then $A \in B(X, Y)$ is called the **Gateaux derivative** of f at $x \in X$ in the direction h if

$$\frac{\|f(x + th) - f(x) - tAh\|_Y}{|t|} \rightarrow 0 \quad \text{as } t \rightarrow 0,$$

which is also denoted by

$$A = D_h f(x).$$

Directional Derivatives

We can also define a directional derivative of an operator $f : X \rightarrow Y$.

Gateaux Derivative

Let $h \in X$ be a unit vector. Then $A \in B(X, Y)$ is called the **Gateaux derivative** of f at $x \in X$ in the direction h if

$$\frac{\|f(x + th) - f(x) - tAh\|_Y}{|t|} \rightarrow 0 \quad \text{as } t \rightarrow 0,$$

which is also denoted by

$$A = D_h f(x).$$

Question: if f is Gateaux differentiable (G.d. in every direction), is f also Fréchet differentiable? Conversely?

Directional Derivatives

Fréchet differentiable implies Gateaux differentiable, but not the other way around