

# Math 5601 Final Project

Jacob Hauck

December 4, 2023

Consider the following second-order ODE with Dirichlet boundary conditions:

$$\frac{d}{dx} \left( c(x) \frac{du(x)}{dx} \right) = f(x), \quad a \leq x \leq b, \quad (1)$$

$$u(a) = g_a, \quad u(b) = g_b. \quad (2)$$

## Problem 1.

---

Consider the second-order ODE (1). Multiplying by  $v \in H^1([a, b])$  and integrating by parts gives

$$\int_a^b f v = c(b) u'(b) v(b) - c(a) u'(a) v(a) - \int_a^b c u' v'. \quad (3)$$

(a) Suppose we have the boundary conditions

$$u'(a) = p_a, \quad u(b) = g_b. \quad (4)$$

Equation (3) still holds, and we can impose the condition  $v(b) = 0$  because we already know that  $u(b) = p_b$ . Since  $u'(a) = p_a$ , equation (3) becomes

$$\int_a^b f v = -c(a) p_a v(a) - \int_a^b c u' v' \quad (5)$$

for all  $v \in H^1([a, b])$  such that  $v(b) = 0$ , which is our weak formulation of (1) with the given boundary conditions.

(b) Suppose we have the boundary conditions

$$u'(a) = p_a, \quad u'(b) + q_b u(b) = p_b. \quad (6)$$

Equation (3) still holds. Since  $u'(b) = p_b - q_b u(b)$ , and  $u'(a) = p_a$ , we get

$$\int_a^b f v = c(b) (p_b - q_b u(b)) v(b) - c(a) p_a v(a) - \int_a^b c u' v' \quad (7)$$

for all  $v \in H^1([a, b])$ , which is our weak formulation of (1) with the given boundary conditions.

(c) Suppose we have the boundary conditions

$$u'(a) = p_a, \quad u'(b) = p_b. \quad (8)$$

Equation (3) still holds. Since  $u'(a) = p_a$ , and  $u'(b) = p_b$ , we get

$$\int_a^b f v = c(b) p_b v(b) - c(a) p_a v(a) - \int_a^b c u' v' \quad (9)$$

for all  $v \in H^1([a, b])$ , which is our weak formulation of (1) with the given boundary conditions.

We note that solutions of this formulation are not unique. Indeed, if  $u \in H^1([a, b])$  satisfies (9) for all  $v \in H^1([a, b])$ , then so does  $u + \alpha$ , where  $\alpha \in \mathbf{R}$  is any real number because  $(u + \alpha)' = u'$  regardless of what  $\alpha$  is, and the weak formulation depends only on  $u'$ .

**Problem 2.**

Consider the Poisson equation

$$\nabla \cdot (c \nabla u) = f \text{ in } D. \quad (10)$$

Using integration by parts, we have

$$\int_D f v = \int_D \nabla \cdot (c \nabla u) v = \int_{\partial D} c v \nabla u \cdot n \, dS - \int_D c \nabla u \cdot \nabla v, \quad (11)$$

where  $dS$  is the surface measure on  $\partial D$ , and  $v \in H^1(\overline{D})$ .

(a) Suppose that we have the boundary condition

$$u = g \text{ on } \partial D. \quad (12)$$

Equation (11) still holds. Since we know the value of  $u$  on  $\partial D$ , we can set  $v = 0$  on  $\partial D$ . Then we get

$$\int_D f v = - \int_D c \nabla u \cdot \nabla v \quad (13)$$

for all  $v \in H^1(\overline{D})$  such that  $v = 0$  on  $\partial D$ , which is our weak formulation of (10) with the given boundary condition.

(b) Suppose that we have the boundary condition

$$\nabla u \cdot n + q u = p \text{ on } \partial D, \quad (14)$$

where  $n$  is the outward unit normal vector to  $\partial D$ , and  $p$  and  $q$  are functions on  $\partial D$ . Equation (11) still holds. Since  $\nabla u \cdot n = p - q u$  on  $\partial D$ , it follows that

$$\int_D f v = \int_{\partial D} c v (p - q u) \, dS - \int_D c \nabla u \cdot \nabla v \quad (15)$$

for all  $v \in H^1(\overline{D})$ , which is our weak formulation of (10) with the given boundary condition.

**Problem 3.**

If  $u \in C^2[a, b]$ , then

$$\|u - I_h u\|_\infty \leq \frac{1}{8} h^2 \|u''\|_\infty, \quad (16)$$

$$\|(u - I_h u)'\|_\infty \leq \frac{1}{2} h \|u''\|_\infty. \quad (17)$$

*Proof.* Consider the interval  $[x_i, x_{i+1}]$ , where  $1 \leq i \leq N$ . Restricted to this interval,  $I_h u$  is the degree-1 Lagrange polynomial interpolation of  $u$  on with nodes  $x_i$  and  $x_{i+1}$ . By the error formula for Lagrange polynomial approximation in the slides,

$$u(x) - I_h u(x) = \frac{f''(\xi(x))(x - x_i)(x - x_{i+1})}{2} \quad (18)$$

for some  $\xi(x) \in [x_i, x_{i+1}]$ . Then

$$|u(x) - I_h u(x)| \leq \|f''\|_\infty \cdot \frac{1}{2} (x - x_i)(x_{i+1} - x). \quad (19)$$

The function  $g(x) = (x - x_i)(x_{i+1} - x)$  is a downward-opening parabola, so it achieves maximum halfway between its roots  $x_i$  and  $x_{i+1}$ . Therefore,

$$|u(x) - I_h u(x)| \leq \|f''\|_\infty \cdot \frac{\left(\frac{x_i + x_{i+1}}{2} - x_i\right) \left(x_{i+1} - \frac{x_i + x_{i+1}}{2}\right)}{2} \quad (20)$$

$$= \|f''\|_\infty \frac{(x_{i+1} - x_i)^2}{8} = \frac{h^2}{8} \|f''\|_\infty. \quad (21)$$

Since this holds for all  $x \in [x_i, x_{i+1}]$  and all  $1 \leq i \leq N$ , it holds for all  $x \in [a, b]$ . Therefore, the inequality (16) follows.

Let  $1 \leq i \leq N$ , and let  $x \in [x_i, x_{i+1}]$ . By the Mean Value Theorem, there exists  $c \in [x_i, x_{i+1}]$  such that

$$u'(c) = \frac{u(x_{i+1}) - u(x_i)}{x_{i+1} - x_i} = (I_h u)'(x). \quad (22)$$

Then  $(I_h u)'$  is the degree-0 Lagrange polynomial interpolation of  $u'$  with node  $c$ . By the error formula for Lagrange polynomial approximation in the slides,

$$u'(x) - (I_h u)'(x) = u''(\xi(x))(x - c) \quad (23)$$

□

#### Problem 4.

Consider the weak formulation of

$$\nabla \cdot (c \nabla u) = f \text{ in } D, \quad u = g \text{ on } \partial D \quad (24)$$

derived in problem 2 (a):

$$\int_D f v = - \int_D c \nabla u \cdot \nabla v \quad (25)$$

for all  $v \in H^1(\overline{D})$  such that  $v = 0$  on  $\partial D$ . Suppose that we have basis functions  $\{\phi_i\}_{i=1}^{N+1}$  for a finite element space  $U_h$  on  $\overline{D}$ . To approximate a solution of the weak formulation, we approximate  $H^1$  by  $U_h$ . Thus, we want to find  $u \in U_h$  such that (25) holds for all  $v \in U_h$ .

By the linearity of the problem and the fact that  $U_h = \text{span}\{\phi_i\}$ , this is equivalent to (25) being true for  $v = \phi_i$ , for  $i = 1, \dots, N+1$ . Since we want  $u \in U_h$ , there exist coefficients  $u_j$  such that

$$u = \sum_{j=1}^{N+1} u_j \phi_j. \quad (26)$$

Hence, we need

$$\int_D f \phi_i = - \int_D c \nabla \left( \sum_{j=1}^{N+1} u_j \phi_j \right) \cdot \nabla \phi_i \quad (27)$$

for all  $i = 1, \dots, N+1$ . Using the linearity of  $\nabla$  and rearranging terms, this is equivalent to

$$\sum_{j=1}^{N+1} u_j \left[ - \int_D c \nabla \phi_j \cdot \nabla \phi_i \right] = \int_D f \phi_i \quad (28)$$

for all  $i = 1, \dots, N + 1$ . If we set

$$A_{ij} = - \int_D c \nabla \phi_j \cdot \nabla \phi_i, \quad b_i = \int_D f \phi_i, \quad X_j = u_j, \quad (29)$$

then this is equivalent to the linear system  $AX = b$ .

---

### Problem 5.

Let  $A$  be a nonsingular, lower-triangular matrix; that is,  $i < j$  implies that  $A_{ij} = 0$ . Then  $A^{-1}$  is also lower-triangular.

*Proof.* We use induction on the size of the matrix. All  $1 \times 1$  matrices are trivially lower-triangular, so the base case holds. Now suppose that the claim is true for all matrices of size  $n \times n$ , where  $n \geq 1$ .

Let  $A$  be a nonsingular,  $(n + 1) \times (n + 1)$ , lower-triangular matrix. Then every entry but the last entry of the last column of  $A$  is zero by the lower-triangular condition. That is, we can write  $A$  in block matrix form as

$$A = \begin{bmatrix} B & 0 \\ c & d \end{bmatrix}, \quad (30)$$

where  $B$  is a  $n \times n$  matrix,  $c$  is a  $1 \times n$  row vector, and  $d$  is a scalar. Since  $A_{ij} = B_{ij}$  if  $i, j \leq n$ , it follows that  $B$  is also lower-triangular. Furthermore,  $B$  must be nonsingular.

Indeed, suppose for the sake of contradiction that  $B$  is singular. Then its rows  $\{B_1, \dots, B_n\}$  are linearly dependent. That is, there exist  $\alpha_1, \dots, \alpha_n$  not all zero such that

$$\alpha_1 B_1 + \dots + \alpha_n B_n = 0. \quad (31)$$

Let  $\{A_1, \dots, A_n, A_{n+1}\}$  denote the rows of  $A$ . Then  $A_i = [B_i \ 0]$  for  $1 \leq i \leq n$ . Hence,

$$\alpha_1 A_1 + \dots + \alpha_n A_n = 0 \quad (32)$$

as well. This implies that the rows of  $A$  are linearly dependent, which contradicts the nonsingularity of  $A$ .

Therefore,  $B$  is a nonsingular,  $n \times n$ , lower-triangular matrix, and the induction hypothesis implies that  $B^{-1}$  is lower-triangular.

In addition,  $d \neq 0$  because  $d = 0$  implies that  $\det(A) = 0$  upon expansion by cofactors on the last column of  $A$ , which contradicts the nonsingularity of  $A$ .

We now observe that

$$A \begin{bmatrix} B^{-1} & 0 \\ -cB^{-1}d^{-1} & d^{-1} \end{bmatrix} = \begin{bmatrix} B & 0 \\ c & d \end{bmatrix} \begin{bmatrix} B^{-1} & 0 \\ -cB^{-1}d^{-1} & d^{-1} \end{bmatrix} = \begin{bmatrix} I_{n \times n} & 0 \\ cB^{-1} - cB^{-1}d^{-1}d & 1 \end{bmatrix} = I_{(n+1) \times (n+1)}, \quad (33)$$

so

$$A^{-1} = \begin{bmatrix} B^{-1} & 0 \\ -cB^{-1}d^{-1} & d^{-1} \end{bmatrix}. \quad (34)$$

Then  $A^{-1}$  is lower-triangular because  $B^{-1}$  is lower triangular.  $\square$

---

### Problem 6.

Let

$$A = \begin{bmatrix} \kappa & \lambda \\ \lambda & \mu \end{bmatrix} \quad (35)$$

be a positive definite matrix. Then the Jacobi method for  $Ax = b$  converges.

*Proof.* We recall from the slides that the Jacobi method is the iteration

$$x^{(k+1)} = -D^{-1}Nx^{(k)} + D^{-1}b, \quad (36)$$

where  $D$  is the diagonal of  $A$ , and  $N$  is the off-diagonal of  $A$ . This iteration converges if and only if  $\rho(-D^{-1}N) < 1$ . In this case,

$$-D^{-1}N = -\begin{bmatrix} 0 & \frac{\lambda}{\mu} \\ \frac{\lambda}{\kappa} & 0 \end{bmatrix}, \quad (37)$$

so any eigenvalue  $\rho$  of  $-D^{-1}N$  satisfies  $\rho^2 - \frac{\lambda^2}{\kappa\mu} = 0$ . Therefore  $|\rho| < 1$  if and only if  $\lambda^2 < \kappa\mu$ , or  $\kappa\mu - \lambda^2 > 0$ . Since  $\kappa\mu - \lambda^2 = \det(A)$ , and the positive definiteness of  $A$  implies that  $\det(A) > 0$ , it follows that  $\rho(-D^{-1}N) < 1$ , and the Jacobi method converges.  $\square$

---

**Problem 7.**

---

(a)

(b)

---

**Problem 8.**

---

(a)