

# Exponential Functions

Jacob Hauck

Math 6010

9-20-2023

# Review of Exponents

Let  $a > 0$  and  $m, n$  be integers

$$a^m = \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}}$$

# Review of Exponents

Let  $a > 0$  and  $m, n$  be integers

$$a^m = \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \quad (1.5)^3 = 1.5 \cdot 1.5 \cdot 1.5 = 3.375$$

# Review of Exponents

Let  $a > 0$  and  $m, n$  be integers

$$a^m = \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \quad (1.5)^3 = 1.5 \cdot 1.5 \cdot 1.5 = 3.375$$

►  $a^0 = 1, \quad 1^m = 1, \quad a^1 = a$

# Review of Exponents

Let  $a > 0$  and  $m, n$  be integers

$$a^m = \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \quad (1.5)^3 = 1.5 \cdot 1.5 \cdot 1.5 = 3.375$$

►  $a^0 = 1, \quad 1^m = 1, \quad a^1 = a$

►  $a^m \cdot a^n = a^{m+n}$

# Review of Exponents

Let  $a > 0$  and  $m, n$  be integers

$$a^m = \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \quad (1.5)^3 = 1.5 \cdot 1.5 \cdot 1.5 = 3.375$$

$$\blacktriangleright a^0 = 1, \quad 1^m = 1, \quad a^1 = a$$

$$\blacktriangleright a^m \cdot a^n = a^{m+n} \quad \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \cdot \underbrace{a \cdot a \cdot \dots \cdot a}_{n \text{ times}}$$

# Review of Exponents

Let  $a > 0$  and  $m, n$  be integers

$$a^m = \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \quad (1.5)^3 = 1.5 \cdot 1.5 \cdot 1.5 = 3.375$$

$$\blacktriangleright a^0 = 1, \quad 1^m = 1, \quad a^1 = a$$

$$\blacktriangleright a^m \cdot a^n = a^{m+n} \quad \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \cdot \underbrace{a \cdot a \cdot \dots \cdot a}_{n \text{ times}}$$

$$\blacktriangleright \frac{1}{a^n} = a^{-n}$$

# Review of Exponents

Let  $a > 0$  and  $m, n$  be integers

$$a^m = \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \quad (1.5)^3 = 1.5 \cdot 1.5 \cdot 1.5 = 3.375$$

$$\blacktriangleright a^0 = 1, \quad 1^m = 1, \quad a^1 = a$$

$$\blacktriangleright a^m \cdot a^n = a^{m+n} \quad \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \cdot \underbrace{a \cdot a \cdot \dots \cdot a}_{n \text{ times}}$$

$$\blacktriangleright \frac{1}{a^n} = a^{-n} \quad \text{want } a^{-n} \cdot a^n = a^0 = 1$$



# Review of Exponents

Let  $a > 0$  and  $m, n$  be integers

$$a^m = \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \quad (1.5)^3 = 1.5 \cdot 1.5 \cdot 1.5 = 3.375$$

►  $a^0 = 1, \quad 1^m = 1, \quad a^1 = a$

►  $a^m \cdot a^n = a^{m+n} \quad \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \cdot \underbrace{a \cdot a \cdot \dots \cdot a}_{n \text{ times}}$

►  $\frac{1}{a^n} = a^{-n} \quad \text{want } a^{-n} \cdot a^n = a^0 = 1$

►  $\frac{a^m}{a^n} = a^{m-n}$

# Review of Exponents

Let  $a > 0$  and  $m, n$  be integers

$$a^m = \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \quad (1.5)^3 = 1.5 \cdot 1.5 \cdot 1.5 = 3.375$$

$$\blacktriangleright a^0 = 1, \quad 1^m = 1, \quad a^1 = a$$

$$\blacktriangleright a^m \cdot a^n = a^{m+n} \quad \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \cdot \underbrace{a \cdot a \cdot \dots \cdot a}_{n \text{ times}}$$

$$\blacktriangleright \frac{1}{a^n} = a^{-n} \quad \text{want } a^{-n} \cdot a^n = a^0 = 1$$

$$\blacktriangleright \frac{a^m}{a^n} = a^{m-n}$$

$$\blacktriangleright (a^n)^m = a^{m \cdot n} \quad \underbrace{a^n \cdot a^n \cdot \dots \cdot a^n}_{m \text{ times}}$$

# Review of Exponents

Let  $a > 0$  and  $m, n$  be integers

$$a^m = \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \quad (1.5)^3 = 1.5 \cdot 1.5 \cdot 1.5 = 3.375$$

$$\blacktriangleright a^0 = 1, \quad 1^m = 1, \quad a^1 = a$$

$$\blacktriangleright a^m \cdot a^n = a^{m+n} \quad \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \cdot \underbrace{a \cdot a \cdot \dots \cdot a}_{n \text{ times}}$$

$$\blacktriangleright \frac{1}{a^n} = a^{-n} \quad \text{want } a^{-n} \cdot a^n = a^0 = 1$$

$$\blacktriangleright \frac{a^m}{a^n} = a^{m-n}$$

$$\blacktriangleright (a^n)^m = a^{m \cdot n} \quad \underbrace{a^n \cdot a^n \cdot \dots \cdot a^n}_{m \text{ times}}$$

$$\blacktriangleright a^{\frac{1}{n}} = \sqrt[n]{a} \quad \text{want } \left(a^{\frac{1}{n}}\right)^n = a^1 = a$$

# Review of Exponents

Let  $a > 0$  and  $m, n$  be integers

$$a^m = \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \quad (1.5)^3 = 1.5 \cdot 1.5 \cdot 1.5 = 3.375$$

$$\blacktriangleright a^0 = 1, \quad 1^m = 1, \quad a^1 = a$$

$$\blacktriangleright a^m \cdot a^n = a^{m+n} \quad \underbrace{a \cdot a \cdot \dots \cdot a}_{m \text{ times}} \cdot \underbrace{a \cdot a \cdot \dots \cdot a}_{n \text{ times}}$$

$$\blacktriangleright \frac{1}{a^n} = a^{-n} \quad \text{want } a^{-n} \cdot a^n = a^0 = 1$$

$$\blacktriangleright \frac{a^m}{a^n} = a^{m-n}$$

$$\blacktriangleright (a^n)^m = a^{m \cdot n} \quad \underbrace{a^n \cdot a^n \cdot \dots \cdot a^n}_{m \text{ times}}$$

$$\blacktriangleright a^{\frac{1}{n}} = \sqrt[n]{a} \quad \text{want } \left(a^{\frac{1}{n}}\right)^n = a^1 = a$$

$$\blacktriangleright a^{\frac{m}{n}} = \sqrt[n]{a^m}$$

# Exponential Functions

Want a function like

$$f(x) = C \cdot a^x$$

# Exponential Functions

Want a function like

$$f(x) = C \cdot a^x$$

but  $x$  might not be  $\frac{m}{n}$ , for integers  $m, n \dots$

# Exponential Functions

Want a function like

$$f(x) = C \cdot a^x$$

but  $x$  might not be  $\frac{m}{n}$ , for integers  $m, n \dots$

►  $x = \pi, \sqrt{2}$ , and so on

# Exponential Functions

Want a function like

$$f(x) = C \cdot a^x$$

but  $x$  might not be  $\frac{m}{n}$ , for integers  $m, n$ ...

►  $x = \pi, \sqrt{2}$ , and so on

...but *every* number has a decimal expansion

$$3 = 3, \quad \frac{2}{5} = 0.4, \quad \pi = 3.14159265\dots, \quad \sqrt{2} = 1.41421356\dots$$



# Exponential Functions

Want a function like

$$f(x) = C \cdot a^x$$

but  $x$  might not be  $\frac{m}{n}$ , for integers  $m, n$ ...

►  $x = \pi, \sqrt{2}$ , and so on

...but *every* number has a decimal expansion

$$3 = 3, \quad \frac{2}{5} = 0.4, \quad \pi = 3.14159265\dots, \quad \sqrt{2} = 1.41421356\dots$$

Approximate any number  $x$  by finitely many of its decimal digits

$$a^\pi \approx a^{3.14} = a^{\frac{314}{100}} = \sqrt[100]{a^{314}} \quad (1)$$

# Exponential Functions

The more digits we use, the closer we get to the true value of  $a^\pi$

$2^3$	$2^{3.1}$	$2^{3.14}$	$2^{3.141}$	$2^{3.1415}$	$\dots$	$2^\pi$
8.0000	8.5742	8.8152	8.8213	8.8244	$\dots$	8.8250

# Exponential Functions

The more digits we use, the closer we get to the true value of  $a^\pi$

$2^3$	$2^{3.1}$	$2^{3.14}$	$2^{3.141}$	$2^{3.1415}$	$\dots$	$2^\pi$
8.0000	8.5742	8.8152	8.8213	8.8244	$\dots$	8.8250

So we can use any number  $x$  in the exponent, and we also get to keep all the previous properties. Hooray!

# Exponential Functions

The more digits we use, the closer we get to the true value of  $a^\pi$

$2^3$	$2^{3.1}$	$2^{3.14}$	$2^{3.141}$	$2^{3.1415}$	$\dots$	$2^\pi$
8.0000	8.5742	8.8152	8.8213	8.8244	$\dots$	8.8250

So we can use any number  $x$  in the exponent, and we also get to keep all the previous properties. Hooray!

## Exponential Functions

An **exponential function** is a function  $f$  such that

$$f(x) = C \cdot a^x$$

- ▶  $C \neq 0$  is the **initial value**
- ▶  $a > 0$ ,  $a \neq 1$  is the **growth factor**

# Exponential Functions

Consider the exponential function  $f(x) = 5 \cdot 2^x$ . Let's make a table of values

$x$	$f(x)$
-2	1.25
-1	2.5
0	5
1	10
2	20

# Exponential Functions

Consider the exponential function  $f(x) = 5 \cdot 2^x$ . Let's make a table of values

$x$	$f(x)$
-2	1.25
-1	2.5
0	5
1	10
2	20

- $f(x)$  goes up by  $\times 2 = a$  each time  $x$  increases

# Exponential Functions

Consider the exponential function  $f(x) = 5 \cdot 2^x$ . Let's make a table of values

$x$	$f(x)$
-2	1.25
-1	2.5
0	5
1	10
2	20

- ▶  $f(x)$  goes up by  $\times 2 = a$  each time  $x$  increases
- ▶  $f(0) = 5 = C$ .

# Initial Value and Growth Factor

$$f(x) = C \cdot a^x$$

- ▶  $f(0) = C \cdot a^0 = C$ , the initial value



# Initial Value and Growth Factor

$$f(x) = C \cdot a^x$$

- ▶  $f(0) = C \cdot a^0 = C$ , the initial value
- ▶ The growth factor: how much the function grows (or shrinks) every time  $x$  goes up by 1

$$f(x+1) = a \cdot f(x)$$

# Initial Value and Growth Factor

$$f(x) = C \cdot a^x$$

- ▶  $f(0) = C \cdot a^0 = C$ , the initial value
- ▶ The growth factor: how much the function grows (or shrinks) every time  $x$  goes up by 1

$$f(x+1) = a \cdot f(x)$$

because

$$\frac{f(x+1)}{f(x)} = \frac{C \cdot a^{x+1}}{C \cdot a^x}$$

# Initial Value and Growth Factor

$$f(x) = C \cdot a^x$$

- ▶  $f(0) = C \cdot a^0 = C$ , the initial value
- ▶ The growth factor: how much the function grows (or shrinks) every time  $x$  goes up by 1

$$f(x+1) = a \cdot f(x)$$

because

$$\frac{f(x+1)}{f(x)} = \frac{C \cdot a^{x+1}}{C \cdot a^x} = \frac{a^{x+1}}{a^x}$$

# Initial Value and Growth Factor

$$f(x) = C \cdot a^x$$

- ▶  $f(0) = C \cdot a^0 = C$ , the initial value
- ▶ The growth factor: how much the function grows (or shrinks) every time  $x$  goes up by 1

$$f(x+1) = a \cdot f(x)$$

because

$$\frac{f(x+1)}{f(x)} = \frac{C \cdot a^{x+1}}{C \cdot a^x} = \frac{a^{x+1}}{a^x} = a^{x+1-x} = a$$

# Graphing Exponential Functions

Let's try to graph the function  $f(x) = 5 \cdot 2^x$  from before. First, extend the table of values

# Graphing Exponential Functions

Let's try to graph the function  $f(x) = 5 \cdot 2^x$  from before. First, extend the table of values

$x$	$f(x)$
-5	0.16125
-4	0.3125
-3	0.625
-2	1.25
-1	2.5
0	5
1	10
2	20
3	40
4	80
5	160

# Graphing Exponential Functions

Let's try to graph the function  $f(x) = 5 \cdot 2^x$  from before. First, extend the table of values

$x$	$f(x)$
-5	0.16125
-4	0.3125
-3	0.625
-2	1.25
-1	2.5
0	5
1	10
2	20
3	40
4	80
5	160

It seems that  $f(x) \rightarrow 0$  as  $x \rightarrow -\infty$ , and  $f(x) \rightarrow \infty$  as  $x \rightarrow \infty$

# Graphing Exponential Functions

Connect the points continuously and use the asymptotic behavior noted

