



Atlantis Course Universidad de Concepción

Day 2 - Key modelling decisions


Javier Porobic and Beth Fulton

CSIRO Ocean & Atmosphere
www.csiro.au



Introduction

- General model design
 - Define your system
 - Define functional groups
 - Map and polygon structure
- State conditions (CDF file)
 - Biomass distribution
- Rate parameters (PRM file)
 - Diets
 - Species movement (within / outside model domain)
 - Recruitment
- Tips and tricks



Complexity tradeoffs

- Spatial
- Temporal
- Processes details (physical, chemical, ecological, anthropogenic)
- Trophic components (taxonomic resolution)
- Human industries
- Stock sub-structure
- Movement

Model complexity: best practices

Attack problems at appropriate scales, with appropriate theories and methods.
A universal scale is NOT useful

Rudolph Marcus

FAO 2007


- recommendation given for each feature
- only as much as needed for critical dynamics
- feedbacks critical
- uncertainty means try multiple options

Define your ecosystem


- Depends on the questions you want to answer
- If possible, choose the area to be contained in a natural oceanographic or biogeographic feature, with uniform climate & current system
- Larger area is better with many migratory species
- Mortality is dealt with explicitly in model dynamics rather than assumed and static
- Migrators / transients don't respect regional jurisdictions



Define your period



Spin-up method




Define your functional groups

- Good reasons to aggregate functional groups
- Ignorance
 - Biomass, diet composition, functional responses within groups may be poorly understood or data unavailable
- Computational convenience
 - Run-time prohibits species-level representation
- Heuristic advantage
 - Too many functional groups makes it difficult to track through interactions and extract causes and consequences




Define your functional groups : Rules of thumb

- Consider stock structure
- Diet, habitat preference, vulnerability to fishing, fishing interest, conservation interest are most important
- Don't aggregate if they have disparate diets, turn-over rates, or trophic levels
- Target resource species individually (**it will pay off**)



Define your functional groups **Methods**


- Subjective / Intuition
- Rule-based
 - Diet
 - Habitat
 - Vulnerability
- Multivariate statistics
 - Cluster analysis¹
 - Factor analysis²
 - Multidimensional scaling



MDS of stomach content data Gulf of California


How many groups to use? Prescription VS Prediction

- More detail can be prescriptive not predictive
- Large models with lots of noisy data vs slim-line model with few precise data
- Detail vs. ease of parameterization
- Complex systems / complex models



Map design : Some useful spatial data


- Bathymetry & coastlines (can read it in directly)
- Habitat type / substrate composition
- rough, flat, soft, canyon & biogenic
- Locations of new & existing marine reserves
- Ancillary information (human population, port locations)
- Fishing areas / depth



These data will be integrated to build your map based on physical oceanography, biogeography, human activities and data availability

Map design : Polygonal geometry

- Polygon matching physical and ecological features (e.g. bioregionalisation)
- Homogeneous areas = large cells, heterogeneous areas = small cells
- Not so easy to declare but may be more meaningful (detail where needed)
- Can conform to biogeographic features, data reporting regions
- Computationally efficient



Same issues vertically as horizontally

Map design : **spatial resolution**




- Lower variance shown through time and space
- Less heterogeneous environment and fewer refugia
- Increased self-simplification of trophic web
 - (extinction and exclusion – no spatial refuge)
- Loss of spatial segregation of predator/prey
- Increased boundary condition effects
- Altered predictions of response to perturbation

True for all ecosystem models

State conditions

- Initial biomass distribution:
 - Important for spatial segregation of predator and prey
 - Adult locations important for recruitment
- Equilibrium position
 - Initial structure
 - Life-span



Ontogenetic biomass distribution


Choosing a survival-at-age model

- The simplest model for predicting numbers at age.

$$N_t = N_{t-1}e^{-z}$$

- Not appropriate for mammals / birds
- Siler's competing risk model (more generic)

$$S_t = S_Juv_{juvenile,t} \cdot S_Mat_t \cdot S_Senescence_t$$



Thank You

Ocean & Atmosphere

Javier Porobic

t +61362325037

e javier.porobicgarate@csiro.au

w Javier Porobic

CSIRO Ocean & Atmosphere
www.csiro.au

