

# Polarized and Directed Type Theory

Jacob Neumann  
25 April 2025  
Reykholt, Iceland



**0 Background**

Standard Martin-Löf Type Theory includes **identity types**:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash t : A \quad \Gamma \vdash t' : A}{\Gamma \vdash \text{Id}_A(t, t') \text{ type}}$$

Standard Martin-Löf Type Theory includes **identity types**:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash t : A \quad \Gamma \vdash t' : A}{\Gamma \vdash \text{Id}_A(t, t') \text{ type}}$$

$$\frac{\Gamma \text{ ctx} \quad \Gamma \vdash t : A}{\Gamma \vdash \text{refl}_t : \text{Id}(t, t)}$$

Standard Martin-Löf Type Theory includes **identity types**:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash t : A \quad \Gamma \vdash t' : A}{\Gamma \vdash \text{Id}_A(t, t') \text{ type}}$$

$$\frac{\begin{array}{c} \Gamma \vdash t : A \\ \Gamma, x : A, u : \text{Id}(t, x) \vdash M(x, u) \text{ type} \\ \Gamma \vdash m : M(t, \text{refl}) \end{array}}{\Gamma, x : A, u : \text{Id}(t, x) \vdash J \, m \, x \, u : M(x, u)}$$
$$\frac{\Gamma \text{ ctx} \quad \Gamma \vdash t : A}{\Gamma \vdash \text{refl}_t : \text{Id}(t, t)}$$



## Symmetry

$$\frac{\Gamma \vdash p : \text{Id}(t, t')}{\Gamma \vdash p^{-1} := J \text{ refl}_t t' p : \text{Id}(t', t)}$$

## Symmetry

$$\frac{\Gamma \vdash p : \text{Id}(t, t')}{\Gamma \vdash p^{-1} := J \text{ refl}_t t' p : \text{Id}(t', t)}$$

## Transitivity

$$\frac{\begin{array}{l} \Gamma \vdash p : \text{Id}(t, t') \\ \Gamma \vdash q : \text{Id}(t', t'') \end{array}}{\Gamma \vdash p \cdot q := J p t'' q : \text{Id}(t, t')}$$



## Symmetry

$$\frac{\Gamma \vdash p : \text{Id}(t, t')}{\Gamma \vdash p^{-1} := J \text{ refl}_t t' p : \text{Id}(t', t)}$$

## Transitivity

$$\frac{\begin{array}{l} \Gamma \vdash p : \text{Id}(t, t') \\ \Gamma \vdash q : \text{Id}(t', t'') \end{array}}{\Gamma \vdash p \cdot q := J p t'' q : \text{Id}(t, t')}$$

- Assuming the **uniqueness of identity proofs** (if  $p, p' : \text{Id}(t, t')$ , there's some  $\text{UIP}(p, p') : \text{Id}(p, p')$ ), this makes  $A$  into an **equivalence relation**.

## Symmetry

$$\frac{\Gamma \vdash p : \text{Id}(t, t')}{\Gamma \vdash p^{-1} := J \text{ refl}_t t' p : \text{Id}(t', t)}$$

## Transitivity

$$\frac{\begin{array}{l} \Gamma \vdash p : \text{Id}(t, t') \\ \Gamma \vdash q : \text{Id}(t', t'') \end{array}}{\Gamma \vdash p \cdot q := J p t'' q : \text{Id}(t, t')}$$

- Assuming the **uniqueness of identity proofs** (if  $p, p' : \text{Id}(t, t')$ , there's some  $\text{UIP}(p, p') : \text{Id}(p, p')$ ), this makes  $A$  into an **equivalence relation**.
- Assuming UIP for the *identity types of*  $A$ , this makes  $A$  into a **synthetic groupoid** (terms of  $A$  are objects, Id-terms are morphisms,  $\text{refl}_t$  is the identity morphism at  $t$ )

## Inverses

$$\frac{\Gamma \vdash p : \text{Id}(t, t')}{\Gamma \vdash p^{-1} := J \text{ refl}_t t' p : \text{Id}(t', t)}$$

## Transitivity

$$\frac{\begin{array}{l} \Gamma \vdash p : \text{Id}(t, t') \\ \Gamma \vdash q : \text{Id}(t', t'') \end{array}}{\Gamma \vdash p \cdot q := J p t'' q : \text{Id}(t, t')}$$

- Assuming the **uniqueness of identity proofs** (if  $p, p' : \text{Id}(t, t')$ , there's some  $\text{UIP}(p, p') : \text{Id}(p, p')$ ), this makes  $A$  into an **equivalence relation**.
- Assuming UIP for the *identity types of*  $A$ , this makes  $A$  into a **synthetic groupoid** (terms of  $A$  are objects, Id-terms are morphisms,  $\text{refl}_t$  is the identity morphism at  $t$ )

## Inverses

$$\frac{\Gamma \vdash p : \text{Id}(t, t')}{\Gamma \vdash p^{-1} := J \text{ refl}_t t' p : \text{Id}(t', t)}$$

## Composition

$$\frac{\begin{array}{l} \Gamma \vdash p : \text{Id}(t, t') \\ \Gamma \vdash q : \text{Id}(t', t'') \end{array}}{\Gamma \vdash p \cdot q := J p t'' q : \text{Id}(t, t')}$$

- Assuming the **uniqueness of identity proofs** (if  $p, p' : \text{Id}(t, t')$ , there's some  $\text{UIP}(p, p') : \text{Id}(p, p')$ ), this makes  $A$  into an **equivalence relation**.
- Assuming UIP for the *identity types of*  $A$ , this makes  $A$  into a **synthetic groupoid** (terms of  $A$  are objects, Id-terms are morphisms,  $\text{refl}_t$  is the identity morphism at  $t$ )

**Question:** Can we  
have a synthetic  
*category* theory?

Standard Martin-Löf Type Theory includes **identity types**:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash t : A \quad \Gamma \vdash t' : A}{\Gamma \vdash \text{Id}_A(t, t') \text{ type}}$$

$$\frac{\begin{array}{c} \Gamma \vdash t : A \\ \Gamma, x : A, u : \text{Id}(t, x) \vdash M(x, u) \text{ type} \\ \Gamma \vdash m : M(t, \text{refl}) \end{array}}{\Gamma, x : A, u : \text{Id}(t, x) \vdash J \ m \ x \ u : M(x, u)}$$
$$\frac{\Gamma \text{ ctx} \quad \Gamma \vdash t : A}{\Gamma \vdash \text{refl}_t : \text{Id}(t, t)}$$

**Directed Type Theory** includes **identity types**:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash t : A \quad \Gamma \vdash t' : A}{\Gamma \vdash \text{Id}_A(t, t') \text{ type}}$$

$$\frac{\begin{array}{c} \Gamma \vdash t : A \\ \Gamma, x : A, u : \text{Id}(t, x) \vdash M(x, u) \text{ type} \\ \Gamma \vdash m : M(t, \text{refl}) \end{array}}{\Gamma, x : A, u : \text{Id}(t, x) \vdash J \ m \ x \ u : M(x, u)}$$
$$\frac{\Gamma \text{ ctx} \quad \Gamma \vdash t : A}{\Gamma \vdash \text{refl}_t : \text{Id}(t, t)}$$

**Directed Type Theory** includes **hom-types**:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash t : A \quad \Gamma \vdash t' : A}{\Gamma \vdash \text{Id}_A(t, t') \text{ type}}$$

$$\frac{\begin{array}{c} \Gamma \vdash t : A \\ \Gamma, x : A, u : \text{Id}(t, x) \vdash M(x, u) \text{ type} \\ \Gamma \vdash m : M(t, \text{refl}) \end{array}}{\Gamma, x : A, u : \text{Id}(t, x) \vdash J \ m \ x \ u : M(x, u)}$$

$$\frac{\Gamma \text{ ctx} \quad \Gamma \vdash t : A}{\Gamma \vdash \text{refl}_t : \text{Id}(t, t)}$$



**Directed Type Theory** includes **hom-types**:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash t : A \quad \Gamma \vdash t' : A}{\Gamma \vdash \text{Hom}_{\mathbf{A}}(t, t') \text{ type}}$$

$$\Gamma \vdash t : A$$

$$\Gamma, x : A, u : \text{Hom}(t, x) \vdash M(x, u) \text{ type}$$

$$\Gamma \vdash m : M(t, \text{refl})$$

$$\frac{\Gamma \text{ ctx} \quad \Gamma \vdash t : A}{\Gamma \vdash \text{refl}_t : \text{Hom}(t, t)}$$

$$\frac{\Gamma \vdash m : M(t, \text{refl})}{\Gamma, x : A, u : \text{Hom}(t, x) \vdash J \ m \ x \ u : M(x, u)}$$

## Inverses

$$\frac{\Gamma \vdash p: \text{Hom}(t, t')}{\Gamma \vdash J \text{ refl}_t t' p: \text{Hom}(t', t)}$$

## Composition

$$\frac{\begin{array}{l} \Gamma \vdash p: \text{Hom}(t, t') \\ \Gamma \vdash q: \text{Hom}(t', t'') \end{array}}{\Gamma \vdash J p t'' q: \text{Hom}(t, t')}$$


## Inverses

$$\frac{\Gamma \vdash p: \text{Hom}(t, t')}{\Gamma \vdash J \text{ refl}_t t' p: \text{Hom}(t', t)}$$


## Composition

$$\frac{\begin{array}{l} \Gamma \vdash p: \text{Hom}(t, t') \\ \Gamma \vdash q: \text{Hom}(t', t'') \end{array}}{\Gamma \vdash J p t'' q: \text{Hom}(t, t'')} \quad \checkmark$$

## Inverses

$$\frac{\Gamma \vdash p: \text{Hom}(t, t')}{\Gamma \vdash J \text{ refl}_t t' p: \text{Hom}(t', t)}$$


## Composition

$$\frac{\begin{array}{l} \Gamma \vdash p: \text{Hom}(t, t') \\ \Gamma \vdash q: \text{Hom}(t', t'') \end{array}}{\Gamma \vdash J p t'' q: \text{Hom}(t, t'')}$$


**Key Idea:** Need  
*semantics* to prove  
that symmetry is  
unprovable

## Inverses

$$\frac{\Gamma \vdash p : \text{Id}(t, t')}{\Gamma \vdash p^{-1} := J \text{ refl}_t t' p : \text{Id}(t', t)}$$

## Composition

$$\frac{\begin{array}{l} \Gamma \vdash p : \text{Id}(t, t') \\ \Gamma \vdash q : \text{Id}(t', t'') \end{array}}{\Gamma \vdash p \cdot q := J p t'' q : \text{Id}(t, t'')}$$

- Assuming the **uniqueness of identity proofs** (if  $p, p' : \text{Id}(t, t')$ , there's some  $\text{UIP}(p, p') : \text{Id}(p, p')$ ), this makes  $A$  into an **equivalence relation**.

## Inverses

$$\frac{\Gamma \vdash p : \text{Id}(t, t')}{\Gamma \vdash p^{-1} := J \text{ refl}_t t' p : \text{Id}(t', t)}$$

## Composition

$$\frac{\begin{array}{l} \Gamma \vdash p : \text{Id}(t, t') \\ \Gamma \vdash q : \text{Id}(t', t'') \end{array}}{\Gamma \vdash p \cdot q := J p t'' q : \text{Id}(t, t')}$$

- Assuming the **uniqueness of identity proofs** (if  $p, p' : \text{Id}(t, t')$ , there's some  $\text{UIP}(p, p') : \text{Id}(p, p')$ )—**which can't be proved in general from the rules of MLTT**—, this makes  $A$  into an **equivalence relation**.

Hofmann and Streicher prove [HS95] that UIP is *independent from the rules of Martin-Löf Type Theory*.



Hofmann and Streicher prove [HS95] that UIP is *independent from the rules of Martin-Löf Type Theory*.

The **groupoid model** of type theory is an interpretation of MLTT as a mathematical structure:

Hofmann and Streicher prove [HS95] that UIP is *independent from the rules of Martin-Löf Type Theory*.

The **groupoid model** of type theory is an interpretation of MLTT as a mathematical structure:

- The contexts are groupoids

Hofmann and Streicher prove [HS95] that UIP is *independent from the rules of Martin-Löf Type Theory*.

The **groupoid model** of type theory is an interpretation of MLTT as a mathematical structure:

- The contexts are groupoids
- A type in context  $\Gamma$  is a  $\Gamma$ -indexed family of groupoids, i.e. a functor  $\Gamma \rightarrow \mathbf{Grpd}$

Hofmann and Streicher prove [HS95] that UIP is *independent from the rules of Martin-Löf Type Theory*.

The **groupoid model** of type theory is an interpretation of MLTT as a mathematical structure:

- The contexts are groupoids
- A type in context  $\Gamma$  is a  $\Gamma$ -indexed family of groupoids, i.e. a functor  $\Gamma \rightarrow \mathbf{Grpd}$
- A term  $t$  (in context  $\Gamma$ ) of type  $A$  is a section sending objects  $\gamma: |\Gamma|$  to objects  $t \gamma: |A(\gamma)|$  and morphisms  $\gamma_{01}: \Gamma [\gamma_0, \gamma_1]$  to morphisms in  $(A \gamma_1) [A \gamma_{01} (t \gamma_0), t \gamma_1]$ .

$$\text{Id}(t, t') : \Gamma \rightarrow \mathbf{Grpd}$$

$$\text{Id}(t, t') \gamma = (A \gamma) [t \gamma, t' \gamma]$$

$$\text{Id}(t, t') \gamma_{01} x_0 = (t' \gamma_{01}) \circ A \gamma_{01} x_0 \circ (t \gamma_{01})^{-1}$$

$$\begin{array}{ccc}
 A \gamma_{01} (t \gamma_0) & \xrightarrow{A \gamma_{01} x_0} & A \gamma_{01} (t' \gamma_0) \\
 \downarrow t \gamma_{01} & & \downarrow t' \gamma_{01} \\
 t \gamma_1 & & t' \gamma_1
 \end{array}$$

$$\text{Id}(t, t') : \Gamma \rightarrow \mathbf{Grpd}$$

$$\text{Id}(t, t') \gamma = (A \gamma) [t \gamma, t' \gamma]$$

$$\text{Id}(t, t') \gamma_{01} x_0 = (t' \gamma_{01}) \circ A \gamma_{01} x_0 \circ (t \gamma_{01})^{-1}$$

*This type has multiple terms in general, hence UIP cannot be proved from the rules.*

**Idea:** Replace  
groupoids with  
categories

The **groupoid model** of type theory is an interpretation of MLTT as a mathematical structure:

- The contexts are groupoids
- A type in context  $\Gamma$  is a  $\Gamma$ -indexed family of groupoids, i.e. a functor  $\Gamma \rightarrow \mathbf{Grpd}$
- A term  $t$  (in context  $\Gamma$ ) of type  $A$  is a section sending objects  $\gamma: |\Gamma|$  to objects  $t \gamma: |A(\gamma)|$  and morphisms  $\gamma_{01}: \Gamma [\gamma_0, \gamma_1]$  to morphisms in  $(A \gamma_1) [A \gamma_{01} (t \gamma_0), t \gamma_1]$ .



The **category model** of type theory is an interpretation of MLTT as a mathematical structure:

- The contexts are groupoids
- A type in context  $\Gamma$  is a  $\Gamma$ -indexed family of groupoids, i.e. a functor  $\Gamma \rightarrow \mathbf{Grpd}$
- A term  $t$  (in context  $\Gamma$ ) of type  $A$  is a section sending objects  $\gamma: |\Gamma|$  to objects  $t \gamma: |A(\gamma)|$  and morphisms  $\gamma_{01}: \Gamma [\gamma_0, \gamma_1]$  to morphisms in  $(A \gamma_1) [A \gamma_{01} (t \gamma_0), t \gamma_1]$ .

The **category model** of type theory is an interpretation of **directedTT** as a mathematical structure:

- The contexts are groupoids
- A type in context  $\Gamma$  is a  $\Gamma$ -indexed family of groupoids, i.e. a functor  $\Gamma \rightarrow \mathbf{Grpd}$
- A term  $t$  (in context  $\Gamma$ ) of type  $A$  is a section sending objects  $\gamma: |\Gamma|$  to objects  $t \gamma: |A(\gamma)|$  and morphisms  $\gamma_{01}: \Gamma [\gamma_0, \gamma_1]$  to morphisms in  $(A \gamma_1) [A \gamma_{01} (t \gamma_0), t \gamma_1]$ .

The **category model** of type theory is an interpretation of **directedTT** as a mathematical structure:

- The contexts are **categories**
- A type in context  $\Gamma$  is a  $\Gamma$ -indexed family of groupoids, i.e. a functor  $\Gamma \rightarrow \mathbf{Grpd}$
- A term  $t$  (in context  $\Gamma$ ) of type  $A$  is a section sending objects  $\gamma: |\Gamma|$  to objects  $t \gamma: |A(\gamma)|$  and morphisms  $\gamma_{01}: \Gamma [\gamma_0, \gamma_1]$  to morphisms in  $(A \gamma_1) [A \gamma_{01} (t \gamma_0), t \gamma_1]$ .

The **category model** of type theory is an interpretation of **directedTT** as a mathematical structure:

- The contexts are **categories**
- A type in context  $\Gamma$  is a  $\Gamma$ -indexed family of **categories**, i.e. a functor  $\Gamma \rightarrow \mathbf{Cat}$
- A term  $t$  (in context  $\Gamma$ ) of type  $A$  is a section sending objects  $\gamma: |\Gamma|$  to objects  $t \gamma: |A(\gamma)|$  and morphisms  $\gamma_{01}: \Gamma [\gamma_0, \gamma_1]$  to morphisms in  $(A \gamma_1) [A \gamma_{01} (t \gamma_0), t \gamma_1]$ .


$$\text{Id}(t, t') : \Gamma \rightarrow \mathbf{Grpd}$$

$$\text{Id}(t, t') \gamma = (A \gamma) [t \gamma, t' \gamma]$$


$$\text{Id}(t, t') \gamma_{01} x_0 = (t' \gamma_{01}) \circ A \gamma_{01} x_0 \circ (t \gamma_{01})^{-1}$$

$$\begin{array}{ccc}
 A \gamma_{01} (t \gamma_0) & \xrightarrow{A \gamma_{01} x_0} & A \gamma_{01} (t' \gamma_0) \\
 \downarrow t \gamma_{01} & & \downarrow t' \gamma_{01} \\
 t \gamma_1 & & t' \gamma_1
 \end{array}$$

## Inverses

$$\frac{\Gamma \vdash p: \text{Hom}(t, t')}{\Gamma \vdash J \text{ refl}_t t' p: \text{Hom}(t', t)}$$


## Composition

$$\frac{\begin{array}{l} \Gamma \vdash p: \text{Hom}(t, t') \\ \Gamma \vdash q: \text{Hom}(t', t'') \end{array}}{\Gamma \vdash J p t'' q: \text{Hom}(t, t'')}$$


# 1 Polarity Calculus

It is well-known that **Cat** does not provide semantics for dependent type theory (e.g. it is not a LCCC), so we have a choice:



It is well-known that **Cat** does not provide semantics for dependent type theory (e.g. it is not a LCCC), so we have a choice:

- Give up having semantics in **Cat**, and work in a more suitable setting (e.g. simplicial sets, bicubical sets);

It is well-known that **Cat** does not provide semantics for dependent type theory (e.g. it is not a LCCC), so we have a choice:

- Give up having semantics in **Cat**, and work in a more suitable setting (e.g. simplicial sets, bicubical sets);
  - ▶ This is the approach taken by [RS17] and subsequent works [Wei22b, Wei24b, Wei22a, Wei24a, GWB24, GWB25, KRW23]

It is well-known that **Cat** does not provide semantics for dependent type theory (e.g. it is not a LCCC), so we have a choice:

- Give up having semantics in **Cat**, and work in a more suitable setting (e.g. simplicial sets, bicubical sets);
  - ▶ This is the approach taken by [RS17] and subsequent works [Wei22b, Wei24b, Wei22a, Wei24a, GWB24, GWB25, KRW23]
- OR: modify our type theory (by adding a type discipline of *polarities*) to still have semantics in **Cat**

It is well-known that **Cat** does not provide semantics for dependent type theory (e.g. it is not a LCCC), so we have a choice:

- Give up having semantics in **Cat**, and work in a more suitable setting (e.g. simplicial sets, bicubical sets);
  - ▶ This is the approach taken by [RS17] and subsequent works [Wei22b, Wei24b, Wei22a, Wei24a, GWB24, GWB25, KRW23]
- OR: modify our type theory (by adding a type discipline of *polarities*) to still have semantics in **Cat**
  - ▶ What we do [NA24, Neu25], following [LH11, Nuy15, Nor19]

It is well-known that **Cat** does not provide semantics for dependent type theory (e.g. it is not a LCCC), so we have a choice:

- Give up having semantics in **Cat**, and work in a more suitable setting (e.g. simplicial sets, bicubical sets);
  - ▶ This is the approach taken by [RS17] and subsequent works [Wei22b, Wei24b, Wei22a, Wei24a, GWB24, GWB25, KRW23]
- OR: modify our type theory (by adding a type discipline of *polarities*) to still have semantics in **Cat**
  - ▶ What we do [NA24, Neu25], following [LH11, Nuy15, Nor19]
- OR: a little of both
  - ▶ E.g. virtual double category approaches [NL23, Nas25, Nas24]

We annotate **variances/polarities** by adding operations to MLTT

We annotate **variances/polarities** by adding operations to MLTT

- **Type negation**

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash A^- \text{ type}}$$

We annotate **variances/polarities** by adding operations to MLTT

- **Type negation**

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash A^- \text{ type}} \quad \overline{(A^-)^-} = A$$



We annotate **variances/polarities** by adding operations to MLTT

- **Type negation**

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash A^- \text{ type}} \quad \overline{(A^-)^- = A}$$

$$\Gamma \xrightarrow{A} \mathbf{Cat} \xrightarrow{(\_)^\text{op}} \mathbf{Cat}$$

We annotate **variances/polarities** by adding operations to MLTT

- **Type negation**

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash A^- \text{ type}} \quad \overline{(A^-)^- = A}$$

- **Context negation**

$$\frac{\Gamma \text{ ctx}}{\Gamma^- \text{ ctx}}$$

We annotate **variances/polarities** by adding operations to MLTT

- **Type negation**

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash A^- \text{ type}} \quad \overline{(A^-)^-} = A$$

- **Context negation**

$$\frac{\Gamma \text{ ctx}}{\Gamma^- \text{ ctx}} \quad \overline{(\Gamma^-)^-} = \Gamma$$

We annotate **variances/polarities** by adding operations to MLTT

- **Type negation**

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash A^- \text{ type}} \quad \overline{(A^-)^-} = A$$

- **Context negation**

$$\frac{\Gamma \text{ ctx}}{\Gamma^- \text{ ctx}} \quad \overline{(\Gamma^-)^-} = \Gamma$$

$$\Gamma^- := \Gamma^{\text{op}}$$

We annotate **variances/polarities** by adding operations to MLTT

- **Type negation**

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash A^- \text{ type}} \quad \overline{(A^-)^-} = A$$

- **Context negation**

$$\frac{\Gamma \text{ ctx}}{\Gamma^- \text{ ctx}} \quad \overline{(\Gamma^-)^-} = \Gamma$$

- **Negative context extension**

We annotate **variances/polarities** by adding operations to MLTT

- **Type negation**

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash A^- \text{ type}} \quad \overline{(A^-)^-} = A$$

- **Context negation**

$$\frac{\Gamma \text{ ctx}}{\Gamma^- \text{ ctx}} \quad \overline{(\Gamma^-)^-} = \Gamma$$

- **Negative context extension**

$$\frac{\Gamma^- \vdash A \text{ type}}{\Gamma, - \ x : A \vdash}$$

We annotate **variances/polarities** by adding operations to MLTT

- **Type negation**

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash A^- \text{ type}} \quad \overline{(A^-)^- = A}$$

- **Context negation**

$$\frac{\Gamma \text{ ctx}}{\Gamma^- \text{ ctx}} \quad \overline{(\Gamma^-)^- = \Gamma}$$

- **Negative context extension**

$$\frac{\Gamma^- \vdash A \text{ type}}{\Gamma, - \ x: A \vdash} \quad (\Gamma, - \ x: A)^- = \Gamma^-, x: A^-$$

We annotate **variances/polarities** by adding operations to MLTT

- **Type negation**

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash A^- \text{ type}} \quad \overline{(A^-)^- = A}$$

- **Context negation**

$$\frac{\Gamma \text{ ctx}}{\Gamma^- \text{ ctx}} \quad \overline{(\Gamma^-)^- = \Gamma}$$

- **Negative context extension**

$$\frac{\Gamma^- \vdash A \text{ type}}{\Gamma, _- x : A \vdash} \quad \frac{\Gamma^- \vdash A \text{ type}}{(\Gamma, _- x : A)^- \vdash x : A^-}$$



$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma, x: A \vdash B \text{ type}}{\Gamma \vdash \Pi(A, B) \text{ type}}$$

$$\frac{\Gamma^- \vdash A \text{ type} \quad \Gamma, x : A \vdash B \text{ type}}{\Gamma \vdash \Pi(A, B) \text{ type}}$$

$$\frac{\Gamma^- \vdash A \text{ type} \quad \Gamma, \textcolor{red}{-} x: A \vdash B \text{ type}}{\Gamma \vdash \Pi(A, B) \text{ type}}$$

$$\frac{\Gamma^- \vdash A \text{ type} \quad \Gamma, \textcolor{red}{-} x: A \vdash B \text{ type}}{\Gamma \vdash \Pi(A, B) \text{ type}}$$

$$\frac{\Gamma, \textcolor{red}{-} x: A \vdash e: B}{\Gamma \vdash \lambda x^A. e: \Pi(A, B)}$$

$$\frac{\Gamma^- \vdash A \text{ type} \quad \Gamma, \textcolor{red}{-} x: A \vdash B \text{ type}}{\Gamma \vdash \Pi(A, B) \text{ type}}$$

$$\frac{\Gamma, \textcolor{red}{-} x: A \vdash e: B}{\Gamma \vdash \lambda x^A. e: \Pi(A, B)}$$

$$\frac{\Gamma \vdash f: \Pi(A, B) \quad \Gamma^- \vdash t: A^-}{\Gamma \vdash f(t): B(t)}$$

**Problem:** We can't  
do anything with  
these

- No identity function

- No identity function

- ▶  $A \rightarrow A$  isn't a well-formed type (domain has to be in  $\Gamma^-$ , codomain in  $\Gamma, - x: A$ )



$$\frac{\Gamma^- \vdash A \text{ type} \quad \Gamma, \textcolor{red}{-} x: A \vdash B \text{ type}}{\Gamma \vdash \Pi(A, B) \text{ type}}$$

$$\frac{\Gamma, \textcolor{red}{-} x: A \vdash e: B}{\Gamma \vdash \lambda x^A. e: \Pi(A, B)}$$

$$\frac{\Gamma \vdash f: \Pi(A, B) \quad \Gamma^- \vdash t: A^-}{\Gamma \vdash f(t): B(t)}$$

- No identity function

- ▶  $A \rightarrow A$  isn't a well-formed type (domain has to be in  $\Gamma^-$ , codomain in  $\Gamma, - x: A$ )

- No identity function

- ▶  $A \rightarrow A$  isn't a well-formed type (domain has to be in  $\Gamma^-$ , codomain in  $\Gamma, - x : A$ )
- ▶  $\lambda x^A. x$  isn't a well-formed term ( $x$  is a term in  $(\Gamma, - x : A)^-$ , not  $\Gamma, - x : A$ )

$$\frac{\Gamma^- \vdash A \text{ type} \quad \Gamma, \textcolor{red}{-} x: A \vdash B \text{ type}}{\Gamma \vdash \Pi(A, B) \text{ type}}$$

$$\frac{\Gamma, \textcolor{red}{-} x: A \vdash e: B}{\Gamma \vdash \lambda x^A. e: \Pi(A, B)}$$

$$\frac{\Gamma \vdash f: \Pi(A, B) \quad \Gamma^- \vdash t: A^-}{\Gamma \vdash f(t): B(t)}$$

We annotate **variances/polarities** by adding operations to MLTT

- **Type negation**

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash A^- \text{ type}} \quad \overline{(A^-)^- = A}$$

- **Context negation**

$$\frac{\Gamma \text{ ctx}}{\Gamma^- \text{ ctx}} \quad \overline{(\Gamma^-)^- = \Gamma}$$

- **Negative context extension**

$$\frac{\Gamma^- \vdash A \text{ type}}{\Gamma, - \ x: A \vdash} \quad \frac{\Gamma^- \vdash A \text{ type}}{(\Gamma, - \ x: A)^- \vdash x: A^-}$$

- No identity function

- ▶  $A \rightarrow A$  isn't a well-formed type (domain has to be in  $\Gamma^-$ , codomain in  $\Gamma, - x : A$ )
- ▶  $\lambda x^A. x$  isn't a well-formed term ( $x$  is a term in  $(\Gamma, - x : A)^-$ , not  $\Gamma, - x : A$ )

- No identity function
  - ▶  $A \rightarrow A$  isn't a well-formed type (domain has to be in  $\Gamma^-$ , codomain in  $\Gamma, - x : A$ )
  - ▶  $\lambda x^A. x$  isn't a well-formed term ( $x$  is a term in  $(\Gamma, - x : A)^-$ , not  $\Gamma, - x : A$ )
- Forget about composition...

We instead work in a **neutral context**, i.e. a groupoid (but types are still valued in categories)



We instead work in a **neutral context**, i.e. a groupoid (but types are still valued in categories)

- Can substitute between  $\Gamma$  and  $\Gamma^-$  (so types like  $A \rightarrow A$  can be well-formed)

We instead work in a **neutral context**, i.e. a groupoid (but types are still valued in categories)

- Can substitute between  $\Gamma$  and  $\Gamma^-$  (so types like  $A \rightarrow A$  can be well-formed)
- Can substitute between  $\Gamma, - \ x : A$  and  $\Gamma, x : A$  (so  $\lambda x^A. x$  is well-formed)

We instead work in a **neutral context**, i.e. a groupoid (but types are still valued in categories)

- Can substitute between  $\Gamma$  and  $\Gamma^-$  (so types like  $A \rightarrow A$  can be well-formed)
- Can substitute between  $\Gamma, -x : A$  and  $\Gamma, x : A$  (so  $\lambda x^A. x$  is well-formed)
- Get coercion operator

$$\frac{\Gamma \text{ Nctx} \quad \Gamma \vdash t : A^-}{\Gamma \vdash -t : A}$$

We instead work in a **neutral context**, i.e. a groupoid (but types are still valued in categories)

- Can substitute between  $\Gamma$  and  $\Gamma^-$  (so types like  $A \rightarrow A$  can be well-formed)
- Can substitute between  $\Gamma, - \ x: A$  and  $\Gamma, x: A$  (so  $\lambda x^A. x$  is well-formed)
- Get coercion operator

$$\frac{\Gamma \text{ Nctx} \quad \Gamma \vdash t: A^-}{\Gamma \vdash -t: A} \quad - \ -t = t$$

# 2 Directed Equality

$$\text{Id}(t, t') : \Gamma \rightarrow \mathbf{Grpd}$$

$$\text{Id}(t, t') \gamma = (A \gamma) [t \gamma, t' \gamma]$$

$$\text{Id}(t, t') \gamma_{01} x_0 = (t' \gamma_{01}) \circ A \gamma_{01} x_0 \circ (t \gamma_{01})^{-1}$$

$$\begin{array}{ccc}
 A \gamma_{01} (t \gamma_0) & \xrightarrow{A \gamma_{01} x_0} & A \gamma_{01} (t' \gamma_0) \\
 \downarrow t \gamma_{01} & & \downarrow t' \gamma_{01} \\
 t \gamma_1 & & t' \gamma_1
 \end{array}$$

**Directed Type Theory** includes **hom-types**:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash t : A \quad \Gamma \vdash t' : A}{\Gamma \vdash \text{Hom}_A(t, t') \text{ type}}$$

$$\Gamma \vdash t : A$$

$$\Gamma, x : A, u : \text{Hom}(t, x) \vdash M(x, u) \text{ type}$$

$$\Gamma \vdash m : M(t, \text{refl})$$

$$\frac{\Gamma \text{ ctx} \quad \Gamma \vdash t : A}{\Gamma \vdash \text{refl}_t : \text{Hom}(t, t)}$$

$$\frac{\Gamma \vdash m : M(t, \text{refl})}{\Gamma, x : A, u : \text{Hom}(t, x) \vdash J \ m \ x \ u : M(x, u)}$$

**Directed Type Theory** includes **hom-types**:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash t : A^- \quad \Gamma \vdash t' : A}{\Gamma \vdash \text{Hom}_{\mathbf{A}}(t, t') \text{ type}}$$

$$\Gamma \vdash t : A$$

$$\Gamma, x : A, u : \text{Hom}(t, x) \vdash M(x, u) \text{ type}$$

$$\Gamma \vdash m : M(t, \text{refl})$$

$$\frac{\Gamma \text{ ctx} \quad \Gamma \vdash t : A}{\Gamma \vdash \text{refl}_t : \text{Hom}(t, t)}$$

$$\frac{\Gamma \vdash m : M(t, \text{refl})}{\Gamma, x : A, u : \text{Hom}(t, x) \vdash J \ m \ x \ u : M(x, u)}$$



$$\text{Id}(t, t') : \Gamma \rightarrow \mathbf{Grpd}$$

$$\text{Id}(t, t') \gamma = (A \gamma) [t \gamma, t' \gamma]$$

$$\text{Id}(t, t') \gamma_{01} x_0 = (t' \gamma_{01}) \circ A \gamma_{01} x_0 \circ (t \gamma_{01})^{-1}$$

$$\begin{array}{ccc}
 A \gamma_{01} (t \gamma_0) & \xrightarrow{A \gamma_{01} x_0} & A \gamma_{01} (t' \gamma_0) \\
 \downarrow t \gamma_{01} & & \downarrow t' \gamma_{01} \\
 t \gamma_1 & & t' \gamma_1
 \end{array}$$

$$\mathbf{Hom}(t, t') : \Gamma \rightarrow \mathbf{Grpd}$$

$$\mathbf{Hom}(t, t') \gamma = (A \gamma) [t \gamma, t' \gamma]$$

$$\mathbf{Hom}(t, t') \gamma_{01} x_0 = (t' \gamma_{01}) \circ A \gamma_{01} x_0 \circ (t \gamma_{01})^{-1}$$

$$\begin{array}{ccc}
 A \gamma_{01} (t \gamma_0) & \xrightarrow{A \gamma_{01} x_0} & A \gamma_{01} (t' \gamma_0) \\
 \downarrow t \gamma_{01} & & \downarrow t' \gamma_{01} \\
 t \gamma_1 & & t' \gamma_1
 \end{array}$$

$$\mathbf{Hom}(t, t') : \Gamma \rightarrow \mathbf{Cat}$$

$$\mathbf{Hom}(t, t') \gamma = (A \gamma) [t \gamma, t' \gamma]$$

$$\mathbf{Hom}(t, t') \gamma_{01} x_0 = (t' \gamma_{01}) \circ A \gamma_{01} x_0 \circ (t \gamma_{01})^{-1}$$

$$\begin{array}{ccc}
 A \gamma_{01} (t \gamma_0) & \xrightarrow{A \gamma_{01} x_0} & A \gamma_{01} (t' \gamma_0) \\
 \downarrow t \gamma_{01} & & \downarrow t' \gamma_{01} \\
 t \gamma_1 & & t' \gamma_1
 \end{array}$$

$$\mathbf{Hom}(t, t') : \Gamma \rightarrow \mathbf{Cat}$$

$$\mathbf{Hom}(t, t') \gamma = (A \gamma) [t \gamma, t' \gamma]$$

$$\mathbf{Hom}(t, t') \gamma_{01} x_0 = (t' \gamma_{01}) \circ A \gamma_{01} x_0 \circ (t \gamma_{01})^{-1}$$

$$\begin{array}{ccc}
 A \gamma_{01} (t \gamma_0) & \xrightarrow{A \gamma_{01} x_0} & A \gamma_{01} (t' \gamma_0) \\
 \uparrow t \gamma_{01} & & \downarrow t' \gamma_{01} \\
 t \gamma_1 & & t' \gamma_1
 \end{array}$$

$$\mathbf{Hom}(t, t') : \Gamma \rightarrow \mathbf{Cat}$$

$$\mathbf{Hom}(t, t') \gamma = (A \gamma) [t \gamma, t' \gamma]$$

$$\mathbf{Hom}(t, t') \gamma_{01} x_0 = (t' \gamma_{01}) \circ A \gamma_{01} x_0 \circ (t \gamma_{01})$$

$$\begin{array}{ccc}
 A \gamma_{01} (t \gamma_0) & \xrightarrow{A \gamma_{01} x_0} & A \gamma_{01} (t' \gamma_0) \\
 \uparrow t \gamma_{01} & & \downarrow t' \gamma_{01} \\
 t \gamma_1 & & t' \gamma_1
 \end{array}$$

**Directed Type Theory** includes **hom-types**:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash t : A^- \quad \Gamma \vdash t' : A}{\Gamma \vdash \text{Hom}_A(t, t') \text{ type}}$$

$$\Gamma \vdash t : A$$

$$\Gamma, x : A, u : \text{Hom}(t, x) \vdash M(x, u) \text{ type}$$

$$\Gamma \vdash m : M(t, \text{refl})$$

$$\frac{\Gamma \text{ ctx} \quad \Gamma \vdash t : A}{\Gamma \vdash \text{refl}_t : \text{Hom}(t, t)}$$

$$\frac{\Gamma \vdash m : M(t, \text{refl})}{\Gamma, x : A, u : \text{Hom}(t, x) \vdash J \ m \ x \ u : M(x, u)}$$

**Directed Type Theory** includes **hom-types**:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash t : A^- \quad \Gamma \vdash t' : A}{\Gamma \vdash \text{Hom}_A(t, t') \text{ type}}$$

$$\Gamma \vdash t : A$$

$$\Gamma, x : A, u : \text{Hom}(t, x) \vdash M(x, u) \text{ type}$$

$$\Gamma \vdash m : M(t, \text{refl})$$

$$\frac{\Gamma \text{ Nctx} \quad \Gamma \vdash t : A}{\Gamma \vdash \text{refl}_t : \text{Hom}(t, t)}$$

$$\frac{\Gamma \vdash m : M(t, \text{refl})}{\Gamma, x : A, u : \text{Hom}(t, x) \vdash J \ m \ x \ u : M(x, u)}$$

**Directed Type Theory** includes **hom-types**:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash t: A^- \quad \Gamma \vdash t': A}{\Gamma \vdash \text{Hom}_A(t, t') \text{ type}}$$

$$\frac{\Gamma \vdash t: A \quad \Gamma, x: A, u: \text{Hom}(t, x) \vdash M(x, u) \text{ type} \quad \Gamma \vdash m: M(t, \text{refl})}{\Gamma, x: A, u: \text{Hom}(t, x) \vdash J \ m \ x \ u: M(x, u)}$$

$$\frac{\Gamma \text{ Nctx} \quad \Gamma \vdash t: A^-}{\Gamma \vdash \text{refl}_t: \text{Hom}(t, -t)}$$



**Directed Type Theory** includes **hom-types**:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash t : A^- \quad \Gamma \vdash t' : A}{\Gamma \vdash \text{Hom}_A(t, t') \text{ type}}$$

$$\frac{\Gamma \vdash t : A^-}{\Gamma, x : A, u : \text{Hom}(t, x) \vdash M(x, u) \text{ type}}$$

$$\frac{\Gamma \vdash m : M(t, \text{refl})}{\Gamma, x : A, u : \text{Hom}(t, x) \vdash J \ m \ x \ u : M(x, u)}$$

$$\frac{\Gamma \text{ Nctx} \quad \Gamma \vdash t : A^-}{\Gamma \vdash \text{refl}_t : \text{Hom}(t, -t)}$$

**Directed Type Theory** includes **hom-types**:


$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash t : A^- \quad \Gamma \vdash t' : A}{\Gamma \vdash \text{Hom}_A(t, t') \text{ type}}$$

$$\frac{\Gamma \vdash t : A^-}{\Gamma, x : A, u : \text{Hom}(t, x) \vdash M(x, u) \text{ type}}$$


$$\frac{\Gamma \vdash m : M(-t, \text{refl})}{\Gamma, x : A, u : \text{Hom}(t, x) \vdash J \ m \ x \ u : M(x, u)}$$

$$\frac{\Gamma \text{ Nctx} \quad \Gamma \vdash t : A^-}{\Gamma \vdash \text{refl}_t : \text{Hom}(t, -t)}$$

## Inverses

$$\frac{\Gamma \vdash p: \text{Hom}(t, t')}{\Gamma \vdash J \text{ refl}_t t' p: \text{Hom}(t', t)}$$


## Composition

$$\frac{\begin{array}{l} \Gamma \vdash p: \text{Hom}(t, t') \\ \Gamma \vdash q: \text{Hom}(t', t'') \end{array}}{\Gamma \vdash J p t'' q: \text{Hom}(t, t'')}$$


## Inverses

$$\frac{\Gamma \vdash p: \text{Hom}(t, t')}{\Gamma \vdash J \text{ refl}_t t' p: \text{Hom}(t', t)}$$

## Composition

$$\frac{\begin{array}{l} \Gamma \vdash p: \text{Hom}(t, t') \\ \Gamma \vdash q: \text{Hom}(-t', t'') \end{array}}{\Gamma \vdash J p t'' q: \text{Hom}(t, t')}$$

## Inverses

$$\frac{\Gamma \vdash p: \text{Hom}(t, t')}{\Gamma \vdash J \text{ refl}_t t' p: \text{Hom}(t', t)}$$

## Composition

$$\frac{\begin{array}{l} \Gamma \vdash p: \text{Hom}(t, t') \\ \Gamma \vdash q: \text{Hom}(-t', t'') \end{array}}{\Gamma \vdash J p t'' q: \text{Hom}(t, t'')} \quad \checkmark$$

## Inverses

$$\frac{\Gamma \vdash p: \text{Hom}(t, t')}{\Gamma \vdash J \text{ refl}_t t' p: \text{Hom}(-t', -t)}$$

## Composition

$$\frac{\begin{array}{l} \Gamma \vdash p: \text{Hom}(t, t') \\ \Gamma \vdash q: \text{Hom}(-t', t'') \end{array}}{\Gamma \vdash J p t'' q: \text{Hom}(t, t'')} \quad \checkmark$$

$$\begin{array}{c}
\Gamma \vdash t : A^- \\
\Gamma, x : A, u : \text{Hom}(t, x) \vdash M(x, u) \text{ type} \\
\Gamma \vdash m : M(-t, \text{refl}) \\
\hline
\Gamma, x : A, u : \text{Hom}(t, x) \vdash J \ m \ x \ u : M(x, u)
\end{array}$$

$$\begin{array}{c}
\Gamma \vdash t : A^- \\
\Gamma, x : A, u : \text{Hom}(t, x) \vdash M(x, u) \text{ type} \\
\Gamma \vdash m : M(-t, \text{refl}) \\
\hline
\Gamma, x : A, u : \text{Hom}(t, x) \vdash J \ m \ x \ u : \text{Id}(-x, -t)
\end{array}$$



$$\begin{array}{c}
\Gamma \vdash t : A^- \\
\Gamma, x : A, u : \text{Hom}(t, x) \vdash \text{Id}(-x, -t) \text{ type} \\
\Gamma \vdash m : M(-t, \text{refl}) \\
\hline
\Gamma, x : A, u : \text{Hom}(t, x) \vdash J \ m \ x \ u : \text{Id}(-x, -t)
\end{array}$$

$$\begin{array}{c}
\Gamma \vdash t : A^- \\
\Gamma, x : A, u : \text{Hom}(t, x) \vdash \text{Id}(-x, -t) \text{ type} \\
\Gamma \vdash m : M(-t, \text{refl}) \\
\hline
\Gamma, x : A, u : \text{Hom}(t, x) \vdash J \ m \ x \ u : \text{Id}(-x, -t)
\end{array}$$

We cannot negate  $x$ , since  $\Gamma, x : A$  is not neutral!

$$\begin{array}{c}
\Gamma \vdash t : A^- \\
\Gamma, x : A, u : \text{Hom}(t, x) \vdash \text{Id}(-x, -t) \text{ type} \\
\Gamma \vdash m : M(-t, \text{refl}) \\
\hline
\Gamma, x : A, u : \text{Hom}(t, x) \vdash m \ x \ u : \text{Id}(-x, -t)
\end{array}$$

We cannot negate  $x$ , since  $\Gamma, x : A$  is not neutral!


## Inverses

$$\frac{\Gamma \vdash p: \text{Hom}(t, t')}{\Gamma \vdash J \text{ refl}_t t' p: \text{Hom}(-t', -t)}$$


## Composition

$$\frac{\begin{array}{l} \Gamma \vdash p: \text{Hom}(t, t') \\ \Gamma \vdash q: \text{Hom}(-t', t'') \end{array}}{\Gamma \vdash J p t'' q: \text{Hom}(t, t'')} \quad \checkmark$$

## Inverses

$$\frac{\Gamma \vdash p: \text{Hom}(t, t')}{\Gamma \vdash J \text{ refl}_t t' p. \text{Hom}(-t', -t)}$$


## Composition

$$\frac{\begin{array}{l} \Gamma \vdash p: \text{Hom}(t, t') \\ \Gamma \vdash q: \text{Hom}(-t', t'') \end{array}}{\Gamma \vdash J p t'' q. \text{Hom}(t, t'')}$$


**Key Idea:** Use model  
to prove symmetry is  
independent

$$\mathbf{Hom}(t, t') : \Gamma \rightarrow \mathbf{Cat}$$

$$\mathbf{Hom}(t, t') \gamma = (A \gamma) [t \gamma, t' \gamma]$$

$$\mathbf{Hom}(t, t') \gamma_{01} x_0 = (t' \gamma_{01}) \circ A \gamma_{01} x_0 \circ (t \gamma_{01})$$

*This type has multiple terms in general, hence UIP cannot be proved from the rules.*

$$\text{Hom}(t, t') : \Gamma \rightarrow \mathbf{Cat}$$

$$\text{Hom}(t, t') \gamma = (A \gamma) [t \gamma, t' \gamma]$$

$$\text{Hom}(t, t') \gamma_{01} x_0 = (t' \gamma_{01}) \circ A \gamma_{01} x_0 \circ (t \gamma_{01})$$

*This type **is asymmetric** in general, hence UIP cannot be proved from the rules.*



$$\text{Hom}(t, t') : \Gamma \rightarrow \mathbf{Cat}$$

$$\text{Hom}(t, t') \gamma = (A \gamma) [t \gamma, t' \gamma]$$

$$\text{Hom}(t, t') \gamma_{01} x_0 = (t' \gamma_{01}) \circ A \gamma_{01} x_0 \circ (t \gamma_{01})$$

*This type **is asymmetric** in general, hence **symmetry** cannot be proved from the rules.*



- Synthetic category theory

- Synthetic category theory
  - ▶ *Inductive* style, where universal mapping properties are expressed as principles of induction

- Synthetic category theory
  - ▶ *Inductive* style, where universal mapping properties are expressed as principles of induction
- Directed homotopy theory

- Synthetic category theory
  - ▶ *Inductive* style, where universal mapping properties are expressed as principles of induction
- Directed homotopy theory
  - ▶ Analogous to interpretation of types in HoTT as homotopy spaces

- Synthetic category theory
  - ▶ *Inductive* style, where universal mapping properties are expressed as principles of induction
- Directed homotopy theory
  - ▶ Analogous to interpretation of types in HoTT as homotopy spaces
  - ▶ Application to concurrency

- Synthetic category theory
  - ▶ *Inductive* style, where universal mapping properties are expressed as principles of induction
- Directed homotopy theory
  - ▶ Analogous to interpretation of types in HoTT as homotopy spaces
  - ▶ Application to concurrency
- Synthetic rewriting systems



- Synthetic category theory
  - ▶ *Inductive* style, where universal mapping properties are expressed as principles of induction
- Directed homotopy theory
  - ▶ Analogous to interpretation of types in HoTT as homotopy spaces
  - ▶ Application to concurrency
- Synthetic rewriting systems
  - ▶ Can have intensional and extensional identity types in the system simultaneously, with hom-types sitting between them

- 📄 Daniel Gratzer, Jonathan Weinberger, and Ulrik Buchholtz.  
Directed univalence in simplicial homotopy type theory, 2024.
- 📄 Daniel Gratzer, Jonathan Weinberger, and Ulrik Buchholtz.  
The yoneda embedding in simplicial type theory, 2025.
- 📄 Martin Hofmann and Thomas Streicher.  
The groupoid interpretation of type theory.  
*Twenty-five years of constructive type theory (Venice, 1995)*, 36:83–111, 1995.
- 📄 Nikolai Kudasov, Emily Riehl, and Jonathan Weinberger.  
Formalizing the  $\infty$ -categorical yoneda lemma, 2023.
- 📄 Daniel R Licata and Robert Harper.  
2-dimensional directed dependent type theory.  
2011.

- 📄 Jacob Neumann and Thorsten Altenkirch.  
Synthetic 1-categories in directed type theory.  
*arXiv preprint arXiv:2410.19520*, 2024.
- 📄 Hayato Nasu.  
An internal logic of virtual double categories.  
*arXiv preprint arXiv:2410.06792*, 2024.
- 📄 Hayato Nasu.  
Logical aspects of virtual double categories.  
*arXiv preprint arXiv:2501.17869*, 2025.
- 📄 Jacob Neumann.  
*A Generalized Algebraic Theory of Directed Equality*.  
PhD thesis, University of Nottingham, 2025.

- 📄 Max S New and Daniel R Licata.  
A formal logic for formal category theory.  
*Foundations of Software Science and Computation Structures LNCS 13992*,  
page 113, 2023.
- 📄 Paige Randall North.  
Towards a directed homotopy type theory.  
*Electronic Notes in Theoretical Computer Science*, 347:223–239, 2019.
- 📄 Andreas Nuyts.  
Towards a directed homotopy type theory based on 4 kinds of variance.  
*Mém. de mast. Katholieke Universiteit Leuven*, 2015.
- 📄 E. Riehl and M. Shulman.  
A type theory for synthetic  $\infty$ -categories.  
*Higher Structures*, 1(1):116–193, 2017.

- Jonathan Weinberger.  
Strict stability of extension types, 2022.
- Jonathan Weinberger.  
A synthetic perspective on  $(\infty, 1)$ -category theory: Fibrational and semantic aspects.  
2022.
- Jonathan Weinberger.  
Internal sums for synthetic fibered  $(\infty, 1)$ -categories.  
*Journal of Pure and Applied Algebra*, 228(9):107659, 2024.
- Jonathan Weinberger.  
Two-sided cartesian fibrations of synthetic  $(\infty, 1)$ -categories.  
*Journal of Homotopy and Related Structures*, 19(3):297–378, 2024.



- Semantics of type theory in **Cat**

- Semantics of type theory in **Cat**
- Adopt a system of **polarity**, annotating variances



- Semantics of type theory in **Cat**
- Adopt a system of **polarity**, annotating variances
- Too much polarity, need to weaken somewhat with **neutrality**

- Semantics of type theory in **Cat**
- Adopt a system of **polarity**, annotating variances
- Too much polarity, need to weaken somewhat with **neutrality**
- Directed J-rule, which *can* prove transitivity/composition, but not symmetry/inverses

- Semantics of type theory in **Cat**
- Adopt a system of **polarity**, annotating variances
- Too much polarity, need to weaken somewhat with **neutrality**
- Directed J-rule, which *can* prove transitivity/composition, but not symmetry/inverses
- Synthetic category theory, directed homotopy theory, concurrency and rewriting

**Thank you!**