ChatGPT

WP1 – Foundations: Symplectic Reduction, Moment Maps, Principal Bundles, Magnetic Terms

Hamiltonian Group Actions and Momentum Maps

A **Hamiltonian G-action** on a symplectic manifold (M,ω) is a group action by symplectomorphisms that admits a **momentum map** $J: M \to \mathfrak{g}^*, where $\mathfrak{g}^$ is the dual of the Lie algebra of G. In practical terms, each Lie algebra element $\xi \in \mathfrak{g}$ generates an infinitesimal action (vector field X_ξ on M), and the momentum map is defined such that for all ξ [1]:

- **Defining equation:** $d\langle J,\xi\rangle = \iota_{X_\xi}\,\omega$, where $\iota_{X_\xi}\omega$ is the contraction of ω with the vector field X_ξ [1]. Equivalently, $\omega(X_\xi, \cdot) = d(J^\xi)$ where $J^\xi(p) = \langle J(p), \xi \rangle$ is the ξ-component of J at point p. This means the flow generated by X_ξ is (locally) governed by the Hamiltonian function J^ξ.

- **Noether's theorem link:** If $H: M\to \mathbb{R}$ is a G-invariant Hamiltonian, then each component $\langle J,\xi\rangle$ is a conserved quantity. The momentum map generalizes classical linear and angular momentum [2], and its existence is essentially the modern geometric formulation of Noether's theorem.

- **Existence conditions:** A symplectic G-action need not always be Hamiltonian (momentum map may not exist globally). The necessary and sufficient condition is that each one-form $\iota_{X_\xi} \omega$ is **exact** on M (not just closed) [3]. In simply connected or contractible manifolds this holds automatically, and for compact connected G, any symplectic action can often be adjusted (by choosing an appropriate constant) to admit a momentum map [4]. If G is compact or semisimple, one can usually choose the integration constants so that J is also **G-equivariant** (commutes with the coadjoint action) [4]. In what follows we assume these conditions are met so that J exists.

- **Example:** For $G = S^1$ (an abelian one-parameter symmetry), the Lie algebra dual is $\mathbb{R}$. A momentum map $J: M \to \mathbb{R}$ in this case is just an invariant Hamiltonian function generating the S^1-action [5]. For instance, if S^1 acts by rotations, J may be the angular momentum around that axis. The defining equation $dJ = \iota_{X}\omega$ says the vector field X (the rotation generator) is Hamiltonian with J as its Hamiltonian function.

Marsden–Weinstein Symplectic Reduction

Given a Hamiltonian G-action with momentum map J, **symplectic reduction** (Marsden–Weinstein–Meyer theorem) constructs a lower-dimensional phase space by "modding out" the symmetry while enforcing a fixed momentum value [6]. In brief [6] [7]:

- For a chosen value $\mu\in \mathfrak{g}^$ *(often $\mu=0$ for simplicity), consider the level set $J^{-1}(\mu)$, which is G-invariant. Under regularity conditions (0 should be a regular value of J, and G should act freely and properly on $J^{-1}(\mu)$), the quotient $M_\mu := J^{-1}(\mu)/G_\mu$ is a smooth manifold (here G_μ is the stabilizer of μ under coadjoint action; for μ in the center of $ \mathfrak{g}$^$ or G abelian, $G_\mu = G$)* [8].

- **Reduced symplectic form:** M_μ inherits a unique symplectic 2-form $\omega_{\rm red}$ such that $\iota^*omega* = *pi*^ \omega_{\rm red}$, where $\iota: J^{-1}(\mu)\hookrightarrow M$ and $\pi: J^{-1}(\mu)\twoheadrightarrow M_\mu$ are inclusion and projection. Intuitively, one restricts $ \omega$ to the constraint surface $J^{-1}(\mu)$ and then descends it to the quotient. The result $ (M_\mu,\omega_{\rm red})$ is called the **symplectic quotient** or **Marsden–Weinstein quotient** [9]. Its dimension is $\dim M_\mu = \dim M - 2\dim G$ (for μ a regular value) [10].

- This reduction can be viewed dynamically: if H is a G-invariant Hamiltonian on M, one can restrict H to $J^{-1}(\mu)$ and then project to M_μ. The **Dirac–Poisson brackets** on the constrained surface coincide with the Poisson structure of $\omega_{\rm red}$, so that H induces well-defined Hamiltonian flow on M_μ. In physical terms, one is "freezing out" the conserved quantity μ and eliminating the cyclic coordinates associated with G. (This is analogous to Routh reduction in Lagrangian mechanics, where ignorable coordinates are eliminated by fixing their conjugate momenta.)

- **Remark:** If μ is not zero and G is nonabelian, one must also incorporate the symplectic structure of the coadjoint orbit through μ. The general formula for the reduced space in that case is $M_\mu = J^{-1}(\mu)/G_\mu$ and its symplectic form includes an additional term coming from the Kirillov–Kostant–Souriau form on the coadjoint orbit $G\cdot\mu$. In this review (and especially for $G = S^1$ or other abelian groups relevant to KS theory) that subtlety is minimal because μ's orbit is a point and $G_\mu=G$.

Principal Bundles, Connections, and Curvature in Reduction

A common situation – crucial for KS regularization – is when the configuration space Q itself is a **principal G-bundle** over some base manifold $B = Q/G$. In that case, the phase space is naturally T^*Q *(with its canonical symplectic form), and one can perform cotangent bundle reduction by G. The momentum map $J: T*^Q \to \mathfrak{g}^$ in this case has a very concrete description* [11]*: at a cotangent vector $p_q \in T*^_q Q$,* $J(p_q)$ is just the **fiber momentum** – the projection of p_q onto the cotangent of the G-fiber (i.e. the component of p_q along the group orbits). Intuitively, $J(p_q)$ measures the conserved "angular" or gauge momentum associated with moving in the G-direction.

Connections and curvature: A principal G-bundle $\pi: Q \to B$ comes equipped with the concept of a **connection** – a 1-form A on Q with values in $\mathfrak{g}$ that prescribes a horizontal subspace at

each point. The connection A allows one to “lift” paths from the base B to horizontal paths in Q. Its **curvature** F_A is a $\mathfrak{g}$-valued 2-form on B defined (for an abelian G like S^1) simply as $F = dA$ (more generally $F = dA + \tfrac{1}{2}[A\wedge A]$, but $[A,A]=0$ for abelian groups). In physical terms, A is like a gauge potential and $F = dA$ is the field strength on the base.

When reducing T^*Q by the G-action, the connection and curvature play a direct role in the reduced symplectic form. Assume G is abelian (so that μ can be identified with a real parameter or a weight in $\mathfrak{g}^\cong \mathbb{R}$). Marsden–Weinstein **cotangent bundle reduction** yields the following important result [12] [13]:

- If $\mu = 0$, the reduced space is canonically isomorphic to T^*B with the usual symplectic form. In other words, imposing $J=0$ effectively enforces the “no momentum in the fiber” constraint, so one is just left with base motions (shapes) and their conjugate momenta [14].

- If $\mu \neq 0$, the reduced space is **still** diffeomorphic to T^*B, but its symplectic form is twisted by the curvature. Specifically, one finds a “magnetic” symplectic form on T^*B: $$\omega_{\text{red}} \;=\; \omega_{\text{can}} \;-\; \beta_\mu,$$ where ω_{can} is the canonical 2-form on T^*B, and β_μ is the pullback (to T^*B) of a 2-form σ_μ on B given by the curvature. In fact $\sigma_\mu = \mu\,F_A$ in many conventions. More precisely, if A is a chosen principal connection 1-form on Q, then σ_μ is the μ-component of its curvature [13]. In formula form, for $G = S^1$ identified with $U(1)$, one can write **$$\sigma_\mu \;=\; d\langle \mu, A\rangle = \mu\,dA,$$** since μ is just a constant factor in the Lie algebra (here $\langle\mu,A\rangle$ means we pair $\mu\in\mathfrak{g}^\cong \mathbb{R}$ with the $\mathfrak{g}$-valued A) [13]. Then $\beta_\mu = \pi_B^*(\sigma_\mu)$ is its pullback to T^*B via the bundle projection $\pi_B: T^*B \to B$ [15]. The sign of β_μ (adding vs. subtracting) depends on conventions for J and ω; here it is written as $\omega_{\rm red} = \omega_{\rm can} - \beta_\mu$ in agreement with the cited reference [16].

- This extra term β_μ is often called a **magnetic 2-form** on the base B [17]. It is closed ($d\sigma_\mu=0$ because $dF_A=0$ for a connection) but generally not exact on B if the bundle has nontrivial topology (in fact $[\sigma_\mu]$ in $H^2(B)$ captures the **Chern class** or monopole charge). In local coordinates, one can think of A like a vector potential and $\mu\,F_A$ like a magnetic field. The reduced equations of motion on B thus describe a particle moving under the influence of a magnetic field (hence the term “magnetic term” in the symplectic form) [17]. Geometrically, what was a straight free motion upstairs (with constant momentum μ along the fiber) becomes a charged particle motion downstairs – straight motion deflected by curvature.

Explicit example (Dirac monopole): A classic instance of this construction is the **Hopf fibration** $S^3 \to S^2$ with fiber S^1. Take $Q=S^3$ (thought of as the unit sphere in $\mathbb{R}^4$) and $G = S^1$ acting by rotations on the Hopf fiber. The base $B = S^2$ can be viewed as $\mathbb{R}^3\setminus{0}$ modulo radial rescaling. The standard connection on S^3 (viewed as a principal $U(1)$ bundle over S^2) has curvature equal to the area 2-form on S^2 – effectively a magnetic monopole field on $\mathbb{R}^3\setminus{0}$. If one takes the cotangent bundle T^*S^3 and reduces by S^1 at momentum μ, the reduced phase space is T^*S^2 carrying a symplectic form [18]:

$$ \omega_{\rm red} \;=\; \sum_{i=1}^3 dp_i \wedge dq_i \;-\; \mu\,\frac{q_1\,dq_2\wedge dq_3 + q_2\,dq_3\wedge dq_1 + q_3\,dq_1\wedge dq_2}{r^3}\,, $$

where (q_1,q_2,q_3) are Cartesian coordinates on $B=\mathbb{R}^3\setminus{0}$ (with $r=\sqrt{q_1^2+q_2^2+q_3^2}$) and p_i are conjugate momenta [19] [20]. The second term is **exactly** μ times the pullback of the area form on the unit 2-sphere (the Dirac monopole 2-form). This $\omega_{\rm red}$ is nondegenerate and closed, but it is *not* the standard canonical form $dp_i\wedge dq_i$ – it has been "twisted" by the magnetic term. In physical terms, a free particle on S^3 (with fixed $U(1)$ momentum μ) descends to a particle moving in $\mathbb{R}^3$ under a central magnetic field of charge μ [18]. Notably, the presence of this curvature term breaks the explicit rotational symmetry that a free particle on S^2 would have had; it is the geometric trace of having reduced a larger symmetry.

Magnetic Terms and the Role of Connections

The appearance of the curvature 2-form σ_μ in $\omega_{\rm red}$ has profound implications:

- **Twisted Poisson brackets:** Observables on the base B acquire modified Poisson brackets due to the term μF_A. For instance, if (q_i,p_i) are canonical coordinates on T^*B, the only nonzero fundamental Poisson brackets in absence of magnetic term are ${q_i,p_j}=\delta_{ij}$. With the magnetic 2-form, the brackets get extra terms* like ${p_i,p_j} = \mu\,(\ast F_{ij})(q)$ reflecting the Lorentz force law. In the Dirac monopole example, ${p_i,p_j} = \mu\,\epsilon_{ijk}q_k/r^3$, meaning the momentum components fail to commute and in fact realize the $\mathfrak{so}(3)$ Lie algebra (this is the origin of the conserved Runge–Lenz vector in the Kepler problem with a $1/r$ potential [21]).

- **No global momentum map on base:** The so-called "magnetic" flow on $(B,\omega_{\rm red})$ typically does *not* admit a global momentum map for the G-symmetry that was reduced. In the monopole case, the $SO(3)$ rotation symmetry on S^2 has no single-valued Hamiltonian potential: the angular momentum is "missing" an component because of the monopole field (one encounters a Dirac string singularity if one tries to define it globally). In general, the presence of a non-exact 2-form means that the lifted symmetry (which *did* have a momentum map upstairs, by construction) fails to be Hamiltonian downstairs – this is an important observation in reduction theory [22] [23]. The lack of an equivariant momentum map is precisely tied to the nontrivial cohomology class of the magnetic term. This is one way to understand how a problem (like the full three-body problem) can lack conserved quantities "downstairs" even if some higher-dimensional lift has hidden symmetries: the obstruction is carried by the curvature two-form.

- **Mechanical connection:** In many mechanical systems with symmetry, one chooses a connection called the *mechanical connection* (or *Locking connection*) such that $\langle J(q,\dot q), \xi \rangle = \langle \text{(momentum)}, \xi\rangle$ vanishes for horizontal velocities. In essence, the mechanical connection identifies the condition $J=0$ with "zero angular momentum" directions (horizontally moving in shape space). The curvature of the mechanical connection then appears as **Coriolis forces** or geometric phases in the reduced dynamics. For example, in the context of two-body Kepler problem, the mechanical $SO(3)$-connection on $Q=\mathbb{R}^3\setminus{0}$ (center-of-mass frame) has curvature proportional to the solid angle form, which is related to the Lenz vector precession [21]. For the three-body problem, the curvature of the appropriate connection on shape space would similarly encode the anholonomy in the dynamics (sometimes described as the **holonomy of the zero-angular-momentum frame**). In short, a nonzero curvature ($F_A \neq 0$) is a harbinger of nontrivial **geometric phase** and often of **nonintegrability** in the reduced system.

- **Curvature as integrability obstacle:** From a modern viewpoint (e.g. Morales–Ramis theory of differential Galois integrability conditions), a nonzero magnetic term can obstruct integrability. The reason is that the lift to a covering space (adding an extra angle coordinate) might restore enough symmetry to allow additional first integrals. If one suspects a hidden symmetry in a problem (like the KS or other lifts in the 3-body problem), a good strategy is to add coordinates so that the troublesome magnetic term becomes **exact** (or vanishes) in an extended phase space. This is exactly what the KS transformation accomplishes in the two-body problem: it introduces additional phase variables (a phase angle, or an $SU(2)$ spinor, etc.) such that the effective magnetic monopole in S^2 disappears upstairs (replaced by trivial linear motion on S^3). Our conjecture is that something analogous can be done for the 3-body problem by finding a suitable extension of phase space where the "curvature" lives in an extra degree of freedom and the downstairs chaos might be tamed by upstairs integrals.

Jacobi–Maupertuis and Eisenhart Lifts: Geodesics in Extended Spaces

In the broader context of regularization and "lifts", it is worth mentioning two classical techniques that relate to adding dimensions or restructuring the problem geometry:

- **Jacobi–Maupertuis principle:** For a conservative mechanical system with total energy E, Jacobi's theorem says the trajectories can be reparametrized and seen as geodesics of a Riemannian metric on configuration space (the **Jacobi metric** $g_{ij}^{J} = (E - V(q))\,g_{ij}^{\text{kin}}$) [24]. In effect, given $H = T + V$, the solution curves in configuration space (for fixed E) extremize the **Maupertuis action** $\int ! \sqrt{2(E-V)}\,ds$ and thus are geodesics of the conformally scaled metric. This is a different kind of "reduction": it eliminates time by using energy as a constant and turns the dynamics into a geodesic flow on a curved manifold (Q, g^J). While Jacobi–Maupertuis (JM) is a *reformulation* rather than an expansion of phase space, it provides insight into integrability (geodesic flows on certain manifolds are integrable if the metric has special properties). For example, the two-body Kepler problem corresponds to geodesic flow on S^3 (for bound orbits) or hyperbolic H^3 (for $E>0$), explaining the extra symmetries (Runge–Lenz vector) as Killing vectors of these spaces [25] [24]. In the three-body problem, Jacobi–Maupertuis metrics have been used to investigate chaotic curvature fluctuations and geodesic instability on the shape space manifold [26].

- **Eisenhart–Duval lift:** This is a method to embed a time-dependent or potential-driven system into a higher-dimensional *geodesic* system by adding auxiliary coordinates. Originally developed by L.P. Eisenhart (1928) and later popularized by Duval et al., the idea is to introduce an extra configuration coordinate (and sometimes the time as a coordinate) such that the motion in the extended space has zero potential and zero curvature (free motion or geodesic in a higher dimension). The potential energy and time reparameterization are encoded in the geometry of the extended manifold. **In essence, the Eisenhart lift converts a natural Hamiltonian $H=\frac{1}{2}g^{ij}p_ip_j + V(q)$ into a geodesic Hamiltonian in one extra dimension.** A concise description [27] is: *"The Jacobi–Eisenhart lift eliminates the potential in an n-dimensional Hamiltonian system, reducing it to a free particle geodesic in $(n+1)$ dimensions"*. All conserved quantities of the original system become geometric constants of motion (e.g. Killing vectors or first integrals) in the extended space, and integrability can only improve or stay the same under the lift (never worsen) [28]. The Eisenhart lift is closely related to the Kaluza–Klein mechanism in general relativity – for instance, a charged particle in a

electromagnetic field can be seen as a free geodesic in one higher dimension (with the extra momentum representing the electric charge). Likewise, for mechanical systems, adding a cyclic coordinate whose momentum stays fixed can mimic the effect of a gauge field on the base dynamics.

In summary, **Jacobi–Maupertuis and Eisenhart lifts complement symplectic reduction: they are ways of using geometry to handle forces.** Jacobi's trick turns potential forces into curvature of a Riemannian metric on configuration space (no new dimensions, but a new metric), whereas Eisenhart's trick turns potential or magnetic forces into curvature-free geodesic motion on an extended phase space (one or two extra dimensions accounting for the potential or gauge potential). Both are valuable in understanding how adding dimensions or structure can simplify dynamics. In the context of our conjecture, the **Kustaanheimo–Stiefel (KS) transformation** can be viewed as a mix of these ideas: it adds coordinates (like an Eisenhart lift) to remove a singular potential and yields an S^3 constraint (a curved space where geodesics correspond to Kepler orbits, akin to Jacobi's metric picture). By studying WP2–WP4 in light of the foundational concepts above, we aim to pinpoint how a **generalized KS lift for three-body dynamics** might similarly convert hard "magnetic" or curvature terms into simpler linear motion in a higher-dimensional space, making hidden symmetries manifest.

Sources:

- Marsden, R. & Weinstein, A. (1974). *Reduction of symplectic manifolds with symmetry*. **Rep. Math. Phys., 5**, 121–130. (Marsden–Weinstein reduction theorem: $J^{-1}(\mu)/G_\mu$ inherits a symplectic form) [6] [9].

- Abraham, R. & Marsden, J. (1978). *Foundations of Mechanics*. (Definitions of Hamiltonian actions and momentum maps; connection between symmetry and conserved quantities via Noether's theorem) [3] [29].

- Ortega, J.-P. & Ratiu, T. (2004). *Momentum Maps and Hamiltonian Reduction*. (Comprehensive treatment of momentum maps; cotangent bundle reduction with connections, see Theorem 2.3.3 for explicit construction) [30] [15].

- Cushman, R. & Bates, L. (1997). *Global Aspects of Classical Integrable Systems*. (Examples of non-trivial magnetic terms in reduced systems and their effect on integrability).

- Yang, C. & Khesin, B. (2020). "Averaging, symplectic reduction, and central extensions." *Nonlinearity, 33*(3), 1342–1369. (Theorem 2.1 gives the cotangent bundle reduction with magnetic term formula; Remark 2.2 names the magnetic term and its physical interpretation) [12] [17].

- Marsden, J. et al. (1990). "Hamiltonian systems with nonholonomic constraints" *Arch. Rational Mech. Anal., 115*, 1–25. (Mechanical connection and curvature forces in reduced dynamics).

- Chanda, S., Gibbons, G.W., Guha, P. (2017). *Jacobi–Maupertuis metric and Kepler equation*. (Discusses Jacobi and Eisenhart lifts in integrable systems; the Jacobi–Eisenhart lift as a way to remove potentials by extending dimension) [27] [24].

- Duval, C., Burdet, G., Künzle, H., & Perrin, M. (1985). “Bargmann structures and Newton–Cartan theory.” *Phys. Rev. D, 31*(8), 1841–1853. (Classic paper on Eisenhart lift and its relation to Bargmann (Newton-Cartan) geometry).

- Cushman, R. & Duistermaat, J.J. (1988). “The quantum mechanical spherical pendulum.” *Bull. Amer. Math. Soc.*, 19(2), 475–479. (Magnetic monopole terms appearing in the reduced phase space of a pendulum – an analogy to our discussion).

- Montgomery, R. (1990). “The argument of perihelion and the problem of attitude.” *Amer. J. Phys.*, 59(5), 394–398. (Geometric phase in the Kepler problem; discusses the curvature associated with the mechanical connection on $SO(3)$ and how the Runge–Lenz vector arises from it).

---

1 2 3 4 5 6 9 10 29 Momentum map - Wikipedia
https://en.wikipedia.org/wiki/Momentum_map

7 8 11 12 13 14 15 16 17 30 arxiv.org
https://arxiv.org/pdf/1806.01755

18 19 20 arxiv.org
https://arxiv.org/pdf/1109.5449

21 The four-dimensional conformal Kepler problem reduces to the three
https://pubs.aip.org/aip/jmp/article-pdf/27/6/1523/19154268/1523_1_online.pdf

22 The reduced spaces of a symplectic Lie group action - math - arXiv
https://arxiv.org/abs/math/0501098

23 [PDF] Symplectic Reduction for Semidirect Products and Central Extensions.
https://www.cds.caltech.edu/~marsden/bib/1998/05-MaMiPeRa1998/MaMiPeRa1998.pdf

24 25 26 27 28 [1612.07395] Jacobi-Maupertius metric and Kepler equation
https://ar5iv.org/pdf/1612.07395