

ALGEBRA

Lines

Slope of the line through $P_1 = (x_1, y_1)$ and $P_2 = (x_2, y_2)$:

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

Slope-intercept equation of line with slope m and y -intercept b :

$$y = mx + b$$

Point-slope equation of line through $P_1 = (x_1, y_1)$ with slope m :

$$y - y_1 = m(x - x_1)$$

Point-point equation of line through $P_1 = (x_1, y_1)$ and $P_2 = (x_2, y_2)$:

$$y - y_1 = m(x - x_1) \quad \text{where } m = \frac{y_2 - y_1}{x_2 - x_1}$$

Lines of slope m_1 and m_2 are parallel if and only if $m_1 = m_2$.

Lines of slope m_1 and m_2 are perpendicular if and only if $m_1 = -\frac{1}{m_2}$.

Circles

Equation of the circle with center (a, b) and radius r :

$$(x - a)^2 + (y - b)^2 = r^2$$

Distance and Midpoint Formulas

Distance between $P_1 = (x_1, y_1)$ and $P_2 = (x_2, y_2)$:

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

Midpoint of $\overline{P_1 P_2}$: $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$

Laws of Exponents

$$x^m x^n = x^{m+n}$$

$$\frac{x^m}{x^n} = x^{m-n}$$

$$(x^m)^n = x^{mn}$$

$$x^{-n} = \frac{1}{x^n}$$

$$(xy)^n = x^n y^n$$

$$\left(\frac{x}{y}\right)^n = \frac{x^n}{y^n}$$

$$x^{1/n} = \sqrt[n]{x}$$

$$\sqrt[n]{xy} = \sqrt[n]{x} \sqrt[n]{y}$$

$$\sqrt[n]{\frac{x}{y}} = \frac{\sqrt[n]{x}}{\sqrt[n]{y}}$$

$$x^{m/n} = \sqrt[n]{x^m} = (\sqrt[n]{x})^m$$

Special Factorizations

$$x^2 - y^2 = (x + y)(x - y)$$

$$x^3 + y^3 = (x + y)(x^2 - xy + y^2)$$

$$x^3 - y^3 = (x - y)(x^2 + xy + y^2)$$

Binomial Theorem

$$(x + y)^2 = x^2 + 2xy + y^2$$

$$(x - y)^2 = x^2 - 2xy + y^2$$

$$(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$$

$$(x - y)^3 = x^3 - 3x^2y + 3xy^2 - y^3$$

$$(x + y)^n = x^n + nx^{n-1}y + \frac{n(n-1)}{2}x^{n-2}y^2 + \dots + \binom{n}{k}x^{n-k}y^k + \dots + nx y^{n-1} + y^n$$

$$\text{where } \binom{n}{k} = \frac{n(n-1)\dots(n-k+1)}{1 \cdot 2 \cdot 3 \cdot \dots \cdot k}$$

Quadratic Formula

$$\text{If } ax^2 + bx + c = 0, \text{ then } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Inequalities and Absolute Value

If $a < b$ and $b < c$, then $a < c$.

If $a < b$, then $a + c < b + c$.

If $a < b$ and $c > 0$, then $ca < cb$.

If $a < b$ and $c < 0$, then $ca > cb$.

$$|x| = x \quad \text{if } x \geq 0$$

$$|x| = -x \quad \text{if } x \leq 0$$



GEOMETRY

Formulas for area A , circumference C , and volume V

Triangle

$$A = \frac{1}{2}bh$$

$$= \frac{1}{2}ab \sin \theta$$



Circle

$$A = \pi r^2$$

$$C = 2\pi r$$




Sector of Circle

$$A = \frac{1}{2}r^2\theta$$

$$s = r\theta$$


(θ in radians)



Sphere


$$V = \frac{4}{3}\pi r^3$$

$$A = 4\pi r^2$$



Cylinder

$$V = \pi r^2 h$$



Cone

$$V = \frac{1}{3}\pi r^2 h$$

$$A = \pi r \sqrt{r^2 + h^2}$$



Cone with arbitrary base

$$V = \frac{1}{3}Ah$$

where A is the area of the base



Pythagorean Theorem: For a right triangle with hypotenuse of length c and legs of lengths a and b , $c^2 = a^2 + b^2$.