


Machine Learning for Gravitational Waveforms from Binary Neutron Star mergers

Jacopo Tissino

Advisors: Dr. Sebastiano Bernuzzi, Dr. Michela Mapelli


2021-10-21

GW170817: the first BNS merger detection




Time-frequency representation of the chirping waveform (Abbott et al. 2017).

BBH merger waveform



BNS merger waveform



The basics of GW data analysis

The signal is modelled as $s(t) = h_\theta(t) + n(t)$, where:

- the noise $n(t)$ is taken to be stationary, with zero mean, and Gaussian with power spectral density $S_n(f)$;
- the signal $h_\theta(t)$ from a binary neutron star merger depends on:
 - intrinsic parameters: mass ratio $q = m_1/m_2$, spins $\vec{\chi}_1$ and $\vec{\chi}_2$, tidal polarizabilities Λ_1 and Λ_2 ;
 - extrinsic parameters: total mass $M = m_1 + m_2$, luminosity distance D_L , inclination $\iota \dots$

The Wiener distance

The likelihood used in parameter estimation reads (Maggiore 2007):

$$\Lambda(s|\theta) \propto \exp\left((h_\theta|s) - \frac{1}{2}(h_\theta|h_\theta)\right), \quad (1)$$

where $(a|b)$ is the Wiener product:

$$(a|b) = 4 \operatorname{Re} \int_0^\infty \frac{\tilde{a}^*(f) \tilde{b}(f)}{S_n(f)} \mathrm{d}f. \quad (2)$$

Models and accuracy


The main strategies for the generation of theoretical waveforms are:

- numerical relativity;
- effective one body;
- post-Newtonian.


Their accuracy is computed according to the **mismatch**

$$\mathcal{F}[h_1, h_2] = 1 - \max_{t_0, \phi_0} \frac{(h_1 | h_2(t_0, \phi_0))}{\underbrace{\sqrt{(h_1 | h_1)(h_2 | h_2)}}_{\cos \theta}} . \quad (3)$$


Amplitudes




Phases




mlgw_bns structure




Residuals: amplitude



Residuals: phase



PCA mismatches




Hyperparameter optimization


Pareto-front Plot




Fidelity




Amplitude reconstruction residuals




Phase reconstruction residuals




Evaluation time, varying initial frequency.




Evaluation time, varying initial frequency.




Evaluation time: $f_0 = 12$ Hz, downsampling.



Profiling the evaluation: 2×10^6 interpolation points




Profiling the evaluation: 8×10^3 interpolation points




Conclusions

- `mlgw_bns` is a surrogate model trained on effective one body waveforms;
- it is accurate (enough not to bias parameter estimation);
- it is fast(er than the state-of-the-art model it is trained on).


Fourier transform issues



Fourier transform issues



Fourier transform issues



Natural units and quadrupolar approximation

$$\frac{|\tilde{h}(f)|}{M} \approx \frac{1}{\pi^{2/3}} \sqrt{\frac{5}{24}} (Mf)^{-7/6} \frac{M}{r} \sqrt{\frac{m_1 m_2}{M^2}} \quad (4)$$

$$|h_+(f)| = \frac{1 + \cos^2 \iota}{2} |h(f)| \quad (5)$$

$$|h_\times(f)| = \cos \iota |h(f)|. \quad (6)$$

Fidelity requirements

For detection: with a mismatch \mathcal{F} we miss approximately a fraction

$$1 - (1 - \mathcal{F})^3 \quad (7)$$


of the signals.

For parameter estimation: in order for the modelling error to be at a 1σ level we need


$$\mathcal{F} \lesssim \frac{D}{2\text{SNR}^2}, \quad (8)$$

where D is the number of intrinsic parameters we estimate.

Power Spectral densities and GW170817



Fidelity compared to TF2



Bibliography



Abbott, B. P. et al. (Oct. 16, 2017). “GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral”. In: *Physical Review Letters* 119.16, p. 161101.



Maggiore, Michele (Nov. 24, 2007). *Gravitational Waves: Volume 1: Theory and Experiments*. 1 edition. Oxford: Oxford University Press. 576 pp.