

# Path Integral

Alessandro Lovo, Alberto Facheris, Francesco Gentile,  
Giorgio Mentasti, Jacopo Tissino, Leonardo Zampieri

Autumn 2019

## 1 Lesson 1

In lesson 1...

## 2 Lesson 4

In the previous lessons we introduced the Dirac representation, where we defined:

$$|\alpha t_0 t\rangle = U(t - t_0) |\alpha t_0\rangle \quad (1a)$$

$$= e^{-\frac{iH(t-t_0)}{\hbar}} |\alpha\rangle \quad (1b)$$

$$= \sum_{a'} e^{-\frac{iE_{a'}(t-t_0)}{\hbar}} |a'\rangle \langle a' | \alpha t_0 \rangle \quad (1c)$$

$$= \sum c'_a(t_0) e^{-\frac{iE_{a'}(t-t_0)}{\hbar}} |a'\rangle \quad (1d)$$

$$= \sum c'_a(t) |a'\rangle \quad (1e)$$

and

$$\langle \bar{x}' | a' t_0 t \rangle = \sum c'_a(t_0) e^{-\frac{iE_{a'}(t-t_0)}{\hbar}} \langle \bar{x}' | a' \rangle, \quad (2)$$

such that  $\psi(\bar{x}', t) = \sum c'_a(t_0) e^{-\frac{iE_{a'}(t-t_0)}{\hbar}} u_{a'}(\bar{x}')$ . Than from Dirac completeness relation,

$$c_{a'}(t_0) = \langle a' | \alpha t_0 \rangle = \int d^3x' \langle a' | \bar{x}' \rangle \langle \bar{x}' | \alpha t_0 \rangle, \quad (3)$$

and for this reason

$$\langle x'' | \alpha t_0 t \rangle = \sum e^{-\frac{iE_{a'}(t-t_0)}{\hbar}} \langle x'' | a' \rangle \int d^3 x' \langle a' | x' \rangle \langle x' | \alpha t_0 t \rangle \quad (4a)$$

$$= \int d^3 x' \sum \langle x'' | a' \rangle \langle a' | x' \rangle e^{-\frac{iE_{a'}(t-t_0)}{\hbar}} \langle x' | \alpha t_0 t \rangle \quad (4b)$$

$$= \int d^3 x' K \langle x' | \alpha t_0 t \rangle \quad (4c)$$

where we defined the propagator  $K$ . We write otherwise  $\psi(x'', t) = \int d^3 x' K(x'', t, x', t_0) \psi(x', t_0)$ , such that the propagator satisfies:

- $K$  respects Schrodinger equation, since  $i\hbar \langle x' | \alpha t_0 t \rangle = \langle x' | H | \alpha t_0 t \rangle$  and  $\langle x' | a' t_0 t \rangle = e^{-\frac{iE_{a'}(t-t_0)}{\hbar}} \langle x'' | a' \rangle$
- $\lim_{t \rightarrow t_0} K = \delta^3(x'' - x')$ , i.e. the evolution of a particle which was in  $x'$  at  $t = 0$

## 2.1 Free particle propagator

Now we want to obtain the propagator  $K$  aforementioned for the particular case of a free particle hamiltonian. In the space of momenta we have

$$P |p'\rangle = p' |p'\rangle, \langle x' | p' \rangle = \frac{e^{\frac{ip'x'}{\hbar}}}{\sqrt{2\pi\hbar}}, \quad H |p'\rangle = \frac{p'^2}{2m} |p'\rangle. \quad (5)$$

In this case we have

$$K = \int dp' \langle x'' | p' \rangle \langle p' | x' \rangle e^{-\frac{ip'^2(t-t_0)}{2m\hbar}} = \int_{-\infty}^{+\infty} dp' \frac{e^{\frac{i}{\hbar} \left( p'(x''-x') - \frac{p'^2}{2m}(t-t_0) \right)}}{2\pi\hbar} \quad (6)$$

Considering that

$$\frac{i}{\hbar} \left( p'(x''-x') - \frac{p'^2}{2m}(t-t_0) \right) = -\frac{i(t-t_0)}{2m\hbar} \left( p'^2 - \frac{2mp'(x''-x')}{t-t_0} \right) = -\frac{i(t-t_0)}{2m\hbar} \left( p' - \frac{m(x''-x')}{t-t_0} \right)^2 + \frac{i(t-t_0)}{2m\hbar} \frac{m^2(x''-x')^2}{(t-t_0)^2} \quad (7)$$

Now, defining the new variable  $\xi = p' - \frac{m(x''-x')}{t-t_0}$ , we finally obtain

$$K = \frac{e^{\frac{im(x''-x')^2}{2m\hbar}}}{2\pi\hbar} \int_{-\infty}^{+\infty} d\xi e^{-\frac{i(t-t_0)\xi^2}{2m\hbar}} = \frac{e^{\frac{im(x''-x')^2}{2m\hbar}}}{2\pi\hbar} \sqrt{\frac{2m\pi\hbar}{i(t-t_0)}} \quad (8)$$

## 2.2 Harmonic oscillator

In the same way we studied the free particle problem, we want to obtain the propagator for the harmonic oscillator, i.e. a single particle system under the evolution with the hamiltonian

$$H = \frac{p^2}{2m} + \frac{m\omega^2 x^2}{2}, \quad (9)$$

where we know that defining the operator distruction  $a = \sqrt{\frac{m\omega}{2\hbar}} \left( x + \frac{ip}{m\omega} \right)$ , we obtain an explicit writing for the eigenvalues  $E_N = \hbar\omega(N + \frac{1}{2})$  and the eigenfunctions respect the relations  $|n+1\rangle = \frac{a^\dagger}{\sqrt{n+1}}|n\rangle$ , in such a way that

$$\langle x' | a | 0 \rangle = \sqrt{\frac{m\omega}{2\hbar}} \langle x' | \left( x + \frac{ip}{m\omega} \right) | 0 \rangle. \quad (10)$$

It follows that  $\left( x' + \frac{i}{m\omega} (-i\hbar \frac{d}{dx'}) \right) \langle x' | 0 \rangle$  and, defining  $x_0 = \sqrt{\frac{\hbar}{m\omega}}$ , we have  $\langle x' | 0 \rangle = \eta e^{-\frac{x'^2}{2x_0^2}}$  with  $\eta = \frac{1}{(x_0\pi)^{\frac{1}{4}}}$ . In this case we obtain

$$\langle x' | n \rangle = \frac{1}{\pi^{\frac{1}{4}} \sqrt{2^n n}} \frac{\left( x' - x_0^2 \frac{d}{dx'} \right)^n}{x_0^{n+\frac{1}{2}}} e^{-\frac{x'^2}{2x_0^2}} = \frac{1}{\pi^{\frac{1}{4}} \sqrt{2^n n!}} H_n(y) e^{-\frac{x'^2}{2x_0^2}}, \quad (11)$$

where we defined  $y = \frac{x'}{x_0}$ , and the propagator

$$K = \sum_{n=0}^{\infty} \langle x' | n \rangle \langle n | x'' \rangle e^{-i\omega(n+\frac{1}{2})(t-t_0)} = \frac{e^{-\frac{i\omega\delta t}{2}}}{\sqrt{\pi x_0}} e^{-\frac{1}{2} \left( \frac{x''^2}{x_0^2} + \frac{x'^2}{x_0^2} \right)} \sum \frac{e^{-i\omega\delta t n}}{2^n n!} H_n\left(\frac{x''}{x_0}\right) H_n\left(\frac{x'}{x_0}\right). \quad (12)$$

Now, considering that, for  $\zeta = e^{-i\omega\delta t}$ ,  $\xi = \frac{x''}{x_0}$  and  $\eta = \frac{x'}{x_0}$

$$e^{-(\xi^2+\eta^2)} \sum \frac{\zeta^n}{2^n n!} H_n \xi H_n \eta = \frac{e^{-\frac{\xi^2+\eta^2-2\xi\eta\zeta}{1-\zeta^2}}}{\sqrt{i-\zeta^2}}, \quad (13)$$

we have

$$K = \frac{e^{-\frac{i\omega\delta t}{2}}}{\sqrt{\pi x_0}} e^{\frac{1}{2}(\xi^2+\eta^2)} e^{-\frac{\xi^2+\eta^2-2\xi\eta\zeta}{1-\zeta^2}} = \frac{e^{\left( -\frac{(\xi^2+\eta^2)e^{i\omega\delta t}-2\xi\eta}{e^{i\omega\delta t}-e^{-i\omega\delta t}} + \frac{\xi^2+\eta^2}{2} \right)}}{x_0 \sqrt{2\pi i \sin(\omega\delta t)}} = \sqrt{\frac{m\omega}{2\pi i \hbar \sin(\omega\delta t)}} e^{-\frac{(\xi^2+\eta^2)(\cos(\omega\delta t)+i\sin(\omega\delta t))-2\xi\eta-is(\xi^2+\eta^2)}{2i\sin(\omega\delta t)}} \quad (14)$$