```
============================================================
  PARAMETER MANIFEST
============================================================

  Model           : Hubbard-Holstein
  Ansatz          : warm:hh_hva_ptw -> adapt:PoolB(UCCSD+HVA+PAOP_FULL) -> final:hh_hva_ptw
  Drive enabled   : True

  t (hopping)     : 1.0
  U (interaction) : 2.0
  dv (disorder)   : 0.0

  L               : 2
  n_ph_max        : 1
  omega0          : 1.0
  g_ep            : 1.0
  noise_modes     : ['ideal']
  trotter_steps   : 8
  num_times       : 11
  magnus_variants : midpoint_magnus2,cfqm4,cfqm6
  hardcoded_family: included(static+drive)

  Command:
    /opt/anaconda3/bin/python /Users/jakestrobel/Documents/Holstein_implementation/Holstein_test/pipelines/exa
    ct_bench/hh_noise_robustness_seq_report.py --output-json /Users/jakestrobel/Documents/Holstein_implementat
    ion/Holstein_test/artifacts/json/hh_noise_robustness_L2_report.json --output-pdf
    '/Users/jakestrobel/Documents/Holstein_implementation/Holstein_test/docs/HH noise robustness report.PDF'
    --smoke-test-intentionally-weak --warm-reps 1 --warm-restarts 1 --warm-maxiter 20 --adapt-max-depth 3
    --adapt-maxiter 20 --final-reps 1 --final-restarts 1 --final-maxiter 20 --num-times 11 --trotter-steps 8
    --noise-modes ideal --oracle-repeats 1 --shots 256 --noisy-mode-timeout-s 60 --use-fake-backend

============================================================
```

```
SECTION: RESULTS SUMMARY

HH Noise Robustness Report Summary

final delta_abs: 0.10610594826147146
warm stop: Maximum number of function evaluations has been exceeded.
adapt stop: max_depth
final stop: Maximum number of function evaluations has been exceeded.
noisy modes completed: 2 / 2

Transition policies:
  warm->adapt: {'window_k': 5, 'slope_epsilon': 5e-05, 'patience': 3, 'min_points_before_switch': 8}
  adapt->vqe: {'window_k': 5, 'slope_epsilon': 5e-05, 'patience': 3, 'min_points_before_switch': 8}

Definitions:
  DeltaE_k = E_k - E_exact_sector
  switch when abs(slope( |DeltaE| window )) <= epsilon for patience checkpoints

Noisy-vs-noiseless policy:
  noisy uses Suzuki-2 only (ideal/shots/aer_noise); Magnus/CFQM shown as noiseless baselines.

SECTION: HARDCODED FAMILY REMAP
  legacy lane -> warm-start HVA seed
  paop lane   -> ADAPT Pool-B seed
  hva lane    -> final seeded conventional VQE
```

SECTION: RESULTS STAGE TRANSITIONS

Transition traces and Pool-B composition.



[PLOT_CAPTION] Stage switching: $|\Delta E|$ vs checkpoint for warm start and ADAPT.
Dashed trace is the windowed slope used by the plateau rule ($w = 5$, $\varepsilon = 5 \times 10^{-5}$, patience=3).

Pool B Composition (strict union + dedup)

Metric	Value
raw uccsd	3
raw hva	8
raw paop_full	9
dedup total	20
presence uccsd	3
presence hva	8
presence paop_full	9
overlap count	0

[PLOT_CAPTION] Stage switching: $|\Delta E|$ vs checkpoint for warm start and ADAPT.
Dashed trace is the windowed slope used by the plateau rule ($w = 5$, $\varepsilon = 5 \times 10^{-5}$, patience=3).

SECTION: RESULTS STATIC OVERLAYS

Static-profile overlays and hardcoded-family diagnostics.



[PLOT_CAPTION] Static: total occupancy n_i(t) heatmaps (sites i=0..L-1).
Panels: exact GS (left), exact evolution (middle), noiseless ADAPT-HVA + Suzuki-2 (right).



[PLOT_CAPTION] Static: spin-up occupancy n_up_i(t) heatmaps.
Panels: exact GS (left), exact evolution (middle), noiseless ADAPT-HVA + Suzuki-2 (right).



[PLOT_CAPTION] Static: spin-down occupancy n_dn_i(t) heatmaps.
Panels: exact GS (left), exact evolution (middle), noiseless ADAPT-HVA + Suzuki-2 (right).



[PLOT_CAPTION] Static (final branch): top F_sub, middle E_total, bottom doublon vs time.
Exact reference vs noiseless ADAPT-HVA + Suzuki-2.



[PLOT_CAPTION] Static: branch overlays - top E_total, bottom doublon (warm / ADAPT / final).
All curves are noiseless; exact reference shown for comparison.



[PLOT_CAPTION] Static: absolute scalar error channels heatmap (rows are |E|, |D|, and |S| error metrics). Columns run over time; brighter color means larger deviation from the matched exact channel.

generated 2026-03-04 04:28:25Z

SECTION: RESULTS DRIVE OVERLAYS

Drive-profile overlays and hardcoded-family diagnostics.



[PLOT_CAPTION] Drive: total occupancy n_i(t) heatmaps (sites i=0..L-1).
Panels: exact GS (left), exact evolution (middle), noiseless ADAPT-HVA + Suzuki-2 (right).



[PLOT_CAPTION] Drive: spin-up occupancy n_up_i(t) heatmaps.
Panels: exact GS (left), exact evolution (middle), noiseless ADAPT-HVA + Suzuki-2 (right).



[PLOT_CAPTION] Drive: spin-down occupancy n_dn_i(t) heatmaps.
Panels: exact GS (left), exact evolution (middle), noiseless ADAPT-HVA + Suzuki-2 (right).



[PLOT_CAPTION] Drive (final branch): top F_sub, middle E_total, bottom doublon vs time.
Exact reference vs noiseless ADAPT-HVA + Suzuki-2.



[PLOT_CAPTION] Drive: branch overlays - top E_total, bottom doublon (warm / ADAPT / final).
All curves are noiseless; exact reference shown for comparison.



[PLOT_CAPTION] Drive: absolute scalar error channels heatmap (rows are |E|, |D|, and |S| error metrics). Columns run over time; brighter color means larger deviation from the matched exact channel.

generated 2026-03-04 04:28:27Z

SECTION: RESULTS METHOD COMPARISON

Suzuki / Magnus / CFQM comparison overlays versus exact reference.

Legend key (applies to this section):
  Exact: exact reference evolution.
  Noiseless: ADAPT-HVA pipeline on noiseless simulator.
  Integrators: Suzuki-2, Magnus-2, CFQM4, CFQM6.

Noiseless methods overlay (static)



[PLOT_CAPTION] Static: noiseless integrator comparison - top E_total, bottom F_sub.
Methods: Suzuki-2 / Magnus-2 / CFQM4 / CFQM6; exact reference in black.

static: Magnus/CFQM error-to-exact



[PLOT_CAPTION] Static: integrator error vs exact - top |E-E_exact|, bottom (1-F_sub).
Smaller is better for ranking Suzuki-2 vs Magnus/CFQM baselines.

Noiseless methods overlay (drive)



[PLOT_CAPTION] Drive: noiseless integrator comparison - top E_total, bottom F_sub.
Methods: Suzuki-2 / Magnus-2 / CFQM4 / CFQM6; exact reference in black.

drive: Magnus/CFQM error-to-exact



[PLOT_CAPTION] Drive: integrator error vs exact - top |E-E_exact|, bottom (1-F_sub).
Smaller is better for ranking Suzuki-2 vs Magnus/CFQM baselines.

SECTION: RESULTS NOISE DETAILS

Overlay + per-mode detail pages for noise robustness.

Legend key (applies to this section):
  Exact: exact reference evolution.
  Noiseless: ADAPT-HVA pipeline on noiseless simulator.
  Noisy: shot-based estimator under selected mode (256 shots).
  Ideal-reference: baseline used for Delta(noisy-ideal) panels.

static: energy overlays



[PLOT_CAPTION] Static: energy vs time - top E_total, bottom E_static.
Exact vs noiseless (ADAPT-HVA+Suzuki-2) vs noisy (mode=ideal, 256 shots).

Static: site-0 occupation overlays



[PLOT_CAPTION] Static: site-0 occupations vs time - top $n_\uparrow(0)$, bottom $n_\downarrow(0)$. Noiseless vs noisy (mode=ideal, 256 shots).

static: doublon/staggered overlays



[PLOT_CAPTION] Static: correlators vs time - top doublon, bottom staggered order S. Noiseless vs noisy (mode=ideal, 256 shots).


Hubbard-Holstein dimer | noise=ideal | drive=on | ansatz=hva | energy
drive: energy overlays
exact reference
noiseless (Suzuki-2)
noise mode=ideal measured
E_total
E_static
time
[PLOT_CAPTION] Drive: energy vs time - top E_total, bottom E_static.
Exact vs noiseless (ADAPT-HVA+Suzuki-2) vs noisy (mode=ideal, 256 shots).
generated 2026-03-04 04:28:27Z

drive: site-0 occupation overlays



[PLOT_CAPTION] Drive: site-0 occupations vs time - top $n_\uparrow(0)$, bottom $n_\downarrow(0)$. Noiseless vs noisy (mode=ideal, 256 shots).

drive: doublon/staggered overlays



[PLOT_CAPTION] Drive: correlators vs time - top doublon, bottom staggered order S.
Noiseless vs noisy (mode=ideal, 256 shots).

static/ideal: noisy energy audit



[PLOT_CAPTION] Static / mode=ideal: top E_total, middle E_static, bottom Δ(measured-ideal-ref).
Shows estimator deviation from its internal ideal reference (not the ED exact curve).

static/ideal: site-0 occupation audit

[PLOT_CAPTION] Static / mode=ideal: top $n_\uparrow(0)$, middle $n_\downarrow(0)$, bottom Δ(measured-ideal-ref).
Use the Δ panel to isolate estimator/noise bias over time.

static/ideal: doublon & staggered audit

[PLOT_CAPTION] Static / mode=ideal: top doublon, middle staggered S, bottom Δ(measured-ideal-ref).
Use the Δ panel to isolate estimator/noise bias over time.

drive/ideal: noisy energy audit



[PLOT_CAPTION] Drive / mode=ideal: top E_total, middle E_static, bottom Δ(measured-ideal-ref).
Shows estimator deviation from its internal ideal reference (not the ED exact curve).

drive/ideal: site-0 occupation audit



[PLOT_CAPTION] Drive / mode=ideal: top $n_\uparrow(0)$, middle $n_\downarrow(0)$, bottom Δ(measured-ideal-ref).
Use the Δ panel to isolate estimator/noise bias over time.

drive/ideal: doublon & staggered audit



[PLOT_CAPTION] Drive / mode=ideal: top doublon, middle staggered S, bottom Δ(measured-ideal-ref).
Use the Δ panel to isolate estimator/noise bias over time.



[PLOT_CAPTION] Drive waveform v(t) evaluated on the simulation time grid.
Gaussian-envelope sinusoid used in H_drive(t)=sum_i,sigma v_i(t) n_i,sigma.

```
Diagnostics and availability

Comparisons keys: ['noiseless_vs_exact', 'noise_vs_noiseless']
Noisy mode diagnostics: {   "static:ideal": {     "success": true,     "env_blocked": false,     "reason": null,
    "error": null   },   "drive:ideal": {     "success": true,     "env_blocked": false,     "reason": null,
    "error": null   } }

Metric formulas:
  energy_static(t) = <psi(t)|H_static|psi(t)>
  energy_total(t)  = <psi(t)|H_static + H_drive(t)|psi(t)>
  n_up_site0 = <n_{0,up}> , n_dn_site0 = <n_{0,dn}>
  doublon(site0) = <n_{0,up} n_{0,dn}>
  staggered = (1/L) * Sum_j (-1)^j <n_j>

equation_registry_count: 90
plot_contract_count: 43
```

SECTION: APPENDIX DEFINITIONS USED

Short equation definitions used in result pages.

SECTION: APPENDIX DEFINITIONS USED

Formula index for this report.

[eq_cfqm]

$$U_{\mathrm{CFQM}}(\Delta t) \approx \prod_m \exp[-i\, a_m \Delta t\, H(t + c_m \Delta t)]$$

Meaning: Commutator-free Magnus stage product (CFQM4/CFQM6).

Units: unitary

[eq_delta_e]

$$\Delta E_k = E_k - E_{\mathrm{exact,\,sector}},\ \ \delta_k = |\Delta E_k|$$

Meaning: Stage transition error metrics.

Units: energy

[eq_doublon]

$$D_i = \langle n_{i\uparrow}\, n_{i\downarrow} \rangle$$

Meaning: Onsite doublon expectation.

Units: dimensionless

[eq_drive_waveform]

$$v_i(t) = s_i A \sin[\omega(t + t_0) + \phi] \exp\left(-\frac{(t + t_0)^2}{2\tilde{t}^2}\right)$$

Meaning: Gaussian-envelope sinusoidal drive with spatial weights s_i.

Units: energy

[eq_energy_static]

$$E^b_{\mathrm{static}}(t) = \langle \psi_b(t) | H_0 | \psi_b(t) \rangle$$

Meaning: Static-Hamiltonian energy channel for branch b.

Units: energy

[eq_energy_total]

$$E^b_{\mathrm{total}}(t) = \langle \psi_b(t) | H_0 + H_{\mathrm{drive}}(t) | \psi_b(t) \rangle$$

Meaning: Instantaneous total energy under drive.

Units: energy

[eq_error_abs]

$$\varepsilon_X(t) = |X_{\mathrm{approx}}(t) - X_{\mathrm{exact}}(t)|$$

Meaning: Absolute trajectory error metric.

Units: observable units

[eq_h_drive_density]

$$H_{\mathrm{drive}}(t) = \sum_{i,\sigma} v_i(t)\, n_{i\sigma}$$

Meaning: Drive couples to onsite electron number operators.

Units: energy

SECTION: APPENDIX DEFINITIONS USED

Formula index for this report.

[eq_magnus2]

$$U_{\mathrm{Magnus2}}(\Delta t) = \exp\left[-i\Delta t\, H\left(t + \frac{\Delta t}{2}\right)\right]$$

Meaning: Exponential midpoint (Magnus-2) reference approximation.

Units: unitary

[eq_noisy_delta]

$$\Delta_O^{\mathrm{noise-ideal}}(t) = \mu_O^{\mathrm{noise}}(t) - \mu_O^{\mathrm{ideal}}(t)$$

Meaning: Noisy-vs-ideal observable delta.

Units: observable units

[eq_noisy_estimator]

$$\mu_O = \mathrm{agg}(\{\langle O\rangle_r\}_{r=1}^{R}),\ \sigma_O = \mathrm{std}(\{\langle O\rangle_r\}_{r=1}^{R})$$

Meaning: Noisy oracle aggregate and empirical spread.

Units: observable units

[eq_number_operator]

$$n_{i\sigma} = \frac{I - Z_{q(i,\sigma)}}{2}$$

Meaning: JW number-operator identity used for occupancy observables.

Units: dimensionless

[eq_slope_switch]

$$\mathrm{switch} \Leftrightarrow |\mathrm{slope}(\delta_{k-w+1:k})| \le \varepsilon \text{ for } p \text{ consecutivecheckpoints}$$

Meaning: Windowed abs-slope plateau rule for stage transitions.

Units: energy/checkpoint

[eq_staggered]

$$S = \frac{1}{L}\sum_{i=0}^{L-1}(-1)^i\langle n_i\rangle$$

Meaning: Staggered charge-density order parameter.

Units: dimensionless

[eq_subspace_fidelity]

$$F_{\mathrm{sub}}^{b}(t) = \langle\psi_b^{\mathrm{trot}}(t)|P_{\mathrm{GS}}(t)|\psi_b^{\mathrm{trot}}(t)\rangle$$

Meaning: Projected fidelity against sector-filtered exact GS manifold.

Units: probability

[eq_suzuki2]

$$U_{\mathrm{S2}}(\Delta t) \approx \prod_j e^{-i\frac{\Delta t}{2}H_j}\prod_j^{\mathrm{rev}} e^{-i\frac{\Delta t}{2}H_j}$$

Meaning: Second-order symmetric Suzuki product step.

Units: unitary

SECTION: APPENDIX DEFINITIONS USED

Formula index for this report.

[eq_total_density]

$$n_i = \langle n_{i\uparrow} + n_{i\downarrow} \rangle$$

Meaning: Site-resolved total occupancy.

Units: particles

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_cfqm]

$$U_{\mathrm{CFQM}}(\Delta t) \approx \prod_m \exp[-i\, a_m \Delta t\, H(t + c_m \Delta t)]$$

Meaning: Commutator-free Magnus stage product (CFQM4/CFQM6).

Units: unitary

Source: dynamics_noiseless.profiles.*.methods.cfqm4, dynamics_noiseless.profiles.*.methods.cfqm6

[eq_delta_e]

$$\Delta E_k = E_k - E_{\mathrm{exact,\,sector}},\ \ \delta_k = |\Delta E_k|$$

Meaning: Stage transition error metrics.

Units: energy

Source: stage_pipeline.*.transition.delta_abs_trace

[eq_doublon]

$$D_i = \langle n_{i\uparrow} n_{i\downarrow} \rangle$$

Meaning: Onsite doublon expectation.

Units: dimensionless

Source: hardcoded_superset.profiles.*.trajectory.*.doublon_*

[eq_doublon_drive_adapt]

$$doublon_{drive}^{adapt}(t)$$

Meaning: doublon channel for drive profile, adapt branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

[eq_doublon_drive_final]

$$doublon_{drive}^{final}(t)$$

Meaning: doublon channel for drive profile, final branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

[eq_doublon_drive_warm]

$$doublon_{drive}^{warm}(t)$$

Meaning: doublon channel for drive profile, warm branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_doublon_static_adapt]

$$doublon_{static}^{adapt}(t)$$

Meaning: doublon channel for static profile, adapt branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_doublon_static_final]

$$doublon_{static}^{final}(t)$$

Meaning: doublon channel for static profile, final branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_doublon_static_warm]

$$doublon_{static}^{warm}(t)$$

Meaning: doublon channel for static profile, warm branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_drive_waveform]

$$v_i(t) = s_i A \sin[\omega(t + t_0) + \phi] \exp\left(-\frac{(t + t_0)^2}{2\bar{t}^2}\right)$$

Meaning: Gaussian-envelope sinusoidal drive with spatial weights s_i.

Units: energy

Source: settings.drive_profile.A, settings.drive_profile.omega, ... (+1 more)

[eq_energy_static]

$$E_{\text{static}}^{b}(t) = \langle \psi_b(t) | H_0 | \psi_b(t) \rangle$$

Meaning: Static-Hamiltonian energy channel for branch b.

Units: energy

Source: hardcoded_superset.profiles.*.trajectory.*.energy_static_*

[eq_energy_static_drive_adapt]

$$energy_static_{drive}^{adapt}(t)$$

Meaning: energy_static channel for drive profile, adapt branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_energy_static_drive_final]

$$energy_static_{drive}^{final}(t)$$

Meaning: energy_static channel for drive profile, final branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

[eq_energy_static_drive_warm]

$$energy_static_{drive}^{warm}(t)$$

Meaning: energy_static channel for drive profile, warm branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

[eq_energy_static_static_adapt]

$$energy_static_{static}^{adapt}(t)$$

Meaning: energy_static channel for static profile, adapt branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_energy_static_static_final]

$$energy_static_{static}^{final}(t)$$

Meaning: energy_static channel for static profile, final branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_energy_static_static_warm]

$$energy_static_{static}^{warm}(t)$$

Meaning: energy_static channel for static profile, warm branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_energy_total]

$$E_{\text{total}}^{b}(t) = \langle \psi_b(t) | H_0 + H_{\text{drive}}(t) | \psi_b(t) \rangle$$

Meaning: Instantaneous total energy under drive.

Units: energy

Source: hardcoded_superset.profiles.*.trajectory.*.energy_total_*

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_energy_total_drive_adapt]

$$energy_total_{drive}^{adapt}(t)$$

Meaning: energy_total channel for drive profile, adapt branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

[eq_energy_total_drive_final]

$$energy_total_{drive}^{final}(t)$$

Meaning: energy_total channel for drive profile, final branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

[eq_energy_total_drive_warm]

$$energy_total_{drive}^{warm}(t)$$

Meaning: energy_total channel for drive profile, warm branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

[eq_energy_total_static_adapt]

$$energy_total_{static}^{adapt}(t)$$

Meaning: energy_total channel for static profile, adapt branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_energy_total_static_final]

$$energy_total_{static}^{final}(t)$$

Meaning: energy_total channel for static profile, final branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_energy_total_static_warm]

$$energy_total_{static}^{warm}(t)$$

Meaning: energy_total channel for static profile, warm branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_error_abs]

$$\varepsilon_X(t) = |X_{\text{approx}}(t) - X_{\text{exact}}(t)|$$

Meaning: Absolute trajectory error metric.

Units: observable units

Source: comparisons.noiseless_vs_exact, comparisons.noise_vs_noiseless

[eq_h_drive_density]

$$H_{\text{drive}}(t) = \sum_{i,\sigma} v_i(t)\, n_{i\sigma}$$

Meaning: Drive couples to onsite electron number operators.

Units: energy

Source: settings.drive_profile, diagnostics.metric_definitions.energy_total

[eq_h_total]

$$H(t) = H_0 + H_{\text{drive}}(t)$$

Meaning: Total Hamiltonian splits into static + drive terms.

Units: energy

Source: settings.t, settings.u, ... (+2 more)

[eq_magnus2]

$$U_{\text{Magnus2}}(\Delta t) = \exp\left[-i\Delta t\, H\left(t + \tfrac{\Delta t}{2}\right)\right]$$

Meaning: Exponential midpoint (Magnus-2) reference approximation.

Units: unitary

Source: dynamics_noiseless.profiles.*.methods.magnus2

[eq_n_dn_site0_drive_adapt]

$$n_d n_s ite0_{drive}^{adapt}(t)$$

Meaning: n_dn_site0 channel for drive profile, adapt branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

[eq_n_dn_site0_drive_final]

$$n_d n_s ite0_{drive}^{final}(t)$$

Meaning: n_dn_site0 channel for drive profile, final branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_n_dn_site0_drive_warm]

$n_d n_s ite0_{drive}^{warm}(t)$

Meaning: n_dn_site0 channel for drive profile, warm branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

[eq_n_dn_site0_static_adapt]

$n_d n_s ite0_{static}^{adapt}(t)$

Meaning: n_dn_site0 channel for static profile, adapt branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_n_dn_site0_static_final]

$n_d n_s ite0_{static}^{final}(t)$

Meaning: n_dn_site0 channel for static profile, final branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_n_dn_site0_static_warm]

$n_d n_s ite0_{static}^{warm}(t)$

Meaning: n_dn_site0 channel for static profile, warm branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_n_up_site0_drive_adapt]

$n_u p_s ite0_{drive}^{adapt}(t)$

Meaning: n_up_site0 channel for drive profile, adapt branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

[eq_n_up_site0_drive_final]

$n_u p_s ite0_{drive}^{final}(t)$

Meaning: n_up_site0 channel for drive profile, final branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_n_up_site0_drive_warm]

$$n_u p_s ite0_{drive}^{warm}(t)$$

Meaning: n_up_site0 channel for drive profile, warm branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

[eq_n_up_site0_static_adapt]

$$n_u p_s ite0_{static}^{adapt}(t)$$

Meaning: n_up_site0 channel for static profile, adapt branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_n_up_site0_static_final]

$$n_u p_s ite0_{static}^{final}(t)$$

Meaning: n_up_site0 channel for static profile, final branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_n_up_site0_static_warm]

$$n_u p_s ite0_{static}^{warm}(t)$$

Meaning: n_up_site0 channel for static profile, warm branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_noisy_delta]

$$\Delta_O^{\mathrm{noise-ideal}}(t) = \mu_O^{\mathrm{noise}}(t) - \mu_O^{\mathrm{ideal}}(t)$$

Meaning: Noisy-vs-ideal observable delta.

Units: observable units

Source: dynamics_noisy.profiles.*.modes.*.trajectory.*.*_delta_noisy_minus_ideal

[eq_noisy_doublon_drive_aer_noise]

$$\mu_{doublon}^{aer_noise,\,drive}(t)$$

Meaning: Noisy estimate for doublon in drive/aer_noise.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.aer_noise.trajectory

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_noisy_doublon_drive_ideal]

$\mu_{doublon}^{ideal,\,drive}(t)$

Meaning: Noisy estimate for doublon in drive/ideal.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.ideal.trajectory

[eq_noisy_doublon_drive_shots]

$\mu_{doublon}^{shots,\,drive}(t)$

Meaning: Noisy estimate for doublon in drive/shots.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.shots.trajectory

[eq_noisy_doublon_static_aer_noise]

$\mu_{doublon}^{aer_noise,\,static}(t)$

Meaning: Noisy estimate for doublon in static/aer_noise.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.aer_noise.trajectory

[eq_noisy_doublon_static_ideal]

$\mu_{doublon}^{ideal,\,static}(t)$

Meaning: Noisy estimate for doublon in static/ideal.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.ideal.trajectory

[eq_noisy_doublon_static_shots]

$\mu_{doublon}^{shots,\,static}(t)$

Meaning: Noisy estimate for doublon in static/shots.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.shots.trajectory

[eq_noisy_energy_static_drive_aer_noise]

$\mu_{energy_static}^{aer_noise,\,drive}(t)$

Meaning: Noisy estimate for energy_static in drive/aer_noise.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.aer_noise.trajectory

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_noisy_energy_static_drive_ideal]

$\mu^{ideal,\,drive}_{energy_static}(t)$

Meaning: Noisy estimate for energy_static in drive/ideal.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.ideal.trajectory

[eq_noisy_energy_static_drive_shots]

$\mu^{shots,\,drive}_{energy_static}(t)$

Meaning: Noisy estimate for energy_static in drive/shots.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.shots.trajectory

[eq_noisy_energy_static_static_aer_noise]

$\mu^{aer_noise,\,static}_{energy_static}(t)$

Meaning: Noisy estimate for energy_static in static/aer_noise.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.aer_noise.trajectory

[eq_noisy_energy_static_static_ideal]

$\mu^{ideal,\,static}_{energy_static}(t)$

Meaning: Noisy estimate for energy_static in static/ideal.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.ideal.trajectory

[eq_noisy_energy_static_static_shots]

$\mu^{shots,\,static}_{energy_static}(t)$

Meaning: Noisy estimate for energy_static in static/shots.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.shots.trajectory

[eq_noisy_energy_total_drive_aer_noise]

$\mu^{aer_noise,\,drive}_{energy_total}(t)$

Meaning: Noisy estimate for energy_total in drive/aer_noise.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.aer_noise.trajectory

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_noisy_energy_total_drive_ideal]

$$\mu_{energy_total}^{ideal,\,drive}(t)$$

Meaning: Noisy estimate for energy_total in drive/ideal.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.ideal.trajectory

[eq_noisy_energy_total_drive_shots]

$$\mu_{energy_total}^{shots,\,drive}(t)$$

Meaning: Noisy estimate for energy_total in drive/shots.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.shots.trajectory

[eq_noisy_energy_total_static_aer_noise]

$$\mu_{energy_total}^{aer_noise,\,static}(t)$$

Meaning: Noisy estimate for energy_total in static/aer_noise.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.aer_noise.trajectory

[eq_noisy_energy_total_static_ideal]

$$\mu_{energy_total}^{ideal,\,static}(t)$$

Meaning: Noisy estimate for energy_total in static/ideal.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.ideal.trajectory

[eq_noisy_energy_total_static_shots]

$$\mu_{energy_total}^{shots,\,static}(t)$$

Meaning: Noisy estimate for energy_total in static/shots.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.shots.trajectory

[eq_noisy_estimator]

$$\mu_O = \mathrm{agg}(\{\langle O\rangle_r\}_{r=1}^{R}),\ \sigma_O = \mathrm{std}(\{\langle O\rangle_r\}_{r=1}^{R})$$

Meaning: Noisy oracle aggregate and empirical spread.

Units: observable units

Source: settings.oracle_repeats, settings.oracle_aggregate

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_noisy_n_dn_site0_drive_aer_noise]

$\mu_{n_d n_s ite0}^{aer_n oise,\, drive}(t)$

Meaning: Noisy estimate for n_dn_site0 in drive/aer_noise.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.aer_noise.trajectory

[eq_noisy_n_dn_site0_drive_ideal]

$\mu_{n_d n_s ite0}^{ideal,\, drive}(t)$

Meaning: Noisy estimate for n_dn_site0 in drive/ideal.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.ideal.trajectory

[eq_noisy_n_dn_site0_drive_shots]

$\mu_{n_d n_s ite0}^{shots,\, drive}(t)$

Meaning: Noisy estimate for n_dn_site0 in drive/shots.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.shots.trajectory

[eq_noisy_n_dn_site0_static_aer_noise]

$\mu_{n_d n_s ite0}^{aer_n oise,\, static}(t)$

Meaning: Noisy estimate for n_dn_site0 in static/aer_noise.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.aer_noise.trajectory

[eq_noisy_n_dn_site0_static_ideal]

$\mu_{n_d n_s ite0}^{ideal,\, static}(t)$

Meaning: Noisy estimate for n_dn_site0 in static/ideal.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.ideal.trajectory

[eq_noisy_n_dn_site0_static_shots]

$\mu_{n_d n_s ite0}^{shots,\, static}(t)$

Meaning: Noisy estimate for n_dn_site0 in static/shots.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.shots.trajectory

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_noisy_n_up_site0_drive_aer_noise]

$\mu_{n_u p_s ite0}^{aer_n oise,\, drive}(t)$

Meaning: Noisy estimate for n_up_site0 in drive/aer_noise.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.aer_noise.trajectory

[eq_noisy_n_up_site0_drive_ideal]

$\mu_{n_u p_s ite0}^{ideal,\, drive}(t)$

Meaning: Noisy estimate for n_up_site0 in drive/ideal.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.ideal.trajectory

[eq_noisy_n_up_site0_drive_shots]

$\mu_{n_u p_s ite0}^{shots,\, drive}(t)$

Meaning: Noisy estimate for n_up_site0 in drive/shots.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.shots.trajectory

[eq_noisy_n_up_site0_static_aer_noise]

$\mu_{n_u p_s ite0}^{aer_n oise,\, static}(t)$

Meaning: Noisy estimate for n_up_site0 in static/aer_noise.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.aer_noise.trajectory

[eq_noisy_n_up_site0_static_ideal]

$\mu_{n_u p_s ite0}^{ideal,\, static}(t)$

Meaning: Noisy estimate for n_up_site0 in static/ideal.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.ideal.trajectory

[eq_noisy_n_up_site0_static_shots]

$\mu_{n_u p_s ite0}^{shots,\, static}(t)$

Meaning: Noisy estimate for n_up_site0 in static/shots.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.shots.trajectory

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_noisy_staggered_drive_aer_noise]

$\mu_{staggered}^{aer_noise,\,drive}(t)$

Meaning: Noisy estimate for staggered in drive/aer_noise.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.aer_noise.trajectory

[eq_noisy_staggered_drive_ideal]

$\mu_{staggered}^{ideal,\,drive}(t)$

Meaning: Noisy estimate for staggered in drive/ideal.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.ideal.trajectory

[eq_noisy_staggered_drive_shots]

$\mu_{staggered}^{shots,\,drive}(t)$

Meaning: Noisy estimate for staggered in drive/shots.

Units: observable units

Source: dynamics_noisy.profiles.drive.modes.shots.trajectory

[eq_noisy_staggered_static_aer_noise]

$\mu_{staggered}^{aer_noise,\,static}(t)$

Meaning: Noisy estimate for staggered in static/aer_noise.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.aer_noise.trajectory

[eq_noisy_staggered_static_ideal]

$\mu_{staggered}^{ideal,\,static}(t)$

Meaning: Noisy estimate for staggered in static/ideal.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.ideal.trajectory

[eq_noisy_staggered_static_shots]

$\mu_{staggered}^{shots,\,static}(t)$

Meaning: Noisy estimate for staggered in static/shots.

Units: observable units

Source: dynamics_noisy.profiles.static.modes.shots.trajectory

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_number_operator]

$$n_{i\sigma} = \frac{I - Z_{q(i,\sigma)}}{2}$$

Meaning: JW number-operator identity used for occupancy observables.

Units: dimensionless

Source: diagnostics.metric_definitions

[eq_slope_switch]

$$\text{switch} \Leftrightarrow |\text{slope}(\delta_{k-w+1:k})| \leq \varepsilon \text{ for } p \text{ consecutivecheckpoints}$$

Meaning: Windowed abs-slope plateau rule for stage transitions.

Units: energy/checkpoint

Source: settings.transition_policy

[eq_staggered]

$$S = \frac{1}{L} \sum_{i=0}^{L-1} (-1)^i \langle n_i \rangle$$

Meaning: Staggered charge-density order parameter.

Units: dimensionless

Source: hardcoded_superset.profiles.*.trajectory.*.staggered_*

[eq_staggered_drive_adapt]

$$staggered_{drive}^{adapt}(t)$$

Meaning: staggered channel for drive profile, adapt branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

[eq_staggered_drive_final]

$$staggered_{drive}^{final}(t)$$

Meaning: staggered channel for drive profile, final branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

[eq_staggered_drive_warm]

$$staggered_{drive}^{warm}(t)$$

Meaning: staggered channel for drive profile, warm branch.

Units: observable units

Source: hardcoded_superset.profiles.drive.trajectory

SECTION: APPENDIX FORMULA ATLAS

Formula index for this report.

[eq_staggered_static_adapt]

$$staggered_{static}^{adapt}(t)$$

Meaning: staggered channel for static profile, adapt branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_staggered_static_final]

$$staggered_{static}^{final}(t)$$

Meaning: staggered channel for static profile, final branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_staggered_static_warm]

$$staggered_{static}^{warm}(t)$$

Meaning: staggered channel for static profile, warm branch.

Units: observable units

Source: hardcoded_superset.profiles.static.trajectory

[eq_subspace_fidelity]

$$F_{\mathrm{sub}}^{b}(t) = \langle \psi_b^{\mathrm{trot}}(t) | P_{\mathrm{GS}}(t) | \psi_b^{\mathrm{trot}}(t) \rangle$$

Meaning: Projected fidelity against sector-filtered exact GS manifold.

Units: probability

Source: hardcoded_superset.profiles.*.trajectory.*.fidelity*

[eq_suzuki2]

$$U_{\mathrm{S2}}(\Delta t) \approx \prod_j e^{-i\frac{\Delta t}{2}H_j} \prod_j^{\mathrm{rev}} e^{-i\frac{\Delta t}{2}H_j}$$

Meaning: Second-order symmetric Suzuki product step.

Units: unitary

Source: dynamics_noiseless.profiles.*.methods.suzuki2

[eq_total_density]

$$n_i = \langle n_{i\uparrow} + n_{i\downarrow} \rangle$$

Meaning: Site-resolved total occupancy.

Units: particles

Source: hardcoded_superset.profiles.*.trajectory.*.n_site_*

SECTION: APPENDIX PLOT CONTRACTS

Full machine-readable plot contract map.

```
SECTION: APPENDIX PLOT CONTRACTS

[plot_drive_aer_noise_doublon_staggered]
  x: time grid
  y: ['doublon_noisy', 'doublon_ideal', 'staggered_noisy', 'staggered_ideal']
  source: ['dynamics_noisy.profiles.drive.modes.aer_noise.trajectory']
  notes: Noisy doublon/staggered channels and deltas.

[plot_drive_aer_noise_energy]
  x: time grid
  y: ['energy_static_noisy', 'energy_static_ideal', 'energy_total_noisy', 'energy_total_ideal']
  source: ['dynamics_noisy.profiles.drive.modes.aer_noise.trajectory']
  notes: Noisy energy channels and noisy-ideal deltas.

[plot_drive_aer_noise_occupancy]
  x: time grid
  y: ['n_up_site0_noisy', 'n_up_site0_ideal', 'n_dn_site0_noisy', 'n_dn_site0_ideal']
  source: ['dynamics_noisy.profiles.drive.modes.aer_noise.trajectory']
  notes: Noisy site-0 spin channels and deltas.

[plot_drive_branch_overlay_doublon_staggered]
  x: time grid
  y: ['doublon_trotter', 'doublon_trotter_paop', 'doublon_trotter_hva', 'staggered_trotter', 'staggered_trotter_paop',
    'staggered_trotter_hva']
  source: ['hardcoded_superset.profiles.drive.trajectory']
  notes: Three-branch mapped doublon/staggered overlay.

[plot_drive_branch_overlay_energy]
  x: time grid
  y: ['energy_total_trotter', 'energy_total_trotter_paop', 'energy_total_trotter_hva', 'energy_total_exact']
  source: ['hardcoded_superset.profiles.drive.trajectory']
  notes: Three-branch mapped overlay (warm/adapt/final).

[plot_drive_density_dn_heatmaps]
  x: time grid
  y: ['n_dn_site_exact', 'n_dn_site_exact_hva', 'n_dn_site_trotter_hva']
  source: ['hardcoded_superset.profiles.drive.trajectory']
  notes: Spin-down density family.
```

```
[plot_drive_density_total_heatmaps]
  x: time grid
  y: ['n_site_exact', 'n_site_exact_hva', 'n_site_trotter_hva']
  source: ['hardcoded_superset.profiles.drive.trajectory']
  notes: Heatmaps: exact GS, exact final branch, trotter final branch.

[plot_drive_density_up_heatmaps]
  x: time grid
  y: ['n_up_site_exact', 'n_up_site_exact_hva', 'n_up_site_trotter_hva']
  source: ['hardcoded_superset.profiles.drive.trajectory']
  notes: Spin-up density family.
```

SECTION: APPENDIX PLOT CONTRACTS (cont. 8)

```
[plot_drive_energy_fidelity_final]
  x: time grid
  y: ['fidelity_hva_trotter', 'energy_total_exact', 'energy_total_trotter_hva', 'doublon_exact_hva', 'doublon_trotter_hva']
  source: ['hardcoded_superset.profiles.drive.trajectory']
  notes: Final-branch hardcoded family summary.

[plot_drive_ideal_doublon_staggered]
  x: time grid
  y: ['doublon_noisy', 'doublon_ideal', 'staggered_noisy', 'staggered_ideal']
  source: ['dynamics_noisy.profiles.drive.modes.ideal.trajectory']
  notes: Noisy doublon/staggered channels and deltas.

[plot_drive_ideal_energy]
  x: time grid
  y: ['energy_static_noisy', 'energy_static_ideal', 'energy_total_noisy', 'energy_total_ideal']
  source: ['dynamics_noisy.profiles.drive.modes.ideal.trajectory']
  notes: Noisy energy channels and noisy-ideal deltas.

[plot_drive_ideal_occupancy]
  x: time grid
  y: ['n_up_site0_noisy', 'n_up_site0_ideal', 'n_dn_site0_noisy', 'n_dn_site0_ideal']
  source: ['dynamics_noisy.profiles.drive.modes.ideal.trajectory']
  notes: Noisy site-0 spin channels and deltas.

[plot_drive_magnus_cfqm_error]
  x: time grid
  y: ['|E_method-E_exact|', '|F_method-1|']
  source: ['dynamics_noiseless.profiles.drive.methods.*.trajectory']
  notes: Noiseless error channels vs exact.

[plot_drive_magnus_cfqm_overlay]
  x: time grid
  y: ['suzuki2.energy_total_trotter', 'magnus2.energy_total_trotter', 'cfqm4.energy_total_trotter', 'cfqm6.energy_total_trotter',
    'exact.energy_total_exact']
  source: ['dynamics_noiseless.profiles.drive.methods.*.trajectory']
  notes: Noiseless method comparison.
```

```
[plot_drive_overlay_doublon_staggered]
  x: time grid
  y: ['doublon', 'staggered overlays']
  source: ['dynamics_noisy.profiles.drive.modes.*.trajectory', 'dynamics_noiseless.profiles.drive.methods.suzuki2.trajectory']
  notes: Overlay page for doublon and staggered order.

[plot_drive_overlay_energy]
  x: time grid
  y: ['noiseless+suzuki2+exact+noisy energy overlays']
  source: ['dynamics_noisy.profiles.drive.modes.*.trajectory', 'dynamics_noiseless.profiles.drive.methods.suzuki2.trajectory']
  notes: Overlay page for exact/noiseless/noisy energies.
```

SECTION: APPENDIX PLOT CONTRACTS (cont. 16)

```
[plot_drive_overlay_occupancy]
  x: time grid
  y: ['n_up(0)', 'n_dn(0) overlays']
  source: ['dynamics_noisy.profiles.drive.modes.*.trajectory', 'dynamics_noiseless.profiles.drive.methods.suzuki2.trajectory']
  notes: Overlay page for site-0 occupancies.

[plot_drive_scalar_error_heatmap]
  x: time grid
  y: ['|E_*| rows', '|D_*| rows', '|S_*| rows']
  source: ['hardcoded_superset.profiles.drive.trajectory']
  notes: Absolute scalar error channels heatmap.

[plot_drive_shots_doublon_staggered]
  x: time grid
  y: ['doublon_noisy', 'doublon_ideal', 'staggered_noisy', 'staggered_ideal']
  source: ['dynamics_noisy.profiles.drive.modes.shots.trajectory']
  notes: Noisy doublon/staggered channels and deltas.

[plot_drive_shots_energy]
  x: time grid
  y: ['energy_static_noisy', 'energy_static_ideal', 'energy_total_noisy', 'energy_total_ideal']
  source: ['dynamics_noisy.profiles.drive.modes.shots.trajectory']
  notes: Noisy energy channels and noisy-ideal deltas.

[plot_drive_shots_occupancy]
  x: time grid
  y: ['n_up_site0_noisy', 'n_up_site0_ideal', 'n_dn_site0_noisy', 'n_dn_site0_ideal']
  source: ['dynamics_noisy.profiles.drive.modes.shots.trajectory']
  notes: Noisy site-0 spin channels and deltas.

[plot_stage_transition]
  x: checkpoint index
  y: ['delta_abs_trace', 'slope_trace']
  source: ['stage_pipeline.warm_start.transition', 'stage_pipeline.adapt_pool_b.transition']
  notes: Windowed abs-slope switching traces.

[plot_static_aer_noise_doublon_staggered]
```

```
  x: time grid
  y: ['doublon_noisy', 'doublon_ideal', 'staggered_noisy', 'staggered_ideal']
  source: ['dynamics_noisy.profiles.static.modes.aer_noise.trajectory']
  notes: Noisy doublon/staggered channels and deltas.

[plot_static_aer_noise_energy]
  x: time grid
  y: ['energy_static_noisy', 'energy_static_ideal', 'energy_total_noisy', 'energy_total_ideal']
  source: ['dynamics_noisy.profiles.static.modes.aer_noise.trajectory']
  notes: Noisy energy channels and noisy-ideal deltas.
```

SECTION: APPENDIX PLOT CONTRACTS (cont. 24)

```
[plot_static_aer_noise_occupancy]
  x: time grid
  y: ['n_up_site0_noisy', 'n_up_site0_ideal', 'n_dn_site0_noisy', 'n_dn_site0_ideal']
  source: ['dynamics_noisy.profiles.static.modes.aer_noise.trajectory']
  notes: Noisy site-0 spin channels and deltas.

[plot_static_branch_overlay_doublon_staggered]
  x: time grid
  y: ['doublon_trotter', 'doublon_trotter_paop', 'doublon_trotter_hva', 'staggered_trotter', 'staggered_trotter_paop',
    'staggered_trotter_hva']
  source: ['hardcoded_superset.profiles.static.trajectory']
  notes: Three-branch mapped doublon/staggered overlay.

[plot_static_branch_overlay_energy]
  x: time grid
  y: ['energy_total_trotter', 'energy_total_trotter_paop', 'energy_total_trotter_hva', 'energy_total_exact']
  source: ['hardcoded_superset.profiles.static.trajectory']
  notes: Three-branch mapped overlay (warm/adapt/final).

[plot_static_density_dn_heatmaps]
  x: time grid
  y: ['n_dn_site_exact', 'n_dn_site_exact_hva', 'n_dn_site_trotter_hva']
  source: ['hardcoded_superset.profiles.static.trajectory']
  notes: Spin-down density family.

[plot_static_density_total_heatmaps]
  x: time grid
  y: ['n_site_exact', 'n_site_exact_hva', 'n_site_trotter_hva']
  source: ['hardcoded_superset.profiles.static.trajectory']
  notes: Heatmaps: exact GS, exact final branch, trotter final branch.

[plot_static_density_up_heatmaps]
  x: time grid
  y: ['n_up_site_exact', 'n_up_site_exact_hva', 'n_up_site_trotter_hva']
  source: ['hardcoded_superset.profiles.static.trajectory']
  notes: Spin-up density family.
```

```
[plot_static_energy_fidelity_final]
  x: time grid
  y: ['fidelity_hva_trotter', 'energy_total_exact', 'energy_total_trotter_hva', 'doublon_exact_hva', 'doublon_trotter_hva']
  source: ['hardcoded_superset.profiles.static.trajectory']
  notes: Final-branch hardcoded family summary.

[plot_static_ideal_doublon_staggered]
  x: time grid
  y: ['doublon_noisy', 'doublon_ideal', 'staggered_noisy', 'staggered_ideal']
  source: ['dynamics_noisy.profiles.static.modes.ideal.trajectory']
  notes: Noisy doublon/staggered channels and deltas.
```

SECTION: APPENDIX PLOT CONTRACTS (cont. 32)

```
[plot_static_ideal_energy]
  x: time grid
  y: ['energy_static_noisy', 'energy_static_ideal', 'energy_total_noisy', 'energy_total_ideal']
  source: ['dynamics_noisy.profiles.static.modes.ideal.trajectory']
  notes: Noisy energy channels and noisy-ideal deltas.

[plot_static_ideal_occupancy]
  x: time grid
  y: ['n_up_site0_noisy', 'n_up_site0_ideal', 'n_dn_site0_noisy', 'n_dn_site0_ideal']
  source: ['dynamics_noisy.profiles.static.modes.ideal.trajectory']
  notes: Noisy site-0 spin channels and deltas.

[plot_static_magnus_cfqm_error]
  x: time grid
  y: ['|E_method-E_exact|', '|F_method-1|']
  source: ['dynamics_noiseless.profiles.static.methods.*.trajectory']
  notes: Noiseless error channels vs exact.

[plot_static_magnus_cfqm_overlay]
  x: time grid
  y: ['suzuki2.energy_total_trotter', 'magnus2.energy_total_trotter', 'cfqm4.energy_total_trotter', 'cfqm6.energy_total_trotter',
    'exact.energy_total_exact']
  source: ['dynamics_noiseless.profiles.static.methods.*.trajectory']
  notes: Noiseless method comparison.

[plot_static_overlay_doublon_staggered]
  x: time grid
  y: ['doublon', 'staggered overlays']
  source: ['dynamics_noisy.profiles.static.modes.*.trajectory', 'dynamics_noiseless.profiles.static.methods.suzuki2.trajectory']
  notes: Overlay page for doublon and staggered order.

[plot_static_overlay_energy]
  x: time grid
  y: ['noiseless+suzuki2+exact+noisy energy overlays']
  source: ['dynamics_noisy.profiles.static.modes.*.trajectory', 'dynamics_noiseless.profiles.static.methods.suzuki2.trajectory']
  notes: Overlay page for exact/noiseless/noisy energies.
```

```
[plot_static_overlay_occupancy]
  x: time grid
  y: ['n_up(0)', 'n_dn(0) overlays']
  source: ['dynamics_noisy.profiles.static.modes.*.trajectory', 'dynamics_noiseless.profiles.static.methods.suzuki2.trajectory']
  notes: Overlay page for site-0 occupancies.

[plot_static_scalar_error_heatmap]
  x: time grid
  y: ['|E_*| rows', '|D_*| rows', '|S_*| rows']
  source: ['hardcoded_superset.profiles.static.trajectory']
  notes: Absolute scalar error channels heatmap.
```

SECTION: APPENDIX PLOT CONTRACTS (cont. 40)

```
[plot_static_shots_doublon_staggered]
  x: time grid
  y: ['doublon_noisy', 'doublon_ideal', 'staggered_noisy', 'staggered_ideal']
  source: ['dynamics_noisy.profiles.static.modes.shots.trajectory']
  notes: Noisy doublon/staggered channels and deltas.

[plot_static_shots_energy]
  x: time grid
  y: ['energy_static_noisy', 'energy_static_ideal', 'energy_total_noisy', 'energy_total_ideal']
  source: ['dynamics_noisy.profiles.static.modes.shots.trajectory']
  notes: Noisy energy channels and noisy-ideal deltas.

[plot_static_shots_occupancy]
  x: time grid
  y: ['n_up_site0_noisy', 'n_up_site0_ideal', 'n_dn_site0_noisy', 'n_dn_site0_ideal']
  source: ['dynamics_noisy.profiles.static.modes.shots.trajectory']
  notes: Noisy site-0 spin channels and deltas.
```

Executed Command

Reference: pipelines/run_guide.md
Script: pipelines/exact_bench/hh_noise_robustness_seq_report.py

```
/opt/anaconda3/bin/python /Users/jakestrobel/Documents/Holstein_implementation/Holstein_test/pipelines/exact_b
    ench/hh_noise_robustness_seq_report.py --output-json /Users/jakestrobel/Documents/Holstein_implementation/
    Holstein_test/artifacts/json/hh_noise_robustness_L2_report.json --output-pdf
    '/Users/jakestrobel/Documents/Holstein_implementation/Holstein_test/docs/HH noise robustness report.PDF'
    --smoke-test-intentionally-weak --warm-reps 1 --warm-restarts 1 --warm-maxiter 20 --adapt-max-depth 3
    --adapt-maxiter 20 --final-reps 1 --final-restarts 1 --final-maxiter 20 --num-times 11 --trotter-steps 8
    --noise-modes ideal --oracle-repeats 1 --shots 256 --noisy-mode-timeout-s 60 --use-fake-backend
```