

Can ENSO predict dengue outbreaks? A causal-based analysis on the role of climate patterns in *Aedes*-borne disease transmission

Javier Corvillo^{1, 2*}, Verónica Torralba^{2*}, Diego Campos², Ana Riviére-Cinnamond³, and Ángel G. Muñoz⁴

¹Complutense University of Madrid, Department of Earth Science and Astrophysics, Madrid, 28040, Spain

²Barcelona Supercomputing Center, Earth Sciences Department, 08034, Spain

³Pan-American Health Organization, Communicable Diseases and Health Analysis, Panama City, 0843-03441, Panama

⁴International Centre for Theoretical Physics, Trieste, 34151, Italy

*javier.corvillo@bsc.es / veronica.torralba@bsc.es / angel.g.munoz@bsc.es

ABSTRACT

Background: Vector-borne diseases transmitted by *Aedes* mosquitoes such as dengue, Zika, and chikungunya, pose significant public health challenges worldwide in the wake of human-driven climate change. However, while their transmission is known to be susceptible to some climate variables like temperature, rainfall or humidity, the overall role of climate patterns on the emergence of these diseases is not so well understood.

Methods: Using data from a number of sources, we explore and analyse the response of *Aedes*-borne disease transmission to climate patterns. Our analysis is composed of three different studies: 1) a timescale decomposition of disease transmissibility values, thereby guiding officials to understand the behaviour of outbreaks for budget and resource allocation; 2) a correlation analysis between transmissibility values and different seasonal climate variability modes, such as El Niño Southern Oscillation, in order to check for the influence of natural climate patterns onto *Aedes*-borne outbreaks; and 3) a causality analysis to solidify findings obtained through correlation, identifying the most relevant predictors for *Aedes*-borne diseases.

Results: Long-term, man-made climate change is shown to have a significant impact on the environmental suitability for *Aedes*-borne diseases in the tropics, where El Niño Southern Oscillation and the Indian Ocean Basin are key climate patterns conditioning disease emergence. Temperate regions are shown to be more susceptible to seasonal climate variability, where multiscale climate patterns, through teleconnections and compound interactions, can influence transmission dynamics.

Conclusions: Disease transmission of *Aedes*-borne diseases is shown to be susceptible to multiple factors, including long-term climate change and seasonal variability. The results of this study highlight the multi-faceted role of climate patterns in disease emergence, and their potential applicability in the development of early warning systems for *Aedes*-borne disease outbreaks. Future research should focus on integrating these findings into an actionable *Aedes*-borne seasonal prediction system using climate patterns as predictors, which can ultimately better inform public health strategies to manage dengue outbreaks.

Keywords: Public Health, Vector-borne Diseases, Epidemiology, Climate Change, Climate Services, Environmental Sciences

Multilingual abstracts

Please consult the [Additional files](#) section for abstract translations into the other five official languages of the United Nations (Arabic, Chinese, French, Russian and Spanish).

1 Background

1.1 The emergence of vector-borne diseases in the context of climate change

Disease stability and transmissibility under changing climate conditions has long been a topic of interest in the fields of epidemiology and virology. Many viral, bacterial, and

parasitic diseases have been shown to be reactive to changes in environmental conditions across different regions and timescales^{1,2}. This is particularly true for previously pandemic diseases that have now become endemic after public health officials implement proper disease control mechanisms³. Prevalent respiratory viruses, such as H1N1 influenza or the novel SARS-CoV-2 virus, have a reduced dependency on human spread once losing their pandemic status, adopting climate-dependent seasonal patterns and becoming more prominent in temperate climates during winter^{4,5}.

In the case of vector-borne diseases (VBDs), many known socio-economic factors shape up the expanse of pathogens^{6–8}, but the relationship between climate and disease transmissibility is even more intertwined. Carriers such as arthropods, snails and slugs thrive under specific climate thresholds, and in the context of anthropogenic climate change, global warming and changing precipitation patterns affect the overall spatial distribution of these vectors^{9,10}.

Mosquitoes of the *Aedes* genus, are of importance in medium and low income countries¹¹. As the main carriers of diseases like dengue (DENV), chikungunya (CHIKV), and Zika (ZIKV), *Aedes albopictus* and *Aedes aegypti* pose a significant public health threat in tropical and subtropical regions¹². In the Americas, for instance, DENV impacts account for over 2 million disability adjusted life years worldwide¹³, and low estimated annual prevention costs of about US\$2.1 billion¹⁴.

The effects of climate change are causing *Aedes*-related diseases not only to emerge in new, previously unaffected regions, but also to increase their spread in previously endemic areas¹⁵. Along with compound changes in urbanization¹⁶ and population growth¹⁷, climate change is posited to be a major driver of increased DENV incidence in temperate climates¹⁸, with the recent establishment of epidemic activity in parts of North America¹⁹, Southeast Asia²⁰, and detection of local transmission in southern European countries along the Mediterranean Basin²¹. Equatorial tropical and subtropical zones like the Sub-Saharan Africa and northern South America have also been subject to higher incidence over the past 40 years²². While the current yearly incidence of DENV amounts to an average of 400 million cases per year²³, it is believed that the effects of climate change may put an additional 2.5 billion people at risk of DENV if *Aedes* vectors were to be present in every region where climate conditions are suitable for their development²².

1.2 Known drivers on *Aedes*-borne disease transmission dynamics and their impact

The present scientific literature highlights a profound influence of the environmental conditions over *Aedes* mosquito proliferation, with temperature and precipitation being the two known primary drivers of *Aedes*-borne disease transmission, due to their impact on both vector and pathogen biology alike. Many vector physiological processes (e.g., biting frequency, reproduction rates) as well as pathogen characteristics (e.g., extrinsic incubation duration) are conditioned by temperature, increasing until certain temperature thresholds are reached²⁴. In a similar manner, rainfall promotes the availability of mosquito breeding sites, up until a critical point in which excessive rainfall may flood or wash out the mosquito larvae away²⁵. In conjunction, temperature and rainfall modulate the length of the transmission season of DENV, CHIKV

and ZIKV in temperate areas and affected hotspots, with higher incidence in urban areas of the Western Pacific and the Eastern Mediterranean regions²⁶.

Aside from these climate variables, humidity is also another probable factor in the proliferation of *Aedes* mosquito breeding sites. Evidence suggests that elevated humidity levels could play a role in reducing incubation periods and blood-feeding cycle duration in *Aedes* mosquitoes²⁷.

1.3 Climate patterns as an aggregate of unknown transmissibility drivers

While macro climatic factors have been widely used in *Aedes*-borne disease research and modelling^{28–31}, it is acknowledged that these are not enough to fully explain the climate component of *Aedes*-borne disease transmissibility, and that more explanatory variables may aid in refining DENV monitoring and prediction systems^{32–35}. Temperature and rainfall may also affect other possible small-scale climate modulators such as soil moisture or vegetation growth, which in turn may affect the availability of breeding sites for *Aedes* mosquitoes. Additionally, a high-resolution analysis of satellite imagery has suggested that land cover change, through deforestation, as well as stagnant water bodies, may promote *Aedes* breeding sites³⁶.

From a pragmatic perspective, understanding the contribution of each of these additional climate variables, each operating in different spatial and temporal scales, can be a challenging undertaking. Many of these variables lack comprehensive long-term observational records precisely in tropical and southern regions where *Aedes* diseases are most prevalent^{7,37}. Computational constraints may also arise when attempting to model the multiple interactions between variables, which, along with the high number of potential variables, makes it difficult to determine which combinations are most relevant for disease prediction without extensive resources. To add to this complexity, the effects of these variables may not be immediately apparent, as variables like soil moisture, for instance, do depend on many sub-processes as well³⁸.

Climate patterns, in this context, may serve a better role in mapping out the underlying climate processes behind *Aedes*-borne disease transmission. Climate patterns are recurring large-scale phenomena that can influence weather and climate conditions over different timescales (subseasonal, seasonal, inter-annual, decadal, etc.). Often characterized by their periodicity, they are identified through defined climate variability indices, and their effects ripple through the climate system, affecting temperature, precipitation, and other climate variables across different regions through teleconnections³⁹.

As such, climate patterns may serve as aggregators of both large and small climate phenomena by linking various

climate oscillations to regional weather events and other environmental processes⁴⁰. Since they encapsulate broader trends that affect local conditions, climate patterns seem preferable over essential climate variables in the development of future climate-and-health *Aedes* prediction systems^{41,42}. The analysis of seasonal patterns, such as the El Niño Southern Oscillation (ENSO), could provide insight as to how climate system interactions influence precipitation, temperature, and ultimately, *Aedes*-disease transmission and emergence.

1.4 The applicability of climate services in vector-borne disease prevention

Given the worldwide impacts of DENV, ZIKV and CHIKV and their expected potential to spread to unprepared regions, the need for effective *Aedes*-borne disease control is pressing. Many climate-and-health initiatives have been developed in recent years in response, with the aim of providing actionable information to public health officials and decision makers. The Tropical Disease Research initiative from the World Health Organization's (WHO)^{43,44}, as well as its Global Strategic Preparedness, Readiness and Response Plan, with \$55 million in funding from different stakeholders and public officials, are among the most notable. Other examples include the mapping of environmental suitability for DENV, ZIKV and CHIKV^{2,45}, or the development of integrated surveillance systems for *Aedes* arboviruses⁴⁶.

The development of early warning systems (EWS) for *Aedes* vectors is another area of focus in which climate information translates to more actionable health information⁴⁷. While disease EWS can be based on clinical and epidemiological data alone, integration of climate information can improve the accuracy and skill of these systems, while simultaneously extending how far in advance changes in transmissibility can be detected^{48,49}.

Here, we aim to characterise the time variability of the climate-driven component of *Aedes*-borne transmissibility, to understand how the frequency of the different climate fluctuations may influence DENV outbreaks in different regions and seasons in a climate change context. Additionally, we aim to understand the link between of climate patterns and *Aedes*-borne disease emergence. This link can be exploited for the implementation of efficient operational vector-borne disease control programs as these climate patterns are suitable to act as predictors under a possible causal-based outbreak prediction system.

2 Methods

2.1 R_0 as a bridge between climate and health

Three main mechanisms have been discussed to explain the seasonality of VBDs⁵⁰: the effects of the vector's behaviour (absence or presence in the area), the effects of human behaviour (whether an infected host can transmit the

disease by travelling), and the effects of climate (whether the conditions are favourable for the vector to transmit the disease). These three mechanisms are encapsulated in the basic reproduction number (R_0), an epidemiological metric that quantifies transmissibility of VBDs, defined as the average number of secondary cases generated by a single infected individual in a susceptible population⁵¹.

However, while R_0 entangles vector, human, and climate behaviour, by only integrating the climate component of the disease transmission for the computation of R_0 , then the resulting R_0 metric is more so interpreted as the role of environmental conditions in the spread of the disease. Thus, this definition of R_0 , which we can understand as the diseases' environmental suitability, acts as a first approximation of the dynamics of VBDs. This tweaked definition of R_0 is one of the operational outbreak indices commonly used by the WHO, along with several other decision-making institutes and health practitioners in disease control and prevention^{52,53}.

2.2 The *Aedes* Disease Environmental Suitability 2's monitoring system

The *Aedes* Disease Environmental Suitability 2's (AeDES2) monitoring system⁴⁵ was used to obtain the R_0 values for the analysis. Improving over its predecessor², AeDES2 is a climate-and-health service that provides real-time monitoring of the environmental suitability for *Aedes*-borne diseases. The system uses disease climate drivers like temperature and precipitation, and through its integration with entomological models and calibration with recorded DENV cases, R_0 values are computed for different regions and seasons. The monitoring system is designed to be used by public health officials and researchers to assess the risk of disease outbreaks and to inform prevention and control strategies.

2.3 Analysis 1: Multi-timescale climate decomposition of R_0

A timescale decomposition methodology was used to isolate the signal of R_0 data into several time components, in order to obtain the total variance across different timescales. This approach allows us to understand how different climate processes at different timescales condition disease transmissibility, which is essential for the development of effective public health strategies, optimization of resource allocation, or implementation of targeted intervention strategies^{54,55}.

2.3.1 Data

The timescale decomposition analysis was undertaken using R_0 outputs from the AeDES2's monitoring system. The 1951-2024 monthly-mean period of AeDES2's R_0 values was selected for the analysis, for a total of 888 months or 295 full seasons. Considering that vector borne diseases are extending to previously unaffected areas due to the effects of man-made climate change, AeDES2's coverage has been increased since its inception to contain global outputs, allowing for a more




Figure 1. Overview of the climate variability indices and their areas of influence.

Index Name	Abbreviation	Periodicity	Pattern Type
Atlantic 3 Index	ATL3	Several months to a few years	Oceanic
Indian Ocean Basin Mode	IOBM	Several months to a few years	Oceanic
Niño3.4 Index	Niño 3.4	Between 2-7 years	Oceanic
North Pacific Meridional Mode	NPMM	Several months to a few years	Atmospheric
South Atlantic Subtropical Dipole	SASD	Several months to a few years	Oceanic
South Pacific Meridional Mode	SPMM	Several months to a few years	Atmospheric
Tropical North Atlantic	TNA	Several months to a few years	Oceanic

Table 1. Summary of the climate variability indices used in the analysis used for the correlation and causality studies.

comprehensive analysis of the relationship between climate variability indices and R_0 in emerging *Aedes* hotspots.

2.3.2 Methodology

As R_0 doesn't follow a defined probability distribution function, the timescale decomposition analysis filters a given R_0 time-series of any given grid-point by employing the non-parametric locally estimated scatterplot smoothing technique (LOESS)⁵⁶. In order to obtain the best LOESS smoothing parameter for the analysis, we select the value that provides the lowest Generalized Correlated Cross-Validation verification metric (GCV) for the goodness of fit of the model⁵⁷. GCV was chosen amongst different metrics, as it provides a robust assessment of the LOESS model by penalizing overfitting through its leave-one-out validation procedure⁵⁸.

Once the ideal LOESS smoothing parameter is found for the R_0 data, the R_0 time-series for each grid-point is separated into four components: a long-term trend signal (understood to be the trend caused by anthropogenic climate change), a seasonal signal (over-the-year change), a decadal signal (10-30 years), and a residual signal which contains other signals of the data (i.e., inter-annual and inter-decadal variability, among others). Variance maps for each of these

four components capture the overall direction and variability of the data over time in any given grid-point.

After the variance maps are obtained, the trend of R_0 values is removed for following correlation and causality analyses. While detrending through the assumption that R_0 changes linearly over time could be a valid approach, it fails to capture the temperature dependency of the data. In order to account for this intrinsic relationship of R_0 with temperature, a similar timescale decomposition analysis is performed on the detrended R_0 data, though instead of time as the independent variable, temperature data from the AeDES2's monitoring system datasets is used instead (monthly-mean temperature data consisting of the GHCN-CAMS project⁵⁹, the CPC Unified Global Temperature dataset⁶⁰, and the ERA5 and the ERA5Land reanalysis datasets⁶¹). In this way, the obtained trend serves as a functional relationship between temperature and R_0 : a temperature-based description of the R_0 signal, attributed to the warming of the planet⁶².

2.4 Analysis 2: Correlation studies between R_0 and climate variability indices

After analyzing R_0 variability, we assess the impact of several climate patterns on global R_0 values over the chosen 1951-

2024 monthly-mean period.

2.4.1 Data

Correlation studies are performed over global outputs, using the temperature-based detrended R_0 data from the previous analysis over the different seasons. A total of 7 temperature-based climate variability indices have been used for the correlation analysis, computed using the detrended temperature data of Section 2.3.2. Their periodicity and pattern type, as well as area of influence, is summarized and represented in Table 1 and Figure 1.

2.4.2 Methodology

The correlation analysis was performed using the Spearman partial correlation coefficient, which assesses how well an arbitrary monotonic function can describe the relationship between two variables⁶³. For computation of statistical significance in correlation, the non-parametric Monte Carlo method was used⁶⁴, with a p-value threshold of 0.05.

2.5 Analysis 3: Causality studies between R_0 and climate variability indices. Outlining of predictors for disease outbreaks

Causal-based patterns can be identified after this analysis, which allow for another robust foundation for understanding of the underlying mechanisms between climate variability and R_0 patterns. Along with results obtained through correlation, this causality analysis can be used to outline the most relevant predictors for disease outbreaks.

2.5.1 Data

The datasets that were used for the causality analysis are the same detrended datasets as those employed in Section 2.4.1.

2.5.2 Methodology

Causality analyses between R_0 and climate variability indices were performed by using Liang-Kleeman's proposed methodology for computing information flow between two entities of a dynamical system, quantifying the amount of information that one time series (the climate variability indices) can provide about another time series (R_0 patterns)⁶⁵. Once the transfer entropy is computed, it is then normalized in order to account for the different scales of the two time series⁶⁶. Statistical significance is computed using Fisher's information matrix, with a p-value threshold of 0.05.

Both sign and value of correlation and causality outputs give information about the relationship between R_0 and the climate variability indices, making the two analyses non-exclusive and complementary to each other. Positive correlation values indicate that R_0 and the climate pattern index are positively correlated; R_0 increases (decreases) when the climate pattern is in its positive (negative) phase. Alternatively, negative correlation values indicate that R_0 and the climate variability index are negatively correlated; R_0 increases (decreases) when the climate pattern is in its negative (positive) phase.

In the case of causality, positive Liang-Kleeman causality values indicate that variations in any given climate pattern tend to amplify or destabilize fluctuations over R_0 (either favorably or unfavorably), leading to large changes in transmission potential. Alternatively, negative causality values suggest that variations in the climate index tend to dampen or stabilize fluctuations over R_0 , acting as a regulator for disease transmission and bringing R_0 toward more moderate values.

3 Results

3.1 Analysis 1: Multi-timescale decomposition of R_0

Timescale decomposition maps for R_0 global outputs for the whole 1951-2024 monthly-mean time series can be found in Figure 2, along with the component time series for the cities of Iquitos, Peru; and Santa Fe, Argentina, known to be traditional hotspots of *Aedes*-borne diseases. The optimal span values of LOESS smoothing for the timescale decomposition methodology for this sample period was 8 months for the GCV verification metric. It is worth noting that, because the R_0 metric operates in certain temperature ranges, grid points with less than 6 months of data per year were excluded from these and following analyses, in order to perform these analyses with the more robust data.

The long-term trend and residual components of R_0 (Figures 2a and 2d) show the highest variance contribution in tropical and subtropical regions. The sum of these components rounds between 40-80% particularly across central Africa, parts of South America including the Amazon basin, and scattered areas in Southeast Asia and northern Australia. These patterns suggest long-term climate change, along with interannual climate variability, to drive systematic shifts in R_0 in these endemic regions, reflecting the complex interactions between temperature, humidity, and precipitation influencing vector populations and virus transmission. For these regions, there is no clearly defined transmission season over the year, as the seasonal signal of R_0 (Figure 2b and 2e) is rather poor. Instead, the climatology of R_0 in these regions, due to high temperature and humidity values, adopts persistent high R_0 values throughout the year, independently of the season.

Though the decadal component (Figure 2c) shows a negligible contribution across all regions of the globe <5%, the seasonal variance shows a markedly different geographical distribution, with the highest contributions concentrated in regions with pronounced seasonal temperature variations (between 60-100% variance explained, Figure 2b). Notable hotspots include the northern edges of traditional DENV transmission zones in South America, parts of southern Africa, central Asia, and temperate regions that border current endemic areas. It is in these regions where the seasonal temperature fluctuations strongly condition the transmissibil-




Figure 2. a)-d) Global timescale decomposition maps for R_0 values for the monthly mean 1951-2024 period, different components. The time series for the cities of Iquitos, Peru (blue dot), and Santa Fe, Argentina (red dot), are shown as examples in e) and f) respectively.

ity season of R_0 , reaching its peak around the summer months.

Indeed, comparing Figures 2e and 2f highlights the effect of the seasonal component on the overall R_0 dynamics in these strong seasonal regions. While the trend component accounts for a significant portion of the variance, it is the seasonal component that can play a crucial role in defining whether a DENV outbreak can occur ($R_0 > 1$) or not ($R_0 < 1$).

Applying the timescale decomposition methodology for two climatological subsets allows to see how the different

components have evolved overtime. The climatologies corresponding to the 1961-1990 and 1991-2020 periods can be found in Figure 3, along with their respective Köppen-Geiger climate classification maps⁶⁷. For both of these two periods, the GCV-optimal span value of LOESS smoothing is also 8 months, same as for the complete time series. The decadal component is not shown in this figure, as it remains negligible (<5%) for both climatologies.

When examining the variance patterns in relation to the Köppen-Geiger classification, a visual inspection of the maps reveals distinct climate zone signatures. Trend and




Figure 3. a)-c) Global timescale decomposition maps for R_0 values for the monthly mean 1961-1990 period, different components. Its associated Köppen-Geiger climate classification is shown in g). d)-f) Global timescale decomposition maps for R_0 values for the monthly mean 1991-2020 period, different components. Its associated Köppen-Geiger climate classification is shown in h). (Source for Köppen-Geiger climate classification maps:⁶⁷)

residual variance typically dominates in tropical climate zones (A-type climates in the Amazon basin, central Africa, and Southeast Asia), while temperate and humid subtropical zones (C-type climates in southern Brazil, parts of East Asia, and the Mediterranean basin) show a more pronounced seasonal component. At a glance, it may seem that a study on the expansion of A and C type Köppen Geiger climates in the context of climate change may provide a useful framework for predicting the evolution of *Aedes*-borne transmissibility, with each major climate type exhibiting distinct characteristics that condition the behavior of the R_0 signal.

However, this would be a naive approach, since the Köppen-Geiger classification is a static classification system that does not thoroughly reflect the dynamic nature of climate change⁶⁸. Indeed, the two 30-year periods reveal shifts in how climate variability affects R_0 . In the 1961-1990 period, the trend component shows more concentrated high-variance

zones, particularly in central Africa and tropical parts of South America. By the later 1991-2020 period however, these trend patterns have intensified in some regions while diminishing in others, highlighting that the impacts of systematic climate change on transmission potential, while shown to have become more pronounced over time, are not entirely straightforward.

Given the strong seasonality of R_0 in temperate regions through these timescale decomposition analyses, and given the strong relationship between temperature and R_0 , it is not unreasonable to expect R_0 values to be closely related to seasonal, temperature-driven climate patterns. The following analyses aim to explore these relationships further, with the seasonal, temperature-driven climate variability indices previously listed in Table 1.




Figure 4. a)-d) Correlation values between R_0 and the Niño3.4 index, for the 1951-2024 monthly mean period in all seasons. Dots represent statistically significant correlation values, at $p < 0.01$.

3.2 Analysis 2: Correlation studies between R_0 and climate variability indices

Spearman partial correlation outputs between R_0 and the Niño3.4 index for all seasons of the 1951-2024 monthly-mean time series are shown in Figure 4. The most consistent positive correlations across seasons appear in the Amazon basin and northeastern South America, tropical A-type climate regions in the Köppen Geiger classification from section 3.1. The austral summer and autumn seasons (figures 4a and 4b) show the highest positive correlation values in those regions, with Africa exhibiting more heterogeneous patterns with both positive and negative correlation zones in the austral autumn. It is during these months that we also observe notable emergence of negative correlation values in parts of Southeast Asia, along with positive correlation values in some temperate transition areas as well.

Over the austral winter (Figure 4c), the correlation patterns shift in the Southern Hemisphere. Fewer regions, with Africa in particular, show predominantly weaker correlation values. Negative correlations emerge over the Sahel region, India and the East Coast of the United States. The austral spring (Figure 4d) demonstrates a return to stronger correlation patterns, particularly in South America where positive correlations

re-intensify across the Amazon and southern Brazil regions.

An aggregate of the maximum statistically significant correlation values between R_0 and all climate variability indices, as well as the index that gives the maximum correlation value, is shown in Figure 5. The left column shows the maximum correlation values for each index, while the right column shows which index gives the maximum correlation value for each grid point. The percentage of grid points over which each index gives the highest statistically significant correlation values is shown in Figure 6. It is worth noting that, since the difference between correlation values can often be small, the second and third highest correlation attribution percentages are also shown in the figure, in order to provide a more complete picture of the relative importance of each index.

Over the austral summer and autumn months (Figure 5a-b) the Niño3.4 index is the most dominant index in the regions previously highlighted in Figure 4a-b. However, as the Niño3.4 correlation with R_0 weakens during the austral winter and spring months over its zones of influence (Figure 4c-d), other climate indices are vying for dominance in its stead (Figure 5c-d, right). This behavior is also replicated in the regions of more seasonal R_0 variance described in section




Figure 5. a)-d) For all seasons: Left, maximum statistically significant correlation values between R_0 and all indexes of Table 1, for the 1951-2024 monthly mean period ($p < 0.01$). Right, which climate index gives the maximum statistically significant correlation values.




Figure 6. a)-d) For all seasons: Percentage of grid points over which each index gives the highest, second or third highest statistically significant correlation values. Any contribution above 10% is labeled in the barplots.

3.1, where the spread of the indices over which maximum correlation is attributed to becomes more heterogeneous (Figure 5a-d, right). Notable areas include the Sahel region, Mediterranean basin, parts of southern Africa, Australia, the East Coast of the United States and southeastern Asia, where different indices dominate the correlation landscape.

It is therefore evident that multiple seasonal temperature-based climate indices compete for dominance across different regions and seasons, with no single index controlling more than 40% of grid points as the primary correlate, in accordance to Figure 6. Still, other notable indices include the IOBM index, contributing more than 10% of grid points in nearly all categories and being prevalent in India during the austral winter and spring months. These findings and results for both El Niño and IOBM are consistent with previous studies that have highlighted the importance of both ENSO and the Indian Ocean in influencing climate variability and *Aedes*-borne disease transmission⁶⁹⁻⁷¹.

Despite their respective contributions however, the spread of all seven indices is relatively even, with no single index clearly dictating whether R_0 is directly tied to one specific climate pattern, reflecting the multifaceted nature of climate

variability influence on *Aedes*-borne disease transmission.

3.3 Analysis 3: Causality studies between R_0 and climate variability indices. Outlining of predictors for disease outbreaks

Analogue of figures 5 and 6 for section 3.3 are shown in Figures 7 and 8 for the Liang-Kleeman causality analysis. The causality analysis is performed over the same 1951-2024 monthly mean period, with the same climate variability indices of Table 1.

As opposed to the correlation outputs in Figure 5, the causality analyses in Figure 7 show a pronounced attenuation across all different regions and seasons, with the maximum causality values rarely exceeding values over ± 0.25 . Since causality is a more stringent condition than correlation, more diffuse and weaker patterns suggest that the direct causal influence of these climate patterns on transmission potential is more modest than simple correlation analysis would suggest. Additionally, since patterns are more spatially fragmented overall, direct causal climate influences on R_0 seem to operate through more localized mechanisms, instead of the more broader-scale patterns that correlation analyses might imply.




Figure 7. a)-d) For all seasons: Left, maximum statistically significant causality values between R_0 and all indexes of Table 1, for the 1951-2024 monthly mean period ($p < 0.01$). Right, which index gives the maximum statistically significant causality values.




Figure 8. a)-d) For all seasons: Percentage of grid points over which each index gives the highest, second or third highest statistically significant causality values. Any contribution above 10% is labeled in the barplots.

Nonetheless, the causality analysis still provides valuable insight into the relationships between R_0 and climate variability indices. Similarly to the correlation analyses, both the Niño3.4 index and IOBM index are the most dominant causal indices over the austral summer and autumn months over the Amazon basin, central Africa and southeastern regions of Asia (Figure 7a-b), with other indices gaining prominence in these regions during the austral winter and spring months (Figure 7c-d). Niño3.4 shows negative causality values over all seasons (causal and stable). Along with the IOBM index showing negative causality values over the spring and autumn months (Figure 7b, d), both of these climate patterns play a buffering effect, preventing extreme deviations on disease transmission values. On the other hand, the IOBM index shows positive causality values over the summer and winter months (causal but unstable, 7a, c), aligning with the seasonal maximum and minimum of the pattern. It is during these months of peak DENV transmission in both hemispheres that this climate pattern pushes already marginal transmission conditions toward more extreme states: either making cold regions less suitable for transmission, or amplifying transmission windows in suitable or unsuitable areas.

The influence of both Niño3.4 and IOBM indices on R_0 through causality relationships can be seen in Figure 8. While both indices do show a sizable contribution over all categories, similarly to correlation, there is no clear dominance of any index during any season, with the second and third highest causality values being relatively evenly distributed across the different indices. Still, a notable shift occurs during boreal autumn and spring months (8b, d), with ATL3 gaining increased prominence while maintaining the multi-index influence pattern. The NPMM and SASD1 indices maintain substantial influence, while ocean-based patterns show varying degrees of influence across regions with high seasonality (7b, d), suggesting that ocean basins become drivers of transmission variability at different times of the year in these areas.

4 Discussion

4.1 The added value of climate patterns in seasonal forecasting of *Aedes*-borne diseases

The climate system is complex and dynamic, with multiple components that shape both broad and local-scale phenomena interacting with each other at multiple timescales⁷². Accurately forecasting the effects of multiple climate modulators

and their reciprocal interactions was, is, and will continue to be a challenge for the scientific community, with new emerging technologies and methodologies pushing the boundaries of our understanding with time.

For vector-borne diseases, where climate is known to play a necessary - but not sufficient - role in disease transmission, the intertwining relationships between climate, socio-economic factors, and vector presence, makes accurately forecasting disease outbreaks an even more complex task. While many studies have provided academically sound evidence of the skillfulness of vector-borne disease forecasts based on climate information⁷³, public health application of these findings usually have come short in practice. Often times, these systems either lack data of sufficient quality, or don't integrate knowledge of how climate mechanisms drive disease transmission as a whole. Pragmatic policy implementation, often hindered by resource or capacity gaps, can be challenging or slower to materialize.

In choosing climate patterns as the focus of our analysis, we tackle the current technical challenges of climate-driven, vector-borne disease forecasting. Climate patterns provide a framework that can be used to forecast disease transmission more reliably than existing methods that use temperature and precipitation alone, which can be both inaccurate or unreliable in unskilled regions. The findings presented here are designed to, first, establish which variability timeframes are most important for DENV prediction in a global context, and second, identify the specific climate patterns that drive these relationships.

4.2 Analysis 1: Multi-timescale decomposition of R_0

Timescale decomposition results detailed in Section 3.1 reflect the overarching influence of climate drivers on the behavior of *Aedes*-borne emergence. Prior literature reflects that decadal-scale variability has had significant impacts on the climate overall, and hence populations and economies⁷⁴. In the context of DENV emergence, phenology models assess that the world became approximately 1.5% more suitable per decade for the development of *Aedes aegypti* during 1950-2000, and that this trend is predicted to accelerate to 3.2-4.4% per decade by 2050, a near tripling of the expansion rate⁷⁵. Yet, the low decadal variability results obtained in this manuscript, isolating the climate component from disease transmissibility, suggest that other socio-economic factors, such as mosquito migration patterns or population circulation and expansion, are facilitating expansion over the longer timescales.

It is certain that warming of the planet is the most prominent feature of climate change models⁷⁶, a certainty that bleeds into the emergence of DENV, ZIKV and CHIKV over temperate areas such as China or the Americas. Findings of Section 3.1 highlight this reality; while climate modulators

of the disease are most notorious over the seasonal timescale, shaped by climate patterns such as ENSO and the IOD, notable shifts are now causing long-term trends and intrayear signals to have a sizeable contribution in these and other tropical areas. The persistence of suitable environmental conditions across the whole year is in great part responsible for an alarming surge in DENV cases in 2023 and 2024 across endemic countries, facing significant strain from arboviral diseases⁷⁷.

Visual inspection of the different timescale decomposition results with climate classification zones are also consistent with the current literature. Both *Aedes aegypti* and *Aedes albopictus* transmission efficiency and population growth rates increase dramatically when areas transition to tropical conditions. Since the 1980s, anthropogenic accelerated global warming has fundamentally altered the geographic boundaries of climate zones⁷⁸. As tropical climate zones expand poleward, these shifts create new areas where these temperature thresholds are consistently met year-round, enabling permanent establishment for both carriers, rather than seasonal presence⁷⁹.

4.3 Analyses 2 and 3: Correlation and causality studies between R_0 and climate variability indices

Previous studies have highlighted the importance of both ENSO and the Indian Ocean in influencing climate variability and *Aedes*-borne disease transmission. For instance, it is well-known that ENSO-driven climate variability is responsible for periodic major outbreaks of dengue in Venezuela, with disease cycles of 1 and 3-4 years by virtue of ENSO-promoted temperature and rainfall patterns in the region⁷. The obtained positive correlation values for ENSO in this study link its positive phase with the emergence of DENV, while negative causality values portray the index as a stabilizer that keeps DENV transmission potential high, so long as the index is in a positive phase. These results for ENSO are consistent with observed real world events, such as the emergence of ZIKV in Latin America and the Caribbean in 2014-2016 during a period of severe drought and unusually high temperatures, conditions associated with the 2015-2016 El Niño event⁸⁰.

Indian Ocean climate variability modes such as IOBM or the Indian Ocean Dipole (IOD) have been posited to help predict long-term trends in *Aedes*-borne disease transmission through their effects on monsoon onset, duration and demise⁶⁹. Additionally, the Indian Ocean temperature anomalies have been shown to not only impact on health in the form of vector-borne diseases, but also on economic, migration, conflicts, and nutrition outcomes in low and middle income countries from the Indo-Pacific region⁸¹. For instance, a distributed lag nonlinear model revealed that the association between dengue incidence and IOD in

Bangladesh was comparatively stronger after adjustment for local climate variables, seasonality and trend⁷¹. Similarly to ENSO, the obtained positive correlation values for IOBM in this study link its positive phase with the emergence of DENV, though positive causality values portray the index as a destabilizer that can push transmission conditions toward even higher values. These results are in line with the ongoing CHIKV outbreak of ten thousand cases in the Guangdong province of China that started in June 2025; the phase of the IOBM index at the time of the outbreak was positive, and has remained positive for the following months.

Yet, to say that both ENSO and Indian Ocean variability patterns like IOBM are the only climate patterns that influence DENV transmission would be a gross oversimplification. Indeed, while the up-to-date literature mostly points to ENSO, and most recently the IOD, as climate drivers of *Aedes*-borne transmission, the results of Section 3.2 and 3.3 show that other temperature-based seasonal patterns gain prominence during the austral winter and spring months, especially over regions with a stronger seasonal transmission component. Multi-basin climate patterns such as NPMM, SPMM or SASD can modify rainfall patterns, thus influencing DENV transmission.

It is also necessary to state, in lieu of these results, that Figures 6 and 8 showing an even spread of the correlation and causality attribution percentages signals the interplay between all seven patterns modulating DENV transmission. For instance, it is known that the teleconnections between ENSO, IOBM and TNA plays a critical role in extreme drought events in the Amazon Basin⁸². Additionally, enhanced NPMM impact translates into an increased frequency of NPMM that leads to an extreme El Niño or La Niña events, which can have a significant impact on DENV transmission in the tropics under the right greenhouse gas emissions scenarios⁸³. All these differing relationships between climate patterns and DENV transmission can be both exclusive or complementary, and their combined influence should be considered in the development of *Aedes*-borne disease prediction systems.

4.4 Notable constraints and limitations

A number of caveats should be acknowledged when interpreting the results of this study. For one, while the results of this study, which are constrained to the climate-driven component of disease transmission, are indeed valuable, it needs to be emphasized that the interplay between climate and other socioeconomic factors is complex and context-dependent, and that climate, while crucial, is just one piece of the puzzle. In the context of the creation of a climate-driven DENV forecasting system using these results, health officials must take into account the multi-faceted nature of these drivers (mosquito presence, human circulation, climate conditions) in order for disease control programs to be more effective. Second, it is of note that the analysis is based on global

datasets. Applicability of these findings to specific DENV hotspots should be carefully considered, with the use of high-resolution observational data being encouraged to reproduce and build on these results in a local context.

Additionally, while these findings have been obtained with careful data handling in avoiding edge cases, correlation and causality analyses in regions where data is spurious or where the R_0 signal is weak may not yield reliable or realistic results (e.g., in temperate regions with low transmission potential). Computation for both analyses are reliant on the variance or covariance of the data available, and therefore, where there are few data points, these statistics become unstable and sensitive, leading to artificially large or small values.

Lastly, it is worth commenting on the non-exhaustive choice of climate patterns in Table 1. While the seven climate patterns that were selected do share similarities with one another (temperature-based seasonal patterns, with some of them being precursors of ENSO phases), other relevant seasonal patterns have been omitted from the analysis. Including more climate patterns is out of the scope of this study, though we encourage future studies to expand on the results presented here.

5 Conclusions

1. Historical reconstructions of climate-driven R_0 values for *Aedes*-borne diseases provide valuable insight in the role of climate in disease emergence, and can be used to improve the accuracy of seasonal DENV forecasts through the identification of predictors in the form of broad-scale climate patterns.
2. Long-term, man-made climate change is shown to have a significant impact on the environmental suitability for *Aedes*-borne diseases in the tropics, which comprise the Amazon basin, central Africa, and Southeast Asia. As the effects of climate change intensify, bordering regions that shift into A-type climate zones in the Köppen-Geiger classification may experience potential shifts in transmission patterns.
3. Tropical regions show a more stable but high R_0 signal throughout the year, whereas temperate regions exhibit a pronounced seasonal component, with peak transmission values occurring during the summer months.
4. Correlation and causality are non-exclusive, complementary analyses that can be used to understand the role of climate patterns in the dynamics of *Aedes*-borne diseases. While correlation exemplifies the strength and direction of relationships between variables, causality provides insight into its stability.
5. ENSO and IOBM are shown to be the most dominant climate patterns in regions where the long-term climate

signal is stronger (Amazon basin, central Africa and southeastern Asia). Other temperature-based seasonal patterns gain prominence during the austral winter and spring months, especially over regions with a stronger seasonal transmission component. Overall, no single climate pattern dominates the transmission landscape; it is the compound effects of multiple patterns that shape the dynamics of DENV outbreaks, and their combined influence should be considered in the development of Aedes-borne disease prediction systems.

6. In the tropics, periods of El Niño act as a promoter of *Aedes*-borne disease transmission, while periods of La Niña serve as a suppressor. Regardless of the phase however, ENSO is a stabilizing factor for DENV outbreaks in the tropics, preventing extreme deviations and keeping transmission potential high. IOBM shows similar behavior, though during peak DENV transmission months, it can push marginal transmission conditions toward more extreme states, either making cold regions less suitable for transmission or warming suitable or unsuitable areas to amplify transmission windows.

Given the above, future work should focus on the integration of high-resolution climate data and local-scale factors into seasonal DENV forecasting models. A framework for the development of such a model is currently being developed, to be reported in future publications.

Additional files

[Additional file 1](#): Multilingual abstracts in the six official working languages of the United Nations.

Abbreviations

AeDES2: *Aedes* Disease Environmental Suitability 2; AIC: Akaike Information Criterion; ATL3: Atlantic 3 Index; CHIKV: Chikungunya; DENV: Dengue; ENSO: El Niño Southern Oscillation; GCV: Generalized cross-validation; IOBM: Indian Ocean Basin; IOD: Indian Ocean Dipole; LOESS: Locally estimated scatterplot smoothing; NPM: North Pacific Meridional Mode; R^2 : Coefficient of determination; SASD: South Atlantic Subtropical Dipole; SPM: South Pacific Meridional Mode; TNA: Tropical North Atlantic; VB: Vector-borne diseases; WHO: World Health Organization; ZIKV: Zika.

Acknowledgements

(Blank until we figure coauthor contributions)

Funding

(Same here, blank for now)

Availability of data and materials

Code for the generation of R_0 values employed for this study, computed using AeDES2's monitoring system, as well as the necessary datasets, functions and scripts to generate the maps and plots for this manuscript are available under request.

Authors' contributions

Á.M., V.T. and D.C. conceived the methodology to be undertaken in this manuscript. Data sources, code and figures were obtained and developed from the ground up by J.C, who also analysed the results. All authors have reviewed the manuscript.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

References

1. Thomson, M. C., Garcia-Herrera, R. & Beniston, M. (eds.) *Seasonal Forecasts, Climatic Change and Human Health* (Springer Netherlands, Dordrecht, 2008).
2. Malloy, S. S., Horack, J. M., Lee, J. & Newton, E. K. Earth observation for public health: Biodiversity change and emerging disease surveillance. *Acta Astronaut.* **160**, 433–441, DOI: [10.1016/j.actaastro.2018.10.042](https://doi.org/10.1016/j.actaastro.2018.10.042) (2019).
3. Li, Y. *et al.* Global patterns in monthly activity of influenza virus, respiratory syncytial virus, parainfluenza virus, and metapneumovirus: A systematic analysis. *The Lancet. Glob. Heal.* **7**, e1031–e1045, DOI: [10.1016/S2214-109X\(19\)30264-5](https://doi.org/10.1016/S2214-109X(19)30264-5) (2019).
4. Shaman, J., Goldstein, E. & Lipsitch, M. Absolute humidity and pandemic versus epidemic influenza. *Am. J. Epidemiol.* **173**, 127–135, DOI: [10.1093/aje/kwq347](https://doi.org/10.1093/aje/kwq347) (2011).
5. Romero Starke, K. *et al.* The Effect of Ambient Environmental Conditions on COVID-19 Mortality: A Systematic Review. *Int. J. Environ. Res. Public Heal.* **18**, 6665, DOI: [10.3390/ijerph18126665](https://doi.org/10.3390/ijerph18126665) (2021).
6. Lippi, C. A. *et al.* The Social and Spatial Ecology of Dengue Presence and Burden during an Outbreak in Guayaquil, Ecuador, 2012. *Int. J. Environ. Res. Public Heal.* **15**, 827, DOI: [10.3390/ijerph15040827](https://doi.org/10.3390/ijerph15040827) (2018).
7. Ibarra, A. M. S. *et al.* Dengue Vector Dynamics (*Aedes aegypti*) Influenced by Climate and Social Factors in Ecuador: Implications for Targeted Control. *PLOS ONE* **8**, e78263, DOI: [10.1371/journal.pone.0078263](https://doi.org/10.1371/journal.pone.0078263) (2013).

8. Stewart-Ibarra, A. M. *et al.* Spatiotemporal clustering, climate periodicity, and social-ecological risk factors for dengue during an outbreak in Machala, Ecuador, in 2010. *BMC infectious diseases* **14**, 610, DOI: [10.1186/s12879-014-0610-4](https://doi.org/10.1186/s12879-014-0610-4) (2014).
9. Lowe, R. *et al.* Nonlinear and delayed impacts of climate on dengue risk in Barbados: A modelling study. *PLOS Medicine* **15**, e1002613, DOI: [10.1371/journal.pmed.1002613](https://doi.org/10.1371/journal.pmed.1002613) (2018).
10. Messina, J. P. *et al.* Mapping global environmental suitability for Zika virus. *eLife* **5**, e15272, DOI: [10.7554/eLife.15272](https://doi.org/10.7554/eLife.15272) (2016).
11. Campbell-Lendrum, D., Manga, L., Bagayoko, M. & Sommerfeld, J. Climate change and vector-borne diseases: What are the implications for public health research and policy? *Philos. Transactions Royal Soc. B: Biol. Sci.* **370**, 20130552, DOI: [10.1098/rstb.2013.0552](https://doi.org/10.1098/rstb.2013.0552) (2015).
12. Global Strategy for dengue prevention and control, 2012–2020. <https://www.who.int/publications/i/item/9789241504034>.
13. Yang, X., Quam, M. B. M., Zhang, T. & Sang, S. Global burden for dengue and the evolving pattern in the past 30 years. *J. Travel. Medicine* **28**, taab146, DOI: [10.1093/jtm/taab146](https://doi.org/10.1093/jtm/taab146) (2021).
14. Shepard, D. S., Coudeville, L., Halasa, Y. A., Zambrano, B. & Dayan, G. H. Economic Impact of Dengue Illness in the Americas. *The Am. J. Trop. Medicine Hyg.* **84**, 200–207, DOI: [10.4269/ajtmh.2011.10-0503](https://doi.org/10.4269/ajtmh.2011.10-0503) (2011).
15. Quam, M. B. *et al.* Estimating Air Travel–Associated Importations of Dengue Virus Into Italy. *J. Travel. Medicine* **22**, 186–193, DOI: [10.1111/jtm.12192](https://doi.org/10.1111/jtm.12192) (2015).
16. Lee, S. A., Economou, T., Catão, R. d. C., Barcellos, C. & Lowe, R. The impact of climate suitability, urbanisation, and connectivity on the expansion of dengue in 21st century Brazil. *PLOS Neglected Trop. Dis.* **15**, e0009773, DOI: [10.1371/journal.pntd.0009773](https://doi.org/10.1371/journal.pntd.0009773) (2021).
17. Struchiner, C. J., Rocklöv, J., Wilder-Smith, A. & Massad, E. Increasing Dengue Incidence in Singapore over the Past 40 Years: Population Growth, Climate and Mobility. *PLOS ONE* **10**, e0136286, DOI: [10.1371/journal.pone.0136286](https://doi.org/10.1371/journal.pone.0136286) (2015).
18. Kraemer, M. U. *et al.* The global distribution of the arbovirus vectors *Aedes aegypti* and *Ae. albopictus*. *eLife* **4**, e08347, DOI: [10.7554/eLife.08347](https://doi.org/10.7554/eLife.08347) (2015).
19. Franklins, L. H. V., Jones, K. E., Redding, D. W. & Abubakar, I. The effect of global change on mosquito-borne disease. *The Lancet Infect. Dis.* **19**, e302–e312, DOI: [10.1016/S1473-3099\(19\)30161-6](https://doi.org/10.1016/S1473-3099(19)30161-6) (2019).
20. Ooi, E.-E. & Gubler, D. J. Dengue in Southeast Asia: Epidemiological characteristics and strategic challenges in disease prevention. *Cadernos de Saúde Pública* **25**, S115–S124, DOI: [10.1590/S0102-311X2009001300011](https://doi.org/10.1590/S0102-311X2009001300011) (2009).
21. Dengue - Annual Epidemiological Report for 2022. <https://www.ecdc.europa.eu/en/publications-data/dengue-annual-epidemiological-report-2022> (2024).
22. Nakase, T., Giovanetti, M., Obolski, U. & Lourenço, J. Population at risk of dengue virus transmission has increased due to coupled climate factors and population growth. *Commun. Earth & Environ.* **5**, 1–11, DOI: [10.1038/s43247-024-01639-6](https://doi.org/10.1038/s43247-024-01639-6) (2024).
23. Pourzangiabadi, M., Najafi, H., Fallah, A., Goudarzi, A. & Pouladi, I. Dengue virus: Etiology, epidemiology, pathobiology, and developments in diagnosis and control – A comprehensive review. *Infect. Genet. Evol.* **127**, 105710, DOI: [10.1016/j.meegid.2024.105710](https://doi.org/10.1016/j.meegid.2024.105710) (2025).
24. Mordecai, E. A. *et al.* Thermal biology of mosquito-borne disease. *Ecol. Lett.* **22**, 1690–1708, DOI: [10.1111/ele.13335](https://doi.org/10.1111/ele.13335) (2019).
25. Paaijmans, K. P., Wandago, M. O., Githeko, A. K. & Takken, W. Unexpected High Losses of *Anopheles gambiae* Larvae Due to Rainfall. *PLOS ONE* **2**, e1146, DOI: [10.1371/journal.pone.0001146](https://doi.org/10.1371/journal.pone.0001146) (2007).
26. Colón-González, F. J. *et al.* Projecting the risk of mosquito-borne diseases in a warmer and more populated world: A multi-model, multi-scenario intercomparison modelling study. *The Lancet Planet. Heal.* **5**, e404–e414, DOI: [10.1016/S2542-5196\(21\)00132-7](https://doi.org/10.1016/S2542-5196(21)00132-7) (2021).
27. Descoux, E. *et al.* Climate-Based Models for Understanding and Forecasting Dengue Epidemics. *PLOS Neglected Trop. Dis.* **6**, e1470, DOI: [10.1371/journal.pntd.0001470](https://doi.org/10.1371/journal.pntd.0001470) (2012).
28. Caminade, C. *et al.* Global risk model for vector-borne transmission of Zika virus reveals the role of El Niño 2015. *Proc. Natl. Acad. Sci.* **114**, 119–124, DOI: [10.1073/pnas.1614303114](https://doi.org/10.1073/pnas.1614303114) (2017).
29. Liu-Helmersson, J., Stenlund, H., Wilder-Smith, A. & Rocklöv, J. Vectorial Capacity of *Aedes aegypti*: Effects of Temperature and Implications for Global Dengue Epidemic Potential. *PLOS ONE* **9**, e89783, DOI: [10.1371/journal.pone.0089783](https://doi.org/10.1371/journal.pone.0089783) (2014).
30. Mordecai, E. A. *et al.* Detecting the impact of temperature on transmission of Zika, dengue, and chikungunya using mechanistic models. *PLOS Neglected Trop. Dis.* **11**, e0005568, DOI: [10.1371/journal.pntd.0005568](https://doi.org/10.1371/journal.pntd.0005568) (2017).
31. Wesolowski, A. *et al.* Impact of human mobility on the emergence of dengue epidemics in Pakistan. *Proc. Natl. Acad. Sci.* **112**, 11887–11892, DOI: [10.1073/pnas.1504964112](https://doi.org/10.1073/pnas.1504964112) (2015).
32. Erraguntla, M. *et al.* Predictive model for microclimatic temperature and its use in mosquito population

- modeling. *Sci. Reports* **11**, 18909, DOI: [10.1038/s41598-021-98316-x](https://doi.org/10.1038/s41598-021-98316-x) (2021).
33. Lee, H. S. *et al.* Seasonal patterns of dengue fever and associated climate factors in 4 provinces in Vietnam from 1994 to 2013. *BMC Infect. Dis.* **17**, 218, DOI: [10.1186/s12879-017-2326-8](https://doi.org/10.1186/s12879-017-2326-8) (2017).
34. Yavari Nejad, F. & Varathan, K. D. Identification of significant climatic risk factors and machine learning models in dengue outbreak prediction. *BMC Med. Informatics Decis. Mak.* **21**, 141, DOI: [10.1186/s12911-021-01493-y](https://doi.org/10.1186/s12911-021-01493-y) (2021).
35. Sriklin, T., Kajornkasirat, S. & Puttinaovarat, S. Dengue Transmission Mapping with Weather-Based Predictive Model in Three Southernmost Provinces of Thailand. *Sustainability* **13**, 6754, DOI: [10.3390/su13126754](https://doi.org/10.3390/su13126754) (2021).
36. Ali, H. & Charles, H. Mosquito Breeding Hotspots in a Changing World: Remote Sensing and GIS-Based Analysis of Ecological and Anthropogenic Factors. Tech. Rep. (2025). DOI: [10.13140/RG.2.2.29354.94406](https://doi.org/10.13140/RG.2.2.29354.94406).
37. Sen Roy, S. Climate Change in the Global South: Trends and Spatial Patterns. In Sen Roy, S. (ed.) *Linking Gender to Climate Change Impacts in the Global South*, 1–25, DOI: [10.1007/978-3-319-75777-3_1](https://doi.org/10.1007/978-3-319-75777-3_1) (Springer International Publishing, Cham, 2018).
38. Peng, C., Zeng, J., Chen, K.-S., Ma, H. & Bi, H. Spatiotemporal Patterns and Influencing Factors Of Soil Moisture At A Global Scale. In *IGARSS 2023 - 2023 IEEE International Geoscience and Remote Sensing Symposium*, 3174–3177, DOI: [10.1109/IGARSS52108.2023.10282096](https://doi.org/10.1109/IGARSS52108.2023.10282096) (2023).
39. Lübbecke, J. F. *et al.* Equatorial Atlantic variability—Modes, mechanisms, and global teleconnections. *WIREs Clim. Chang.* **9**, e527, DOI: [10.1002/wcc.527](https://doi.org/10.1002/wcc.527) (2018).
40. Lee, D., Ward, P. & Block, P. Attribution of Large-Scale Climate Patterns to Seasonal Peak-Flow and Prospects for Prediction Globally. *Water Resour. Res.* **54**, 916–938, DOI: [10.1002/2017WR021205](https://doi.org/10.1002/2017WR021205) (2018).
41. Easterbrook, D. J. Chapter 21 - Using Patterns of Recurring Climate Cycles to Predict Future Climate Changes. In Easterbrook, D. J. (ed.) *Evidence-Based Climate Science (Second Edition)*, 395–411, DOI: [10.1016/B978-0-12-804588-6.00021-5](https://doi.org/10.1016/B978-0-12-804588-6.00021-5) (Elsevier, 2016).
42. Hallett, T. B. *et al.* Why large-scale climate indices seem to predict ecological processes better than local weather. *Nature* **430**, 71–75, DOI: [10.1038/nature02708](https://doi.org/10.1038/nature02708) (2004).
43. Keating, C. *Tore Godal and the Evolution of Global Health* (Routledge, New York, 2023), 1 edn.
44. Ramirez, B. & TDR-IDRC Research Initiative on Vector Borne Diseases and Climate Change. Support for research towards understanding the population health vulnerabilities to vector-borne diseases: Increasing resilience under climate change conditions in Africa. *Infect. Dis. Poverty* **6**, 164, DOI: [10.1186/s40249-017-0378-z](https://doi.org/10.1186/s40249-017-0378-z) (2017).
45. Guerra, J. C. *et al.* From climate variables to health information - Predicting and monitoring mosquito-borne disease outbreaks with AeDES2. Tech. Rep. EGU25-15204, Copernicus Meetings (2025). DOI: [10.5194/egusphere-egu25-15204](https://doi.org/10.5194/egusphere-egu25-15204).
46. Leandro, A. & Maciel-de-Freitas, R. Development of an Integrated Surveillance System to Improve Preparedness for Arbovirus Outbreaks in a Dengue Endemic Setting: Descriptive Study. *JMIR Public Heal. Surveillance* **10**, e62759, DOI: [10.2196/62759](https://doi.org/10.2196/62759) (2024).
47. Ropelewski, C. F., Janowiak, J. E. & Halpert, M. S. The Analysis and Display of Real Time Surface Climate Data. *Mon. Weather. Rev.* **113**, 1101–1106, DOI: [10.1175/1520-0493\(1985\)113<1101:TAADOR>2.0.CO;2](https://doi.org/10.1175/1520-0493(1985)113<1101:TAADOR>2.0.CO;2) (1985).
48. Thomson, M. C. *et al.* Malaria early warnings based on seasonal climate forecasts from multi-model ensembles. *Nature* **439**, 576–579, DOI: [10.1038/nature04503](https://doi.org/10.1038/nature04503) (2006).
49. (PDF) Detecting Epidemic Malaria, Uganda. *Research-Gate* DOI: [10.3201/eid1305.061410](https://doi.org/10.3201/eid1305.061410).
50. Moriyama, M., Hugentobler, W. J. & Iwasaki, A. Seasonality of Respiratory Viral Infections. *Annu. Rev. Virol.* **7**, 83–101, DOI: [10.1146/annurev-virology-012420-022445](https://doi.org/10.1146/annurev-virology-012420-022445) (2020).
51. Dietz, K. The estimation of the basic reproduction number for infectious diseases. *Stat. Methods Med. Res.* **2**, 23–41, DOI: [10.1177/096228029300200103](https://doi.org/10.1177/096228029300200103) (1993).
52. WHO guidance for surveillance during an influenza pandemic, 2017 update. <https://www.who.int/publications/i/item/9789241513333>.
53. Chiew, M. *et al.* Estimating the measles effective reproduction number in Australia from routine notification data. *Bull. World Heal. Organ.* **92**, 171–177, DOI: [10.2471/BLT.13.125724](https://doi.org/10.2471/BLT.13.125724) (2014).
54. Thomson, M. C., Muñoz, Á. G., Cousin, R. & Shumake-Guillemot, J. Climate drivers of vector-borne diseases in Africa and their relevance to control programmes. *Infect. Dis. Poverty* **7**, 81, DOI: [10.1186/s40249-018-0460-1](https://doi.org/10.1186/s40249-018-0460-1) (2018).
55. Muñoz, Á. G., Thomson, M. C., Goddard, L. & Aldighieri, S. Analyzing climate variations at multiple timescales can guide Zika virus response measures. *Giga-Science* **5**, 41, DOI: [10.1186/s13742-016-0146-1](https://doi.org/10.1186/s13742-016-0146-1) (2016).
56. Jacoby, W. G. Loess:: A nonparametric, graphical tool for depicting relationships between variables. *Elect. Stud.* **19**, 577–613, DOI: [10.1016/S0261-3794\(99\)00028-1](https://doi.org/10.1016/S0261-3794(99)00028-1) (2000).
57. Craven, P. & Wahba, G. Smoothing noisy data with spline functions. *Numer. Math.* **31**, 377–403, DOI: [10.1007/BF01404567](https://doi.org/10.1007/BF01404567) (1978).

58. Marcotte, D. Generalized cross-validation for covariance model selection. *Math. Geol.* **27**, 659–672, DOI: [10.1007/BF02093906](https://doi.org/10.1007/BF02093906) (1995).
59. Fan, Y. & van den Dool, H. A global monthly land surface air temperature analysis for 1948–present. *J. Geophys. Res. Atmospheres* **113**, DOI: [10.1029/2007JD008470](https://doi.org/10.1029/2007JD008470) (2008).
60. Xie, P., Arkin, P. A. & Janowiak, J. E. CMAP: The CPC Merged Analysis of Precipitation. In Levizzani, V., Bauer, P. & Turk, F. J. (eds.) *Measuring Precipitation From Space: EURAINSAT and the Future*, 319–328, DOI: [10.1007/978-1-4020-5835-6_25](https://doi.org/10.1007/978-1-4020-5835-6_25) (Springer Netherlands, Dordrecht, 2007).
61. Chen, M. *et al.* Assessing objective techniques for gauge-based analyses of global daily precipitation. *J. Geophys. Res. Atmospheres* **113**, DOI: [10.1029/2007JD009132](https://doi.org/10.1029/2007JD009132) (2008).
62. (PDF) Web tool deconstructs variability in twentieth-century climate. *ResearchGate* DOI: [10.1029/2011EO450001](https://doi.org/10.1029/2011EO450001).
63. Spearman, C. The Proof and Measurement of Association between Two Things. *The Am. J. Psychol.* **15**, 72–101, DOI: [10.2307/1412159](https://doi.org/10.2307/1412159) (1904). [1412159](https://doi.org/10.2307/1412159).
64. New, M. & Hulme, M. Representing uncertainty in climate change scenarios: A Monte-Carlo approach. *Integr. Assess.* **1**, 203–213, DOI: [10.1023/A:1019144202120](https://doi.org/10.1023/A:1019144202120) (2000).
65. Liang, X. S. Unraveling the cause-effect relation between time series. *Phys. Rev. E* **90**, 052150, DOI: [10.1103/PhysRevE.90.052150](https://doi.org/10.1103/PhysRevE.90.052150) (2014).
66. Liang, X. S. Normalizing the causality between time series. *Phys. Rev. E* **92**, 022126, DOI: [10.1103/PhysRevE.92.022126](https://doi.org/10.1103/PhysRevE.92.022126) (2015).
67. Beck, H. E. *et al.* High-resolution (1 km) Köppen-Geiger maps for 1901–2099 based on constrained CMIP6 projections. *Sci. Data* **10**, 724, DOI: [10.1038/s41597-023-02549-6](https://doi.org/10.1038/s41597-023-02549-6) (2023).
68. Triantafyllou, G. N. & Tsonis, A. A. Assessing the ability of the Köppen System to delineate the general world pattern of climates. *Geophys. Res. Lett.* **21**, 2809–2812, DOI: [10.1029/94GL01992](https://doi.org/10.1029/94GL01992) (1994).
69. Chen, Y. *et al.* Indian Ocean temperature anomalies predict long-term global dengue trends. *Science* **384**, 639–646, DOI: [10.1126/science.adj4427](https://doi.org/10.1126/science.adj4427) (2024).
70. Kurnianingsih, Wirasatriya, A., Lazuardi, L., Kubota, N. & Ng, N. IOD and ENSO-Related Time Series Variability and Forecasting of Dengue and Malaria Incidence in Indonesia. In *2020 International Symposium on Community-centric Systems (CcS)*, 1–8, DOI: [10.1109/CcS49175.2020.9231358](https://doi.org/10.1109/CcS49175.2020.9231358) (2020).
71. Banu, S. *et al.* Impacts Of El Niño Southern Oscillation And Indian Ocean Dipole On Dengue Incidence In Bangladesh. *ISEE Conf. Abstr.* **2015**, 746, DOI: [10.1289/isee.2015.2015-746](https://doi.org/10.1289/isee.2015.2015-746) (2015).
72. Muñoz, Á. *Cross-Timescale Interference and Rainfall Extreme Events in South Eastern South America*. Ph.D. thesis (2015). DOI: [10.7916/D8G73DFQ](https://doi.org/10.7916/D8G73DFQ).
73. Lowe, R. *et al.* The development of an early warning system for climate-sensitive disease risk with a focus on dengue epidemics in Southeast Brazil. *Stat. Medicine* **32**, 864–883, DOI: [10.1002/sim.5549](https://doi.org/10.1002/sim.5549) (2013).
74. Thomson, M. C., Muñoz, Á. G., Cousin, R. & Shumake-Guillemot, J. Climate drivers of vector-borne diseases in Africa and their relevance to control programmes. *Infect. Dis. Poverty* **7**, 81, DOI: [10.1186/s40249-018-0460-1](https://doi.org/10.1186/s40249-018-0460-1) (2018).
75. Iwamura, T., Guzman-Holst, A. & Murray, K. A. Accelerating invasion potential of disease vector *Aedes aegypti* under climate change. *Nat. Commun.* **11**, 2130, DOI: [10.1038/s41467-020-16010-4](https://doi.org/10.1038/s41467-020-16010-4) (2020).
76. (PDF) Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation. Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change. https://www.researchgate.net/publication/266208219_Managing_the_Risks_of_Extreme_Events_and_Disasters_to_Advance_Climate_Change_Adaptation DOI: [10.13140/2.1.3117.9529](https://doi.org/10.13140/2.1.3117.9529) (2025).
77. Lim, A. *et al.* The overlapping global distribution of dengue, chikungunya, Zika and yellow fever. *Nat. Commun.* **16**, 3418, DOI: [10.1038/s41467-025-58609-5](https://doi.org/10.1038/s41467-025-58609-5) (2025).
78. Laporta, G. Z. *et al.* Global Distribution of *Aedes aegypti* and *Aedes albopictus* in a Climate Change Scenario of Regional Rivalry. *Insects* **14**, 49, DOI: [10.3390/insects14010049](https://doi.org/10.3390/insects14010049) (2023).
79. Ryan, S. J., Carlson, C. J., Mordecai, E. A. & Johnson, L. R. Global expansion and redistribution of *Aedes*-borne virus transmission risk with climate change. *PLOS Neglected Trop. Dis.* **13**, e0007213, DOI: [10.1371/journal.pntd.0007213](https://doi.org/10.1371/journal.pntd.0007213) (2019).
80. Santos, L. L. M., de Aquino, E. C., Fernandes, S. M., Ternes, Y. M. F. & Feres, V. C. d. R. Dengue, chikungunya, and Zika virus infections in Latin America and the Caribbean: A systematic review. *Revista Panam. de Salud Pública* **47**, e34, DOI: [10.26633/RPSP.2023.34](https://doi.org/10.26633/RPSP.2023.34) (2023).
81. Floridi, A. *et al.* Effects of El Niño and the Positive Indian Ocean Dipole (+IOD) on Health, Food Security, Economics, and Conflict in Low- and Middle-Income Countries in the Indo-Pacific: A Systematic Review. *Campbell Syst. Rev.* **21**, e70038, DOI: [10.1002/cl2.70038](https://doi.org/10.1002/cl2.70038) (2025).
82. Mamani, L. *et al.* Extreme droughts in the Amazon Basin during cyclic ENSO events coupled with Indian Ocean Dipole modes and Tropical North Atlantic

warming. *Sci. The Total. Environ.* **963**, 178536, DOI: [10.1016/j.scitotenv.2025.178536](https://doi.org/10.1016/j.scitotenv.2025.178536) (2025).

83. Jia, F., Cai, W., Gan, B., Wu, L. & Di Lorenzo, E. Enhanced North Pacific impact on El Niño/Southern Oscillation under greenhouse warming. *Nat. Clim. Chang.* **11**, 840–847, DOI: [10.1038/s41558-021-01139-x](https://doi.org/10.1038/s41558-021-01139-x) (2021).