Day 2: Population growth, species interactions and time series

Day 2: Population growth, species interactions and time series

Please:

This course is for you!

Please interrupt us as often as needed.

Please give us feedback during and after the course.

Day 2: Population growth, species interactions and time series

Schedule:

09:00 – 10:00	Introduction (Emanuel Fronhofer)
10:15 – 11:15	Fitting population dynamics in stan/rstan (Jhelam Deshpande)
11:30 – 12:30	Introducing process error and state-space models (Benjamin Rosenbaum)
14:00 – 17:00	It’s your turn!
17:00 – 17:30	Fitting multiple time series at once (Jhelam Deshpande)
18:00 – 19:00	Evening lecture: Frédéric Barraquand

Mtui et al. 2017 PloS ONE



Mtui et al. 2017 PloS ONE





Mtui et al. 2017 PloS ONE



Saade et al. 2022 PRSB





Saade et al. 2022 PRSB



Sibly et al. 2005 Science



Global Population Dynamics Database

knb ABOUT DATA SUBMIT TOOLS Jump to: DOI or ID Go SIGN IN

White, Owen Smith, John Lawton, Pablo Inchausti, et al. 2010. The Global Population Dynamics Database. Knowledge Network for Biocomplexity. doi:10.5063/F1BZ63Z8.

knb

Citations 0 Views 2.8K Copy Citation Assessment report

Files in this dataset Package: urn:uuid:bced9df4-1398-4768-a0f3-6d50074464ad4

		File type	Size		
opulation Dynamics Database		EML v2.1.1	48 KB	2763 views	Download
	More info	text/csv	8 MB	2433 downloads	Download
	More info	text/csv	794 KB	1710 downloads	Download
timeperiod.csv	More info	text/csv	17 KB	1532 downloads	Download

Download All

Show 5 more items in this data set

General

Identifier: df35b.240.11

Abstract: As a source of animal and plant population data, the Global Population Dynamics Database (GPDD) is unrivalled. Nearly five thousand separate time series are available here. In addition to all the population counts, there are taxonomic details of over 1400 species. The type of data contained in the GPDD varies enormously, from annual counts of mammals or birds at individual sampling sites, to weekly counts of zooplankton and other marine fauna. The project commenced in October 1994, following discussions on ways in which the collaborating partners could make a practical and enduring contribution to research into population dynamics. A small team was assembled and, with assistance and advice from numerous interested parties we decided to construct the database using the popular Microsoft Access platform. After an initial design phase, the major task has been that of locating, extracting, entering and validating the data in all the various tables. Now, nearly 5000 individual datasets have been entered onto the GPDD. The Global Population Dynamics Database comprises six Tables of data and information. The tables are linked to each other as shown in the diagram shown in figure 3 of the GPDD User Guide (GPDD-User-Guide.pdf). Referential integrity is maintained through record ID numbers which are held, along with other information in the Main Table. It's structure obeys all the rules of a standard relational database.

Keywords

Keyword	Type
Time series data	
animal populations	
plant populations	
count data	

Sibly et al. 2005 Science









Sibly et al. 2005 Science

How should we analyse such data?

How should we analyse such data?



How should we analyse such data?



How should we analyse such data?



How should we analyse such data?



Why is this more interesting than less informed models?

How should we analyse such data?



Why is this more interesting than less informed models?

→ parameter estimation
→ test of theory predictions
→ maybe wrong inference?

Back to basics: exponential growth, logistic growth and inter-specific competition

Back to basics: exponential growth, logistic growth and inter-specific competition

Journal of the History of Biology
https://doi.org/10.1007/s10739-019-09570-9

ORIGINAL RESEARCH

How the Modern Synthesis Came to Ecology

Philippe Huneman[1]

© Springer Nature B.V. 2019

Abstract

Ecology in principle is tied to evolution, since communities and ecosystems result from evolution and ecological conditions determine fitness values (and ultimately evolution by natural selection). Yet the two disciplines of evolution and ecology

Back to basics: exponential growth, logistic growth and inter-specific competition

Evolutionary Ecology Research, 2012, **14**: 627–665

The struggle for existence: how the notion of carrying capacity, *K*, obscures the links between demography, Darwinian evolution, and speciation

James Mallet

Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, Massachusetts, USA and Department of Genetics, Evolution and Environment, University College London, London, UK

Back to basics: exponential growth, logistic growth and inter-specific competition





How should we fit such models?









Sibly et al. 2005 Science







$$\frac{dN}{dt} = r_0 \left[1 - \left(\frac{N}{K} \right)^{\theta} \right] N$$

least-squares



Sibly et al. 2005 Science

Methods in Ecology and Evolution

British Ecological Society

Methods in Ecology and Evolution 2010, **1**, 253–262 doi: 10.1111/j.2041-210X.2010.00029.x

The theta-logistic is unreliable for modelling most census data

Francis Clark[1], Barry W. Brook[1], Steven Delean[1], H. Reşit Akçakaya[2] and Corey J. A. Bradshaw[1,3]*

[1]*The Environment Institute and School of Earth & Environmental Sciences, University of Adelaide, Adelaide, SA 5005, Australia;* [2]*Department of Ecology and Evolution, Stony Brook University, Stony Brook, NY 11794, USA; and* [3]*South Australian Research and Development Institute, P.O. Box 120, Henley Beach, SA 5022, Australia*

Methods in Ecology and Evolution

British Ecological Society

Methods in Ecology and Evolution 2010, **1**, 253–262 doi: 10.1111/j.2041-210X.2010.00029.x

The theta-logistic is unreliable for modelling most census data

Francis Clark[1], Barry W. Brook[1], Steven Delean[1], H. Reşit Akçakaya[2] and Corey J. A. Bradshaw[1,3]*

[1]*The Environment Institute and School of Earth & Environmental Sciences, University of Adelaide, Adelaide, SA 5005, Australia;* [2]*Department of Ecology and Evolution, Stony Brook University, Stony Brook, NY 11794, USA; and* [3]*South Australian Research and Development Institute, P.O. Box 120, Henley Beach, SA 5022, Australia*



Fig. 2. Bimodality of fitted θ and relationship with r_m. (a) Fitted θ for 1000 simulations of 20 time-steps length, with generating model parameters: r_m = 0·5, θ = 1·0, σ = 0·1 and K = 100 (vertical line at generating θ = 1·0). Trace shows the effect of 10% measurement error (ME) applied to the abundance values; (b) $\log_{10}$–$\log_{10}$ scatter plot of fitted [r_m, θ] pairs.

Statistical Rethinking

A Bayesian Course with Examples in R and Stan (& PyMC3 & brms & Julia too, see links below)

Most recent set of free lectures: Statistical Rethinking 2022

Second Edition

The second edition is now out in print. Publisher information on the CRC Press page. For more detail about what is new, look here.

Materials



2nd Edition

- Book: CRC Press
- Book sample: Chapters 1 and 2 (2MB PDF)
- Lectures and slides:
  * Winter 2022 materials (ongoing)
  * Winter 2019 materials
- Code and examples:
  * R package: rethinking (github repository)
  * R code examples from the book: code.txt
  * Book examples in Stan+tidyverse
  * brms + tidyverse conversion here
  * PyMC3 code examples: PyMC repository
  * NumPyro!
  * More NumPyro
  * TensorFlow Probability notebooks
  * Julia & Turing examples (both 1st and 2nd edition)
  * Another Julia code translation with clean outline in notebook format
  * R-INLA examples

FOLLOW ME ON TWITTER

My Tweets

Day 2: Population growth, species interactions and time series

Schedule:

09:00 – 10:00	Introduction (Emanuel Fronhofer)
10:15 – 11:15	Fitting population dynamics in stan/rstan (Jhelam Deshpande)
11:30 – 12:30	Introducing process error and state-space models (Benjamin Rosenbaum)
14:00 – 17:00	It’s your turn!
17:00 – 17:30	Fitting multiple time series at once (Jhelam Deshpande)
18:00 – 19:00	Evening lecture: Frédéric Barraquand

https://github.com/theodatasci/demo_day_2_jhelam