

Biology 30 IB

Genetics

Jad Chehimi

December 3, 2020

Unfinished!

Contents

(18.1) Gregor Mendel	3
Peas	3
Introduction	3
General	3
Terms	3
Dominance	4
Crosses	4
Generations	4
Mendel's Laws	4
(18.2) Punnett Squares	5
(18.3) Pedigree Charts	6
Terms	6
(18.4) Other Patterns of Inheritance	6
Pleiotropy	6
Multiple Alleles	7
Example	7
Incomplete Dominance	7
Codominance	8
Blood Type	8

(18.5) Dihybrid Crosses	9
# of Unique Gametes	9
Drawing a Dihybrid Cross	9
Example	10
Probability	11
Monohybrid Method	11
Combination (Monohybrid to Dihybrid)	11
Calculating Probability	11
Gene Interaction	11
(19.1) Chromosomal Theory	11
Autosomal Traits	12
Sex-linked Genes	12
Carriers in X-linked	12
X-Linked Conditions	13
Testes Determing Factor (TDF)	13
Testicular Feminizing Syndrome (TFS)	13
Mitochondrial DNA	13

(18.1) Gregor Mendel

- Created the **Laws of Inheritance of Traits**
- Studied inheritance of traits in **pea plants** (test question)
- Called DNA and chromosomes particles — he didn't know
- **Father of Genetics**

Peas

- Can be grown in a small area
- Produce lots of offspring
- Self-pollinate
- Can be **artificially cross-pollinated** (test question)

Introduction

General

- **Trait** = any **characteristic** that can be passed from **parent to offspring**
- **Heredity** = passing of traits from parent to offspring
- **Genetics** = study of heredity

Terms

- **Alleles** = two forms of a gene — **dominant & recessive**
- **Dominant** = **stronger** of two genes, represented with **capital letter** (R)
- **Recessive** = **weaker** of two genes, represented with **lowercase letter** (r)
- **Genotype** = gene combination for a trait (**RR, Rr, rr**) separated with colons
- **Phenotype** = **physical feature** resulting from genotype (red, white) separated with colons
- **Homozygous genotype** = genotype involving **2 dominant** OR **2 recessive** genes — **pure**, RR or rr
- **Heterozygous genotype** = genotype involving **1 dominant** OR **1 recessive** gene — **hybrid**, Rr

Dominance

- Dominant and recessive alleles can code for different things
- e.x. R = Brown eyes, r = blue eyes
- Any dominant alleles = dominant phenotype — RR, Rr in most cases
- Only recessive alleles = recessive phenotype — rr

Genotypes	RR	Rr	rr
Phenotypes	RED	RED	YELLOW

Crosses

- **Monohybrid cross** = cross involving a single trait
- **Dihybrid cross** = cross involving two traits
- **Test cross** = cross involving always involving a homozygous recessive (rr) parent crossed with an unknown genotype, in order to find the genotype

Generations

- **P1 Generation** = parental generation in a breeding experiment
- **F1 Generation** (1st filial gen) = first-generation offspring in a breeding experiment
- **F2 Generation** (2nd filial gen) = second-generation offspring, and so on...

Mendel's Laws

LAW	PARENT CROSS	OFFSPRING
DOMINANCE	TT x tt tall x short	100% Tt tall
SEGREGATION	Tt x Tt tall x tall	75% tall 25% short
INDEPENDENT ASSORTMENT	RrGg x RrGg round & green x round & green	9/16 round seeds & green pods 3/16 round seeds & yellow pods 3/16 wrinkled seeds & green pods 1/16 wrinkled seeds & yellow pods

(18.2) Punnett Squares




Figure 1: Punnett square demonstrating **law of segregation**, which occurs in **anaphase I**

- One gene from father, one gene from mother
- Dominant and recessive alleles compete in punnett squares
- **Phenotypic ratio** = ratio of **dominant phenotypes to recessive phenotypes**, in this diagram its 3:1

(18.3) Pedigree Charts




Figure 2: All in your data booklet, except known heterozygotes

- **Pedigree** = a family tree used to trace inherited traits from parents to offspring
- Roman numerals denote different generations (rows)
- Arabic numerals denote different individuals in a generation (columns)

Terms

- **Autosomal condition** = almost equal number of males and females are affected by a condition
- **Dominance condition** = condition appears in every generation
- **Recessive condition** = condition does not appear in every generation; skips generations

(18.4) Other Patterns of Inheritance

Pleiotropy

- **Pleiotropic gene** = one gene affects more than one phenotypic characteristic
- Examples of wide-ranging effects from a single gene include...

- dwarfism (achondroplasia)
- gigantism (acromegaly)

Multiple Alleles

- Possible to have more than two alleles for a trait
- Superscript identifies different alleles
- Different alleles are dominant to one another
 - e.g. $E^1 > E^2 > E^3 > E^4$


Example

Predict the genotypic and phenotypic outcomes of crossing $E^1E^4 \times E^2E^3$.

- E^1 = wild, E^2 = apricot, E^3 = honey, E^4 = white

Wild type	$E^1E^1, E^1E^2, E^1E^3, E^1E^4$
Apricot	E^2E^2, E^2E^3, E^2E^4
Honey	E^3E^3, E^3E^4
White	E^4E^4

Figure 3: This is typically given



- Genotype = $1 \times E^1E^2 : 1 \times E^1E^3 : 1 \times E^2E^4 : 1 \times E^3E^4$
- Phenotype = 2 wild : 1 apricot : 1 honey

Incomplete Dominance

- A hybrid (Rr) appearance can sometimes be in between the phenotypes
- For example...

- RR = red
- rr = white
- Rr = pink (normally red)

Codominance

- Both alleles expressed in heterozygous individuals



Figure 4: Red bull + white cow = red and white hair

Blood Type

Example of codominance and multiple alleles

$$I^A \text{ \& } I^B > i$$

- Type A = $I^A I^A$ or $I^A i$
- Type B = $I^B I^B$ or $I^B i$
- Type AB = $I^A I^B$
- Type O = ii

(18.5) Dihybrid Crosses

- A breeding experiment that tracks the inheritance of **two traits**
- **Law of Independent Assortment** = each pair of alleles segregate independently (metaphase)
- The two traits **do not influence one another**, they are independent
- These traits are on different locations — called **loci**, **locus** — of a chromosome

of Unique Gametes

$$2^n$$

n = # of heterozygotes of **each parent**

Examples...

- RrBb = 2 heterozygotes = $2^2 = 4$ gametes
- yypp = 0 heterozygotes = $2^0 = 1$ gamete
- AaBbCCDd = 3 heterozygotes = $2^3 = 8$ gametes
- MmNnOoPPQQRrssTtXx = 6 heterozygotes = $2^6 = 64$ gametes

Drawing a Dihybrid Cross

Determine how many different possible gametes/phenotypes with the formula.

- $2^2 = 4$
- There will be 4 different types of offspring from this cross
- If a question specifically asks for the **adults**, there may be a gene that prevents adulthood — i.e. a fatal gene — so do not include that combination in any percent calculations and ratios
- If you are doing a test cross — there is an unknown gene — make a dihybrid cross for each possible combination; this applies to monohybrid as well

Prepare a cross for every possible combination of all heterozygotes.

1. Top header = **FOIL** between genes of the gene pair of the first parent
2. Left header = **FOIL** between genes of the gene pair of other parent
3. You can skip redundant combinations
4. Fill in the table like usual, including the top and left header, to list every possible combination
5. Make sure to order the alleles dominant to recessive, and same letters together

Using the cross, you can...

- Determine the number/ratio of all possible genotypes by counting
- Determine the number of phenotypes from all possible genotypes

It helps to mark each of the table cells with a symbol to keep track.

Example

- $R_ = \text{round}$, $Y_ = \text{yellow}$
- $rr = \text{wrinkled}$, $yy = \text{green}$

Question: Determine the genotype and phenotype of a $RyYy \times RyYy$ cross.

	RY	Ry	rY	ry
RY	RRYY	RRYy	RrYY	RrYy
Ry	RRYy	RRyy	RrYy	Rryy
rY	RrYY	RrYy	rrYY	rrYy
ry	RrYy	Rryy	rrYy	rryy

- Genotype
 $1 \times RRYY : 2 \times RRYy : 2 \times RrYY : 4 \times RrYy : 1 \times RRyy : 2 \times Rryy : 1 \times rrYY : 2 \times rrYy : 1 \times rryy$
- Phenotype
 - 9 round, yellow
 - 3 round, green
 - 3 wrinkled, yellow
 - 1 wrinkled, green

Probability

$$\text{probability} = \frac{\# \text{ of chances of an event}}{\# \text{ of possible combinations}}$$

- **Independent event** = individual event, random chance, previous events do not influence

Monohybrid Method

Alternative to creating a dihybrid cross.

1. Create a monohybrid cross for each of only the **matching letter genes between both parents**

e.g. $MmNnOo \times MmNnOo = Mn \times Mn, Nn \times Nn, Oo \times Oo$

Combination (Monohybrid to Dihybrid)

- Similar to distributive property
- Multiply **each gene in the first** monohybrid by **every gene in the other** monohybrid
- This should result in the same answers as FOILing in a dihybrid cross

Calculating Probability

- Divide the **number of a desired gene** out of the **number of possible combinations**
- This applies to **both methods** — the entire dihybrid FOIL, and the monohybrids of the monohybrid method
- **Multiply all probabilities** together if looking for the probability of multiple specific independent genes together
e.g. the probability of blue eyes and brown hair

Gene Interaction

- **Polygenic** = more than one genes influence a trait
e.g. skin colour, eye colour, height
- **Epistasis** = one gene masks/epistatic to another, influences ratio

(19.1) Chromosomal Theory

- **Chromosomal Theory** = anything to do with chromosomes (diploma question)
- Sex-linked genes discovered by **Morgan**

Autosomal Traits

- **Cystic fibrosis** = mucus build up in many organs
 - normal alleles = Cc, CC
 - mutated alleles = cc
- **Sickle cell anemia** = sickling of blood cells, cannot carry oxygen
 - normal alleles = Hb^AHb^A
 - mutated alleles = Hb^SHb^S
 - **carrier alleles** = Hb^AHb^S (less extreme effects)

Sex-linked Genes

- **Autosomes** = chromosomes that do not code for proteins related to the sex of the individual
- **Autosomal trait** = traits not located on sex chromosomes
- **Sex-linked trait** = traits located on sex chromosomes

Carriers in X-linked

- Females cannot be affected by an irregular allele, since they have two X chromosomes in which one of them may dominate the other; they **can be carriers** — they carry a dormant recessive irregular allele, which can be given to their offspring (homozygous recessive females are rare)
 - **Barr body** = dormant, **unexpressed X** chromosome; chromosome shrinks into barr body; **not in males**, somatic cells only
 - **Barr bodies** = whichever allele is dormant is not consistent in all body cells; causes spots of dark skin and no sweat glands
- Males can be affected by an irregular allele, but **cannot be carriers**; they only have one X chromosome, which could be the irregular one from their mom
- A mother's recessive alleles are **most likely** to be given to their **sons**

X-Linked Conditions

- At this level, X chromosomes carry traits, while Y chromosomes do not
- **Hemophilia** = blood clotting disorder
 - normal allele = X^H
 - mutated allele = X^h
 - X^hY , X^hX^h
- **Colour-blindness** = typically red-green colourblind
 - normal allele = X^R
 - mutated allele = X^r
 - X^rY , X^rX^r

Testes Determining Factor (TDF)

- Gene located on the Y chromosome
- Develops male gonads while in embryo
- Produces testosterone when activated

Testicular Feminizing Syndrome (TFS)

- Gonad forming tissues ignore testosterone
- Produces female vaginal folds AND penis

Mitochondrial DNA

- From mother
- Can be defective, causing deafness