

Table 1: Heterogeneity of stacked GATE effects: diarrhea

Heterogeneity group	Any Treatment		Boiling		Chlorination/tablets		Straining/settling	
	Clustered folds	Ordinary folds	Clustered folds	Ordinary folds	Clustered folds	Ordinary folds	Clustered folds	Ordinary folds
Panel A: Diarrhea (under-5): Initial source-water E. coli decile								
Decile 1	0.031 (0.031)	0.044 (0.028)	0.027 (0.070)	0.048 (0.083)	-0.002 (0.019)	-0.008 (0.043)	0.084*** (0.021)	0.091*** (0.026)
Decile 5	0.038 (0.045)	0.035 (0.051)	0.065 (0.090)	0.035 (0.094)	0.019 (0.027)	0.083 (0.080)	0.132 (0.131)	0.075 (0.105)
Decile 6	0.028 (0.064)	0.024 (0.061)	0.085* (0.049)	0.087** (0.044)	-0.331 (0.276)	-0.162 (0.234)	-0.040 (0.212)	-0.094 (0.236)
Decile 7	-0.125 (0.114)	-0.089 (0.081)	-0.019 (0.069)	-0.026 (0.079)	0.001 (0.073)	-0.022 (0.050)	-0.378 (0.366)	-0.236 (0.277)
Decile 8	-0.013 (0.039)	-0.033 (0.044)	-0.060 (0.084)	-0.072 (0.090)	0.095** (0.039)	0.044 (0.045)	0.003 (0.086)	-0.013 (0.065)
<i>Sample</i>								
IRM	$N = 3,612$; PSUs = 3,263							
APOS	$N = 3,612$; PSUs = 3,263							
<i>BLP R²</i>								
IRM, Any Treatment	0.001	0.001						
APOS, Boiling			0.000	0.000				
APOS, Chlorination/tablets					0.004	0.001		
APOS, Straining/settling							0.002	0.001

Notes: Coefficient rows report GATE estimates with significance stars; the following row reports standard errors in parentheses. Clustered folds use cluster-level sample splitting; ordinary folds use unclustered folds. N and PSU counts are reported for each outcome and method. $BLP R^2$ is descriptive: it is the R-squared from projecting the APOS/IRM orthogonal contrast signal onto heterogeneity-group indicators, not a causal-model R^2 . The heterogeneity analysis is exploratory.