

Table 1: Stacked water-treatment effects: clustered and ordinary folds

	Some risk		Very high risk		Diarrhea (U5)	
	Clustered	Unclustered	Clustered	Unclustered	Clustered	Unclustered
<i>IRM</i>						
Any treatment	-0.080*** (0.006)	-0.080*** (0.006)	-0.064*** (0.007)	-0.065*** (0.006)	-0.000 (0.008)	0.000 (0.008)
Outcome mean	0.737		0.308		0.139	
Treated (%)	21.3%		21.3%		19.5%	
<i>APOS</i>						
Boiling	-0.123** (0.058)	-0.123*** (0.008)	-0.103* (0.060)	-0.102*** (0.008)	-0.003 (0.058)	-0.002 (0.009)
Chlorination/tablets	-0.046 (0.066)	-0.043*** (0.009)	-0.071 (0.074)	-0.075*** (0.010)	-0.009 (0.073)	-0.009 (0.011)
Straining/settling	-0.016 (0.065)	-0.018** (0.009)	-0.012 (0.089)	-0.016 (0.011)	0.027 (0.075)	0.027** (0.011)
Outcome mean	0.737		0.308		0.139	
Shares: $D = 0$	78.8%		78.8%		80.6%	
Shares: $D = 1$	10.1%		10.1%		7.6%	
Shares: $D = 2$	2.2%		2.2%		2.0%	
Shares: $D = 3$	5.1%		5.1%		7.2%	
Shares: $D = 98$	3.9%		3.9%		2.6%	
<i>Sample</i>						
Observations	59,620		59,620		36,121	
PSUs	18,655		18,655		13,596	

Notes: Cells report coefficients with significance stars and standard errors in parentheses. IRM and APOS are separated into blocks. Clustered specifications use cluster-level sample splitting; ordinary specifications use unclustered folds. APOS reports treatment shares for all estimated treatment levels, including level 98. Cross-fitting uses 5 folds and 3 repetitions. APOS effects are contrasts relative to treatment level 0. *** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$.